



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 08:55 PM UTC

PDB ID : 4BEL / pdb_00004bel
Title : BACE2 XAPERONE COMPLEX
Authors : Banner, D.W.; Kuglstatter, A.; Benz, J.; Stihle, M.; Ruf, A.
Deposited on : 2013-03-11
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	6% 85% 8% 7%
1	B	386	5% 83% 9% 8%
2	D	122	2% 86% 5% 8%
2	E	122	84% 8% 7%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-SECRETASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	6	0
			2821	1816	452	541	12			
1	B	357	Total	C	N	O	S	0	9	0
			2834	1821	454	547	12			


- Molecule 2 is a protein called XA4813.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	112	Total	C	N	O	S	0	4	0
			860	527	155	171	7			
2	E	113	Total	C	N	O	S	0	6	0
			887	544	163	174	6			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is 2-[3-(2-HYDROXY-1,1-DIHYDROXYMETHYL-ETHYLAMINO)-PROPYLAMINO]-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: B3P) (formula: C₁₁H₂₆N₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			19	11	2	6		
4	B	1	Total	C	N	O	0	0
			19	11	2	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	210	Total	O	0	0
			210	210		
5	B	212	Total	O	0	0
			212	212		
5	D	96	Total	O	0	0
			96	96		
5	E	100	Total	O	0	0
			100	100		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: BETA-SECRETASE 2




• Molecule 1: BETA-SECRETASE 2



• Molecule 2: XA4813



• Molecule 2: XA4813



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.03Å 61.76Å 115.62Å 90.00° 101.38° 90.00°	Depositor
Resolution (Å)	43.68 – 1.85 43.68 – 1.85	Depositor EDS
% Data completeness (in resolution range)	90.0 (43.68-1.85) 90.0 (43.68-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.181 , 0.222 0.183 , 0.224	Depositor DCC
R_{free} test set	3757 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8059	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 19.97 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.6362e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, B3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/2889	0.82	0/3925
1	B	0.62	0/2901	0.83	2/3943 (0.1%)
2	D	0.70	0/874	0.83	0/1182
2	E	0.65	0/905	0.78	0/1223
All	All	0.63	0/7569	0.82	2/10273 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	228	LEU	N-CA-C	6.42	120.51	110.17
1	B	290	TRP	N-CA-C	5.72	118.25	111.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	227	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2821	0	2734	24	0
1	B	2834	0	2742	20	0
2	D	860	0	835	8	0
2	E	887	0	869	8	0
3	A	1	0	0	0	0
4	A	19	0	26	2	0
4	B	19	0	26	1	0
5	A	210	0	0	2	0
5	B	212	0	0	4	0
5	D	96	0	0	3	0
5	E	100	0	0	5	0
All	All	8059	0	7232	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:TYR:CE1	1:A:170:MET:HE3	2.11	0.84
2:E:236:ASN:CG	5:E:2080:HOH:O	2.19	0.84
2:D:271:SER:C	5:D:2016:HOH:O	2.22	0.82
1:B:141[B]:THR:HG23	5:B:2091:HOH:O	1.82	0.78
1:A:170:MET:HE1	1:A:347:ALA:CB	2.14	0.77
1:A:344:VAL:HG21	4:A:1399:B3P:H72	1.71	0.71
2:D:242[B]:MET:HE2	2:D:245:LEU:HD21	1.73	0.70
1:A:34:MET:HE1	1:A:54:PHE:CZ	2.27	0.69
1:A:170:MET:HE2	1:A:351:GLU:HG3	1.75	0.69
1:A:170:MET:HE1	1:A:347:ALA:HB1	1.76	0.67
2:E:178[B]:ARG:NH2	5:E:2027:HOH:O	2.29	0.65
1:B:320:GLN:OE1	1:B:333[A]:ARG:NE	2.22	0.65
2:E:198:GLN:NE2	2:E:202:LYS:O	2.32	0.61
1:B:129:ILE:HD12	1:B:131:TRP:O	2.02	0.59
1:A:34:MET:HE3	1:A:101:VAL:HG11	1.85	0.59
1:A:30:TYR:HE1	1:A:170:MET:HE3	1.66	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:SER:HA	1:B:351:GLU:OE2	2.03	0.58
1:A:25:ASP:HB2	5:A:2005:HOH:O	2.04	0.57
1:A:197:TYR:HA	1:A:370:PRO:HD3	1.86	0.56
1:A:111[B]:PHE:CZ	1:A:156:LEU:HD13	2.43	0.54
2:D:271:SER:HB2	5:D:2096:HOH:O	2.08	0.54
5:A:2112:HOH:O	2:D:257:SER:OG	2.19	0.50
2:E:242[B]:MET:HE2	2:E:245:LEU:HD21	1.93	0.50
1:B:61:HIS:HB3	1:B:64:ILE:HG12	1.94	0.49
1:A:158:THR:HG22	2:D:194:THR:HG21	1.94	0.49
1:B:259:GLU:HG3	1:B:263:ARG:HH11	1.78	0.49
1:A:126:LEU:O	1:A:129:ILE:HG12	2.13	0.48
1:A:170:MET:HE1	1:A:347:ALA:CA	2.44	0.48
1:A:30:TYR:CE1	1:A:170:MET:CE	2.93	0.48
1:A:139:TYR:CZ	1:A:209:GLU:HG2	2.49	0.47
1:B:58:GLY:HA3	1:B:117:THR:HG22	1.97	0.47
1:B:210:TRP:CG	1:B:211:TYR:H	2.32	0.47
1:B:64:ILE:HD12	1:B:66[B]:THR:O	2.14	0.47
1:A:246:LEU:HD22	1:A:335:GLY:HA2	1.96	0.46
1:B:362:LYS:NZ	5:B:2145:HOH:O	2.49	0.46
1:B:109[A]:THR:HG22	5:B:2075:HOH:O	2.16	0.45
1:A:170:MET:CE	1:A:347:ALA:HB1	2.44	0.45
2:E:164:GLN:NE2	5:E:2007:HOH:O	2.48	0.45
1:B:35:LEU:HB3	1:B:40:PRO:HB2	1.99	0.44
1:A:26:SER:HB3	1:A:351:GLU:OE2	2.17	0.44
1:A:158:THR:CG2	2:D:194:THR:HG21	2.48	0.44
2:E:236:ASN:OD1	5:E:2080:HOH:O	2.20	0.44
2:D:242[A]:MET:HE1	2:D:267:VAL:HG21	2.00	0.43
1:B:35:LEU:HB2	1:B:102:THR:HB	2.00	0.43
4:A:1399:B3P:H61	4:A:1399:B3P:H31	1.66	0.42
1:A:111[A]:PHE:CE1	1:A:160:ALA:HB2	2.55	0.42
1:A:210:TRP:CG	1:A:211:TYR:H	2.38	0.42
1:A:235:ALA:HA	1:A:236:ASP:HA	1.68	0.41
1:B:158[A]:THR:CG2	2:E:194:THR:HG21	2.50	0.41
1:B:163:PRO:HD2	1:B:190:GLY:O	2.20	0.41
1:B:288:THR:HB	1:B:289:PRO:HD2	2.02	0.41
2:D:202:LYS:HE2	5:D:2004:HOH:O	2.20	0.41
1:B:197:TYR:HA	1:B:370:PRO:HD3	2.03	0.41
1:B:34:MET:HE2	1:B:101:VAL:HG11	2.02	0.40
1:B:78:SER:HB2	5:B:2052:HOH:O	2.20	0.40
1:B:87:TYR:CZ	4:B:1398:B3P:H112	2.57	0.40
1:A:111[A]:PHE:CZ	1:A:160:ALA:HB2	2.57	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:236:ASN:CB	5:E:2080:HOH:O	2.65	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/386 (92%)	350 (99%)	4 (1%)	0	100	100
1	B	356/386 (92%)	351 (99%)	5 (1%)	0	100	100
2	D	114/122 (93%)	111 (97%)	3 (3%)	0	100	100
2	E	117/122 (96%)	116 (99%)	1 (1%)	0	100	100
All	All	941/1016 (93%)	928 (99%)	13 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/320 (96%)	300 (98%)	6 (2%)	48	36
1	B	308/320 (96%)	304 (99%)	4 (1%)	61	51
2	D	94/99 (95%)	93 (99%)	1 (1%)	65	57
2	E	97/99 (98%)	95 (98%)	2 (2%)	47	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	805/838 (96%)	792 (98%)	13 (2%)	59 44

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	210	TRP
1	A	224	GLN
1	A	284	THR
1	A	333[A]	ARG
1	A	333[B]	ARG
1	A	393	SER
1	B	109[A]	THR
1	B	109[B]	THR
1	B	236	ASP
1	B	252	LYS
2	D	271	SER
2	E	221	ASP
2	E	224	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	89	GLN
1	B	169	GLN
1	B	227	ASN
1	B	341	ASN
2	D	164	GLN
2	D	241	GLN
2	D	243	ASN
2	E	164	GLN
2	E	241	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	B3P	B	1398	-	18,18,18	0.55	0	23,23,23	2.07	4 (17%)
4	B3P	A	1399	-	18,18,18	0.52	0	23,23,23	1.38	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	B3P	B	1398	-	-	13/28/28/28	-
4	B3P	A	1399	-	-	17/28/28/28	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1398	B3P	C3-N1-C4	7.27	126.80	116.17
4	B	1398	B3P	C2-N2-C8	5.45	124.13	116.17
4	A	1399	B3P	C3-N1-C4	4.22	122.34	116.17
4	A	1399	B3P	C2-N2-C8	3.75	121.65	116.17
4	B	1398	B3P	O2-C10-C8	-2.23	107.14	111.68
4	B	1398	B3P	C6-C4-C5	-2.14	105.34	110.02

There are no chirality outliers.

All (30) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1399	B3P	C5-C4-N1-C3
4	A	1399	B3P	C6-C4-N1-C3
4	A	1399	B3P	N1-C4-C5-O4
4	A	1399	B3P	C6-C4-C5-O4
4	A	1399	B3P	C7-C4-C5-O4
4	A	1399	B3P	N1-C4-C6-O5
4	A	1399	B3P	C5-C4-C6-O5
4	A	1399	B3P	C7-C4-C6-O5
4	A	1399	B3P	N1-C4-C7-O6
4	A	1399	B3P	C5-C4-C7-O6
4	A	1399	B3P	C6-C4-C7-O6
4	B	1398	B3P	C1-C3-N1-C4
4	B	1398	B3P	C5-C4-N1-C3
4	B	1398	B3P	C6-C4-N1-C3
4	B	1398	B3P	C7-C4-N1-C3
4	B	1398	B3P	N2-C8-C9-O1
4	B	1398	B3P	C11-C8-C9-O1
4	A	1399	B3P	C3-C1-C2-N2
4	B	1398	B3P	C10-C8-C9-O1
4	B	1398	B3P	C2-C1-C3-N1
4	B	1398	B3P	C3-C1-C2-N2
4	B	1398	B3P	N1-C4-C6-O5
4	B	1398	B3P	C7-C4-C6-O5
4	B	1398	B3P	C1-C2-N2-C8
4	A	1399	B3P	C7-C4-N1-C3
4	A	1399	B3P	C1-C3-N1-C4
4	A	1399	B3P	O3-C11-C8-C10
4	B	1398	B3P	C5-C4-C6-O5
4	A	1399	B3P	O3-C11-C8-N2
4	A	1399	B3P	N2-C8-C9-O1


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1398	B3P	1	0
4	A	1399	B3P	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	358/386 (92%)	0.18	23 (6%)	25	27	11, 28, 56, 78	26 (7%)
1	B	357/386 (92%)	0.08	19 (5%)	32	35	11, 26, 53, 76	29 (8%)
2	D	112/122 (91%)	-0.32	2 (1%)	67	71	11, 23, 35, 57	8 (7%)
2	E	113/122 (92%)	-0.32	0	100	100	11, 24, 38, 53	8 (7%)
All	All	940/1016 (92%)	0.02	44 (4%)	36	39	11, 25, 52, 78	71 (7%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	265	SER	7.9
1	B	26	SER	6.2
1	B	27	GLY	5.9
1	A	292	TYR	5.8
1	A	271	SER	5.1
1	A	283	TRP	4.6
1	B	88	THR	4.5
1	B	90	GLY	4.3
1	B	283	TRP	4.2
1	B	227	ASN	3.9
1	A	28	ARG	3.8
1	A	284	THR	3.7
2	D	271	SER	3.4
1	B	28	ARG	3.4
1	A	88	THR	3.4
1	A	289	PRO	3.3
1	A	331	CYS	3.2
1	A	264	ALA	3.0
1	A	281	ALA	2.8
1	A	16	LEU	2.7
1	B	89	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	322	MET	2.6
1	A	276	THR	2.6
1	A	87	TYR	2.6
1	A	275	TRP	2.6
1	B	292	TYR	2.5
1	A	86	LYS	2.5
1	B	174	GLY	2.4
1	A	89	GLN	2.3
1	B	14	ASN	2.3
2	D	270	SER	2.3
1	B	275	TRP	2.3
1	B	288	THR	2.3
1	B	262	ALA	2.3
1	A	397	PRO	2.2
1	A	26	SER	2.2
1	A	104	PRO	2.2
1	B	16	LEU	2.2
1	B	282	CYS	2.1
1	B	280	LEU	2.1
1	B	281	ALA	2.1
1	A	280	LEU	2.1
1	B	331	CYS	2.1
1	A	227	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	B3P	A	1399	19/19	0.78	0.18	41,45,47,47	19
4	B3P	B	1398	19/19	0.84	0.12	42,50,55,66	0
3	CL	A	1398	1/1	0.99	0.03	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around B3P A 1399:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.