



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:57 PM UTC

PDB ID : 7BEB / pdb_00007beb
Title : Unusual structural features in the adduct of dirhodium tetraacetate with lysozyme (4)
Authors : Loreto, D.; Ferraro, G.; Merlino, A.
Deposited on : 2020-12-23
Resolution : 1.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2531 (1.34-1.30)
Clashscore	190562	2585 (1.34-1.30)
Ramachandran outliers	187476	2528 (1.34-1.30)
Sidechain outliers	187428	2528 (1.34-1.30)
RSRZ outliers	180081	2528 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	129	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NO3	AAA	204	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	ACT	AAA	218	-	-	X	-
6	ACT	AAA	223	-	-	X	-
6	ACT	AAA	226	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 1275 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	129	1053	644	205	194	10	0	7	0

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	2	Total	Na	0	0
			2	2		

- Molecule 3 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	1
			8	2	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		
3	AAA	1	Total	N	O	0	0
			4	1	3		

- Molecule 4 is Rhodium (CCD ID: RH) (formula: Rh) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	2	Total	Rh	0	1
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	AAA	1	Total	C	O	0	0
			4	2	2		
5	AAA	1	Total	C	O	0	0
			4	2	2		
5	AAA	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	AAA	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0
6	AAA	1	Total C O 4 2 2	0	0


- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	AAA	120	Total O 122 122	0	2

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysozyme

Chain AAA: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.60Å 78.60Å 37.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.58 – 1.32 55.58 – 1.32	Depositor EDS
% Data completeness (in resolution range)	95.4 (55.58-1.32) 95.4 (55.58-1.32)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 1.32Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.178 , 0.223 0.187 , 0.228	Depositor DCC
R_{free} test set	1286 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtrriage
Anisotropy	0.001	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1275	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RH, EDO, NO3, ACT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	1.06	0/1079	1.18	0/1455

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1053	0	1008	23	1
2	AAA	2	0	0	0	0
3	AAA	52	0	0	3	0
4	AAA	2	0	0	0	0
5	AAA	12	0	16	1	0
6	AAA	32	0	24	10	0
7	AAA	122	0	0	5	0
All	All	1275	0	1048	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:13:LYS:NZ	7:AAA:301:HOH:O	2.07	0.80
1:AAA:116:LYS:NZ	6:AAA:226:ACT:H1	2.03	0.74
1:AAA:74:ASN:ND2	3:AAA:207:NO3:O2	2.33	0.61
1:AAA:116:LYS:HZ3	6:AAA:226:ACT:H1	1.66	0.59
1:AAA:46:ASN:OD1	1:AAA:59[A]:ASN:ND2	2.38	0.57

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:129:LEU:O	1:AAA:129:LEU:O[8_555]	1.93	0.27

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	133/129 (103%)	132 (99%)	1 (1%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	112/105 (107%)	110 (98%)	2 (2%)	51 15

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	55	ILE
1	AAA	125	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 4 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ACT	AAA	218	-	3,3,3	2.14	1 (33%)	3,3,3	0.92	0
6	ACT	AAA	221	-	3,3,3	0.90	0	3,3,3	0.76	0
3	NO3	AAA	203[B]	-	1,3,3	0.56	0	0,3,3	-	-
6	ACT	AAA	219	-	3,3,3	1.00	0	3,3,3	0.87	0
6	ACT	AAA	220	4	3,3,3	0.96	0	3,3,3	0.81	0
6	ACT	AAA	216	4	3,3,3	1.01	0	3,3,3	0.74	0
3	NO3	AAA	210	-	1,3,3	0.14	0	0,3,3	-	-
6	ACT	AAA	226	-	3,3,3	1.02	0	3,3,3	0.72	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	AAA	212	2	3,3,3	0.32	0	2,2,2	0.38	0
3	NO3	AAA	205	-	1,3,3	1.03	0	0,3,3	-	-
3	NO3	AAA	222	-	1,3,3	0.78	0	0,3,3	-	-
3	NO3	AAA	204	-	1,3,3	1.25	0	0,3,3	-	-
3	NO3	AAA	225	-	1,3,3	0.51	0	0,3,3	-	-
3	NO3	AAA	206	-	1,3,3	0.35	0	0,3,3	-	-
5	EDO	AAA	213	-	3,3,3	0.21	0	2,2,2	0.36	0
5	EDO	AAA	214	-	3,3,3	0.08	0	2,2,2	0.16	0
3	NO3	AAA	203[C]	-	1,3,3	1.98	0	0,3,3	-	-
3	NO3	AAA	209	-	1,3,3	0.67	0	0,3,3	-	-
3	NO3	AAA	202	2	1,3,3	0.26	0	0,3,3	-	-
3	NO3	AAA	207	-	1,3,3	0.70	0	0,3,3	-	-
6	ACT	AAA	223	2	3,3,3	0.89	0	3,3,3	0.89	0
3	NO3	AAA	224	-	1,3,3	0.23	0	0,3,3	-	-
3	NO3	AAA	208	-	1,3,3	0.14	0	0,3,3	-	-
6	ACT	AAA	217	-	3,3,3	0.90	0	3,3,3	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	AAA	212	2	-	0/1/1/1	-
5	EDO	AAA	213	-	-	1/1/1/1	-
5	EDO	AAA	214	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	AAA	218	ACT	OXT-C	-3.52	1.14	1.30

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	AAA	213	EDO	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	AAA	218	ACT	5	0
6	AAA	221	ACT	1	0
6	AAA	219	ACT	1	0
6	AAA	226	ACT	3	0
5	AAA	212	EDO	1	0
3	AAA	204	NO3	2	0
3	AAA	207	NO3	1	0
6	AAA	223	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	129/129 (100%)	0.78	8 (6%) 26 34	11, 25, 40, 53	7 (5%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	71	GLY	3.3
1	AAA	126	GLY	3.0
1	AAA	121	GLN	2.7
1	AAA	129	LEU	2.6
1	AAA	62	TRP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NO3	AAA	204	4/4	0.65	0.16	30,31,34,46	0
6	ACT	AAA	219	4/4	0.65	0.21	56,63,66,67	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ACT	AAA	223	4/4	0.70	0.23	45,60,62,68	0
5	EDO	AAA	213	4/4	0.74	0.16	54,57,59,59	0
6	ACT	AAA	226	4/4	0.76	0.15	31,47,51,56	0
6	ACT	AAA	220	4/4	0.79	0.20	51,51,52,54	4
3	NO3	AAA	209	4/4	0.79	0.12	33,45,46,47	0
3	NO3	AAA	210	4/4	0.79	0.11	50,63,65,69	0
3	NO3	AAA	208	4/4	0.80	0.26	22,31,35,42	4
6	ACT	AAA	217	4/4	0.81	0.18	57,63,64,67	0
6	ACT	AAA	216	4/4	0.83	0.14	42,44,53,57	0
6	ACT	AAA	218	4/4	0.83	0.14	24,34,37,38	0
5	EDO	AAA	212	4/4	0.84	0.12	31,32,33,36	0
3	NO3	AAA	206	4/4	0.85	0.09	37,43,44,55	0
5	EDO	AAA	214	4/4	0.86	0.13	47,53,53,55	0
3	NO3	AAA	205	4/4	0.86	0.10	37,38,39,40	0
6	ACT	AAA	221	4/4	0.87	0.17	36,43,46,49	0
3	NO3	AAA	203[C]	4/4	0.88	0.13	22,30,32,38	4
3	NO3	AAA	203[B]	4/4	0.88	0.13	24,31,33,36	4
3	NO3	AAA	225	4/4	0.89	0.11	58,59,60,61	4
3	NO3	AAA	224	4/4	0.89	0.14	30,31,37,47	4
3	NO3	AAA	207	4/4	0.90	0.10	40,50,53,64	0
4	RH	AAA	215[B]	1/1	0.92	0.10	41,41,41,41	1
3	NO3	AAA	222	4/4	0.92	0.08	24,27,28,33	0
2	NA	AAA	227	1/1	0.93	0.18	32,32,32,32	0
3	NO3	AAA	202	4/4	0.93	0.07	27,30,32,33	0
2	NA	AAA	201	1/1	0.94	0.09	25,25,25,25	0
4	RH	AAA	211	1/1	0.96	0.06	35,35,35,35	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around RH AAA 215 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.