



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:24 PM UTC

PDB ID : 9BEZ / pdb_00009bez
Title : MID domain of human Argo2 bound to RNA
Authors : Harp, J.M.; Egli, M.
Deposited on : 2024-04-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	4% 81% 15% ..
1	B	136	13% 73% 21% 6% .
1	C	136	5% 85% 11% ..

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3265 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein argonaute-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	133	Total	C	N	O	S	0	0	0
			1036	659	185	182	10			
1	B	135	Total	C	N	O	S	0	0	0
			1047	666	187	184	10			
1	C	132	Total	C	N	O	S	0	0	0
			1032	657	184	181	10			

- Molecule 2 is [(3 {S},4 {R},5 {R})-5-[5-methyl-2,4-bis(oxidanylidene)pyrimidin-1-yl]-4-oxidanyl-oxolan-3-yl] [oxidanyl(phosphonooxy)phosphoryl] hydrogen phosphate (CCD ID: A1ANT) (formula: C₉H₁₅N₂O₁₄P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			28	9	2	14	3		
2	B	1	Total	C	N	O	P	0	0
			28	9	2	14	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			28	9	2	14	3		



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total	O	0	0
			28	28		
3	B	10	Total	O	0	0
			10	10		
3	C	28	Total	O	0	0
			28	28		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein argonaute-2



- Molecule 1: Protein argonaute-2



- Molecule 1: Protein argonaute-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.55Å 46.90Å 66.21Å 86.00° 74.02° 84.76°	Depositor
Resolution (Å)	20.50 – 1.90 20.50 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.50-1.90) 99.4 (20.50-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.21_5207	Depositor
R, R_{free}	0.250 , 0.288 0.247 , 0.283	Depositor DCC
R_{free} test set	1716 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.332	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,-h+l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3265	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1ANT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	2/1055 (0.2%)	1.10	3/1425 (0.2%)
1	B	1.06	2/1066 (0.2%)	1.11	4/1440 (0.3%)
1	C	1.06	4/1051 (0.4%)	1.15	5/1420 (0.4%)
All	All	1.04	8/3172 (0.3%)	1.12	12/4285 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	4
1	C	0	3
All	All	0	9

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	521	ILE	CG1-CD1	-9.83	1.13	1.51
1	B	521	ILE	CG1-CD1	-8.33	1.19	1.51
1	B	483	MET	SD-CE	7.08	1.97	1.79
1	C	548	GLN	CG-CD	6.57	1.68	1.52
1	A	521	ILE	CG1-CD1	-6.09	1.28	1.51
1	C	548	GLN	CD-OE1	5.62	1.34	1.23
1	A	563	LEU	CB-CG	-5.41	1.42	1.53
1	C	542	MET	SD-CE	5.27	1.92	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	548	GLN	CB-CG-CD	8.01	126.21	112.60
1	B	558	GLN	OE1-CD-NE2	-7.36	115.24	122.60
1	A	486	GLN	N-CA-C	-6.03	100.46	110.17
1	A	499	ASP	CA-CB-CG	5.91	118.51	112.60
1	C	548	GLN	OE1-CD-NE2	-5.79	116.81	122.60
1	C	480	ASP	CA-CB-CG	5.58	118.17	112.60
1	A	502	GLU	CB-CG-CD	5.57	122.07	112.60
1	C	548	GLN	CG-CD-OE1	5.49	131.77	120.80
1	B	558	GLN	CB-CG-CD	5.48	121.91	112.60
1	B	548	GLN	CB-CG-CD	-5.34	103.52	112.60
1	B	491	PHE	CA-CB-CG	5.14	118.94	113.80
1	C	552	VAL	N-CA-C	-5.01	107.50	111.81

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	460	ARG	Sidechain
1	A	506	ARG	Sidechain
1	B	475	ARG	Sidechain
1	B	479	ARG	Sidechain
1	B	506	ARG	Sidechain
1	B	554	ARG	Sidechain
1	C	460	ARG	Sidechain
1	C	479	ARG	Sidechain
1	C	534	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1036	0	1077	16	0
1	B	1047	0	1089	32	0
1	C	1032	0	1074	8	0
2	A	28	0	0	3	0
2	B	28	0	0	4	0
2	C	28	0	0	3	0
3	A	28	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
3	C	28	0	0	2	0
All	All	3265	0	3240	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:450:VAL:CG1	1:B:515:LEU:HA	1.80	1.11
1:B:450:VAL:HG12	1:B:515:LEU:HA	1.33	1.11
1:C:539:VAL:HG12	1:C:540:LEU:HD12	1.47	0.97
1:B:526:THR:CG2	1:B:528:VAL:HG22	1.95	0.96
1:C:539:VAL:HG12	1:C:540:LEU:CD1	2.11	0.81
1:A:529:TYR:HE1	2:A:601:A1ANT:O5	1.61	0.81
1:B:539:VAL:HG12	1:B:540:LEU:HD22	1.64	0.78
1:B:526:THR:HG21	1:B:528:VAL:HG22	1.68	0.76
1:B:463:THR:H	1:B:466:HIS:HD2	1.34	0.76
1:A:542:MET:HA	1:B:525:LYS:HE3	1.71	0.72
1:A:463:THR:H	1:A:466:HIS:HD2	1.40	0.68
1:B:483:MET:O	1:B:483:MET:HG2	1.95	0.66
1:B:526:THR:HG22	1:B:528:VAL:H	1.60	0.66
2:C:601:A1ANT:O11	3:C:701:HOH:O	2.15	0.65
1:B:545:GLN:NE2	2:B:601:A1ANT:O10	2.25	0.65
1:A:529:TYR:CE1	2:A:601:A1ANT:O5	2.46	0.65
1:A:566:LYS:NZ	2:A:601:A1ANT:O11	2.19	0.65
1:B:526:THR:HG22	1:B:528:VAL:HG22	1.78	0.64
2:C:601:A1ANT:O7	3:C:701:HOH:O	2.15	0.63
1:B:450:VAL:CG1	1:B:515:LEU:CA	2.70	0.62
1:B:534:ARG:O	1:B:538:THR:OG1	2.14	0.62
1:C:473:GLN:HA	1:C:476:LYS:HD3	1.83	0.60
1:A:458:PRO:HG2	1:A:461:GLN:HE21	1.69	0.58
1:B:450:VAL:HG12	1:B:515:LEU:CA	2.21	0.57
1:C:476:LYS:HG2	1:C:477:ILE:N	2.19	0.56
1:B:566:LYS:HZ3	2:B:601:A1ANT:C3	2.17	0.56
1:B:502:GLU:HB3	1:B:503:PRO:HD3	1.87	0.56
1:C:502:GLU:HB3	1:C:503:PRO:HD3	1.88	0.55
1:A:458:PRO:HG2	1:A:461:GLN:NE2	2.21	0.54
1:B:509:LYS:HE3	1:B:542:MET:HE2	1.90	0.54
1:B:521:ILE:HD12	1:B:552:VAL:HG21	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:566:LYS:NZ	2:B:601:A1ANT:C3	2.71	0.53
1:A:537:ASP:O	1:B:526:THR:CG2	2.58	0.52
1:B:526:THR:CB	1:B:528:VAL:HG22	2.40	0.51
1:B:471:THR:O	1:B:475:ARG:HG3	2.11	0.51
1:A:537:ASP:O	1:B:526:THR:HG21	2.11	0.50
1:C:539:VAL:CG1	1:C:540:LEU:CD1	2.87	0.49
1:A:543:ALA:H	1:B:525:LYS:HE3	1.78	0.49
1:A:515:LEU:O	1:B:525:LYS:NZ	2.46	0.48
1:A:502:GLU:HB3	1:A:503:PRO:HD3	1.95	0.48
1:B:460:ARG:NH1	1:C:537:ASP:O	2.46	0.48
1:B:491:PHE:CZ	1:B:493:LYS:HG3	2.49	0.48
1:B:533:LYS:HD2	2:B:601:A1ANT:O8	2.14	0.47
1:B:464:GLU:OE2	1:B:468:LYS:NZ	2.49	0.46
1:A:521:ILE:HD12	1:A:552:VAL:HG21	1.99	0.45
1:A:538:THR:O	1:B:458:PRO:HD3	2.18	0.44
1:C:548:GLN:CD	1:C:548:GLN:H	2.27	0.43
1:B:463:THR:H	1:B:466:HIS:CD2	2.25	0.42
1:A:534:ARG:O	1:A:538:THR:HB	2.20	0.41
2:C:601:A1ANT:O11	2:C:601:A1ANT:O7	2.38	0.41
1:A:542:MET:HA	1:B:525:LYS:CE	2.46	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/136 (96%)	129 (98%)	1 (1%)	1 (1%)	16	8
1	B	133/136 (98%)	133 (100%)	0	0	100	100
1	C	130/136 (96%)	130 (100%)	0	0	100	100
All	All	394/408 (97%)	392 (100%)	1 (0%)	1 (0%)	36	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	441	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/117 (98%)	114 (99%)	1 (1%)	70	73
1	B	116/117 (99%)	110 (95%)	6 (5%)	21	13
1	C	115/117 (98%)	112 (97%)	3 (3%)	40	35
All	All	346/351 (99%)	336 (97%)	10 (3%)	37	31

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	547	VAL
1	B	450	VAL
1	B	500	SER
1	B	525	LYS
1	B	539	VAL
1	B	540	LEU
1	B	574	VAL
1	C	476	LYS
1	C	496	GLN
1	C	548	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	GLN
1	A	466	HIS
1	A	488	GLN
1	A	496	GLN
1	A	507	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	443	HIS
1	B	466	HIS
1	B	496	GLN
1	C	488	GLN
1	C	551	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1ANT	C	601	-	28,29,29	0.77	0	38,46,46	0.79	0
2	A1ANT	A	601	-	28,29,29	0.72	0	38,46,46	0.95	0
2	A1ANT	B	601	-	28,29,29	0.74	0	38,46,46	1.13	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1ANT	C	601	-	-	8/21/34/34	0/2/2/2
2	A1ANT	A	601	-	-	11/21/34/34	0/2/2/2
2	A1ANT	B	601	-	-	6/21/34/34	0/2/2/2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	A1ANT	O5-P2-O6	2.92	115.17	107.27
2	B	601	A1ANT	O6-P2-O10	-2.90	101.98	110.70
2	B	601	A1ANT	O5-P2-O4	2.04	112.78	107.27

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	A1ANT	C1-C2-O2-P1
2	A	601	A1ANT	C3-C2-O2-P1
2	A	601	A1ANT	O12-C4-N1-C5
2	A	601	A1ANT	O12-C4-N1-C6
2	B	601	A1ANT	C1-C2-O2-P1
2	B	601	A1ANT	O12-C4-N1-C5
2	C	601	A1ANT	C3-C2-O2-P1
2	C	601	A1ANT	C1-C4-N1-C5
2	C	601	A1ANT	C1-C4-N1-C6
2	C	601	A1ANT	P2-O6-P3-O7
2	B	601	A1ANT	O12-C4-N1-C6
2	C	601	A1ANT	O12-C4-N1-C5
2	C	601	A1ANT	O12-C4-N1-C6
2	A	601	A1ANT	P2-O4-P1-O2
2	B	601	A1ANT	P2-O4-P1-O2
2	A	601	A1ANT	P3-O6-P2-O4
2	A	601	A1ANT	P1-O4-P2-O5
2	B	601	A1ANT	P3-O6-P2-O10
2	A	601	A1ANT	P1-O4-P2-O10
2	B	601	A1ANT	P3-O6-P2-O5
2	C	601	A1ANT	P1-O4-P2-O10
2	A	601	A1ANT	P2-O6-P3-O8
2	C	601	A1ANT	P2-O6-P3-O8
2	A	601	A1ANT	C1-C4-N1-C6
2	A	601	A1ANT	P3-O6-P2-O5


There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	601	A1ANT	3	0
2	A	601	A1ANT	3	0
2	B	601	A1ANT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	133/136 (97%)	0.50	5 (3%) 44 47	16, 30, 47, 84	0
1	B	135/136 (99%)	0.97	18 (13%) 7 7	18, 33, 55, 76	0
1	C	132/136 (97%)	0.59	7 (5%) 32 34	18, 31, 56, 87	0
All	All	400/408 (98%)	0.69	30 (7%) 20 21	16, 31, 55, 87	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	499	ASP	4.3
1	B	527	PRO	3.8
1	B	538	THR	3.7
1	B	490	CYS	3.7
1	B	514	GLY	3.5
1	C	541	GLY	3.2
1	A	513	ALA	3.2
1	B	574	VAL	2.9
1	B	506	ARG	2.8
1	C	538	THR	2.8
1	B	539	VAL	2.8
1	B	529	TYR	2.7
1	C	534	ARG	2.7
1	B	540	LEU	2.7
1	B	536	GLY	2.6
1	B	507	HIS	2.6
1	A	572	GLY	2.6
1	B	510	ASN	2.6
1	B	542	MET	2.6
1	B	530	ALA	2.6
1	C	539	VAL	2.6
1	A	440	LYS	2.5
1	C	445	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	487	GLY	2.4
1	A	444	THR	2.3
1	B	511	THR	2.2
1	B	508	LEU	2.1
1	C	443	HIS	2.0
1	B	513	ALA	2.0
1	B	485	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	A1ANT	A	601	28/28	0.71	0.15	47,110,140,160	0
2	A1ANT	C	601	28/28	0.72	0.18	78,115,153,168	0
2	A1ANT	B	601	28/28	0.73	0.16	48,105,137,150	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1ANT A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1ANT C 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1ANT B 601:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.