



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:53 AM UTC

PDB ID : 1BFN / pdb_00001bfm
Title : BETA-AMYLASE/BETA-CYCLODEXTRIN COMPLEX
Authors : Adachi, M.; Mikami, B.; Katsube, T.; Utsumi, S.
Deposited on : 1998-05-22
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	
2	B	7	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-AMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	0	0
			3915	2510	660	728	17			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	76	LEU	PHE	cloning artifact	UNP P10538
A	202	GLY	ARG	variant	UNP P10538
A	399	ARG	LYS	variant	UNP P10538

- Molecule 2 is an oligosaccharide called Cycloheptakis-(1-4)-(alpha-D-glucopyranose).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	7	Total	C	O	0	0	0
			77	42	35			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	318	Total	O	0	0
			318	318		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-AMYLASE



• Molecule 2: Cycloheptakis-(1-4)-(alpha-D-glucopyranose)



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	86.03Å 86.03Å 144.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.07 10.00 – 2.07	Depositor EDS
% Data completeness (in resolution range)	80.7 (10.00-2.07) 88.7 (10.00-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.34 (at 2.07Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.158 , 0.211 0.162 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.031 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4315	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	20/4021 (0.5%)	1.66	65/5463 (1.2%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	456	HIS	CD2-NE2	-7.22	1.29	1.37
1	A	23	VAL	CA-CB	7.13	1.62	1.54
1	A	273	HIS	CD2-NE2	-6.94	1.30	1.37
1	A	179	VAL	CA-CB	6.72	1.62	1.54
1	A	308	HIS	CD2-NE2	-6.67	1.30	1.37
1	A	146	HIS	CD2-NE2	-6.26	1.30	1.37
1	A	119	HIS	CD2-NE2	-6.25	1.30	1.37
1	A	226	HIS	CD2-NE2	-6.21	1.31	1.37
1	A	442	HIS	CD2-NE2	-6.15	1.31	1.37
1	A	335	HIS	CD2-NE2	-6.14	1.31	1.37
1	A	335	HIS	CG-ND1	-6.00	1.31	1.38
1	A	336	HIS	CD2-NE2	-5.91	1.31	1.37
1	A	273	HIS	CG-ND1	-5.70	1.31	1.38
1	A	93	HIS	CD2-NE2	-5.52	1.31	1.37
1	A	300	HIS	CD2-NE2	-5.41	1.31	1.37
1	A	119	HIS	CG-ND1	-5.18	1.32	1.38
1	A	456	HIS	CG-ND1	-5.16	1.32	1.38
1	A	146	HIS	CG-ND1	-5.15	1.32	1.38
1	A	175	ILE	CA-CB	5.15	1.60	1.54
1	A	308	HIS	CG-ND1	-5.01	1.32	1.38

All (65) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	VAL	N-CA-CB	-12.03	98.97	112.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	419	LEU	O-C-N	10.99	136.13	123.05
1	A	70	ARG	N-CA-C	9.06	120.76	111.07
1	A	30	GLU	N-CA-C	9.02	120.72	111.07
1	A	59	ILE	N-CA-C	8.55	118.57	110.53
1	A	241	VAL	N-CA-CB	-8.51	99.29	111.21
1	A	399	ARG	CA-C-N	7.89	127.81	119.05
1	A	399	ARG	C-N-CA	7.89	127.81	119.05
1	A	176	ASP	CA-CB-CG	7.71	120.31	112.60
1	A	420	ARG	CA-CB-CG	-7.47	99.16	114.10
1	A	420	ARG	N-CA-C	7.46	126.69	110.80
1	A	419	LEU	CA-C-N	7.24	135.37	121.54
1	A	419	LEU	C-N-CA	7.24	135.37	121.54
1	A	255	VAL	CB-CA-C	7.12	119.75	110.84
1	A	102	ILE	N-CA-C	7.12	117.21	110.30
1	A	24	ASN	CA-CB-CG	7.01	119.61	112.60
1	A	407	GLY	CA-C-N	6.41	124.27	119.66
1	A	407	GLY	C-N-CA	6.41	124.27	119.66
1	A	70	ARG	NE-CZ-NH1	6.37	127.87	121.50
1	A	456	HIS	CB-CG-CD2	-6.28	123.04	131.20
1	A	444	ASP	N-CA-C	6.10	120.52	113.38
1	A	409	PRO	CA-C-O	-6.04	114.56	121.56
1	A	93	HIS	CB-CG-CD2	-5.97	123.44	131.20
1	A	101	ASP	CA-C-N	5.91	128.09	120.70
1	A	101	ASP	C-N-CA	5.91	128.09	120.70
1	A	309	ALA	N-CA-C	5.86	117.67	111.28
1	A	102	ILE	CB-CG1-CD1	5.83	126.05	113.80
1	A	428	LYS	N-CA-C	5.82	118.11	111.02
1	A	308	HIS	CB-CG-CD2	-5.74	123.74	131.20
1	A	385	ARG	NE-CZ-NH1	5.74	127.24	121.50
1	A	485	TRP	CB-CG-CD1	-5.69	118.37	126.90
1	A	420	ARG	O-C-N	5.68	130.15	122.59
1	A	321	ASP	CA-CB-CG	5.63	118.23	112.60
1	A	450	ASN	CA-C-N	5.60	125.44	119.28
1	A	450	ASN	C-N-CA	5.60	125.44	119.28
1	A	101	ASP	CA-C-O	5.59	127.06	120.69
1	A	23	VAL	N-CA-C	-5.55	100.40	108.17
1	A	385	ARG	NE-CZ-NH2	-5.54	114.21	119.20
1	A	6	ASN	OD1-CG-ND2	-5.52	117.08	122.60
1	A	118	ASN	OD1-CG-ND2	-5.51	117.09	122.60
1	A	449	ALA	N-CA-C	5.50	118.20	111.82
1	A	376	ARG	CA-CB-CG	-5.46	103.17	114.10
1	A	330	ARG	NE-CZ-NH2	-5.43	114.31	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	241	VAL	CB-CA-C	5.42	121.23	111.36
1	A	101	ASP	O-C-N	5.42	129.38	123.04
1	A	182	GLY	CA-C-N	5.39	125.40	119.90
1	A	182	GLY	C-N-CA	5.39	125.40	119.90
1	A	62	LYS	N-CA-C	5.38	117.89	111.71
1	A	174	ILE	CA-C-N	5.37	127.43	120.56
1	A	174	ILE	C-N-CA	5.37	127.43	120.56
1	A	374	ASP	CA-CB-CG	5.37	117.97	112.60
1	A	495	GLY	N-CA-C	5.31	128.69	113.30
1	A	257	GLU	CA-CB-CG	5.29	124.68	114.10
1	A	164	ASN	OD1-CG-ND2	-5.28	117.33	122.60
1	A	357	SER	N-CA-CB	-5.26	101.86	110.43
1	A	485	TRP	CG-CD2-CE3	5.23	139.13	133.90
1	A	419	LEU	CA-C-O	5.22	125.97	120.54
1	A	62	LYS	CA-CB-CG	-5.19	103.71	114.10
1	A	494	ASP	CA-C-N	5.17	131.00	121.70
1	A	494	ASP	C-N-CA	5.17	131.00	121.70
1	A	486	LEU	CA-C-N	5.10	124.71	119.05
1	A	486	LEU	C-N-CA	5.10	124.71	119.05
1	A	490	ASP	CA-CB-CG	5.05	117.65	112.60
1	A	131	ASN	OD1-CG-ND2	-5.04	117.56	122.60
1	A	346	MET	N-CA-C	5.04	118.17	110.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3915	0	3814	22	0
2	B	77	0	63	0	0
3	A	5	0	0	0	0
4	A	318	0	0	0	0
All	All	4315	0	3877	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:GLU:H	1:A:239:ASN:HD21	1.40	0.69
1:A:7:MET:HE1	1:A:403:VAL:HG11	1.84	0.59
1:A:347:ARG:HH11	1:A:393:GLN:NE2	2.01	0.58
1:A:367:LEU:HD22	1:A:377:VAL:HG11	1.86	0.56
1:A:193:PRO:HG2	1:A:196:GLN:HB2	1.89	0.55
1:A:68:ASP:OD1	1:A:70:ARG:HD3	2.06	0.54
1:A:246:GLY:HA2	1:A:249:LYS:HE3	1.89	0.54
1:A:330:ARG:HG2	1:A:473:LEU:HD12	1.91	0.52
1:A:351:GLN:HG3	1:A:357:SER:OG	2.10	0.52
1:A:294:ILE:HG12	1:A:332:LEU:HD21	2.00	0.44
1:A:131:ASN:HD22	1:A:132:LYS:N	2.15	0.44
1:A:367:LEU:HD22	1:A:377:VAL:CG1	2.48	0.44
1:A:199:GLU:H	1:A:239:ASN:ND2	2.10	0.43
1:A:401:GLN:HE22	1:A:410:LYS:HE3	1.83	0.42
1:A:122:PHE:HB3	1:A:131:ASN:O	2.19	0.42
1:A:199:GLU:N	1:A:239:ASN:HD21	2.12	0.42
1:A:142:GLU:O	1:A:149:THR:HA	2.19	0.42
1:A:298:GLY:H	1:A:342:THR:CG2	2.33	0.41
1:A:436:LYS:O	1:A:440:LYS:HG2	2.21	0.41
1:A:98:ASN:ND2	1:A:195:SER:HB3	2.36	0.41
1:A:286:LEU:HD21	1:A:463:PRO:HD3	2.01	0.40
1:A:458:ILE:HG22	1:A:459:THR:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	488/495 (99%)	475 (97%)	12 (2%)	1 (0%)	43 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	420	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	420/424 (99%)	384 (91%)	36 (9%)	10 4

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	20	LEU
1	A	23	VAL
1	A	35	LEU
1	A	39	LEU
1	A	42	LEU
1	A	70	ARG
1	A	75	LEU
1	A	102	ILE
1	A	103	VAL
1	A	112	LEU
1	A	130	ARG
1	A	135	LEU
1	A	179	VAL
1	A	181	LEU
1	A	201	PRO
1	A	231	LEU
1	A	241	VAL
1	A	255	VAL
1	A	278	LEU
1	A	279	ASP
1	A	286	LEU
1	A	290	VAL
1	A	292	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	330	ARG
1	A	353	SER
1	A	421	LEU
1	A	426	LEU
1	A	427	GLN
1	A	429	SER
1	A	461	LEU
1	A	462	LYS
1	A	473	LEU
1	A	474	LEU
1	A	486	LEU
1	A	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	38	GLN
1	A	66	GLN
1	A	80	GLN
1	A	98	ASN
1	A	118	ASN
1	A	131	ASN
1	A	239	ASN
1	A	276	GLN
1	A	340	ASN
1	A	393	GLN
1	A	401	GLN
1	A	450	ASN
1	A	456	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	11,11,12	1.03	0	15,15,17	1.04	1 (6%)
2	GLC	B	2	2	11,11,12	0.66	0	15,15,17	1.21	1 (6%)
2	GLC	B	3	2	11,11,12	0.87	0	15,15,17	0.98	1 (6%)
2	GLC	B	4	2	11,11,12	0.75	0	15,15,17	0.92	1 (6%)
2	GLC	B	5	2	11,11,12	0.61	0	15,15,17	1.08	1 (6%)
2	GLC	B	6	2	11,11,12	0.98	0	15,15,17	1.03	2 (13%)
2	GLC	B	7	2	11,11,12	1.13	1 (9%)	15,15,17	1.32	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	0/2/19/22	0/1/1/1
2	GLC	B	4	2	-	1/2/19/22	0/1/1/1
2	GLC	B	5	2	-	0/2/19/22	0/1/1/1
2	GLC	B	6	2	-	0/2/19/22	0/1/1/1
2	GLC	B	7	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	7	GLC	C2-C3	2.23	1.55	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	GLC	C1-O5-C5	3.97	117.50	112.19
2	B	7	GLC	C1-O5-C5	3.80	117.28	112.19
2	B	5	GLC	C1-O5-C5	3.17	116.44	112.19
2	B	6	GLC	O4-C4-C3	-2.55	104.36	110.38
2	B	4	GLC	C1-O5-C5	2.54	115.59	112.19
2	B	1	GLC	O4-C4-C3	-2.21	105.18	110.38
2	B	3	GLC	C3-C4-C5	2.14	114.11	110.23
2	B	7	GLC	C3-C4-C5	-2.10	106.42	110.23
2	B	6	GLC	C1-O5-C5	2.07	114.95	112.19

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	4	GLC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection.

RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	503	-	4,4,4	0.39	0	6,6,6	0.56	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/495 (98%)	-0.90	2 (0%) 88 90	11, 18, 36, 67	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	ASN	2.8
1	A	495	GLY	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	2	11/12	0.83	0.09	49,52,55,59	0
2	GLC	B	3	11/12	0.83	0.09	46,53,58,58	0
2	GLC	B	1	11/12	0.86	0.09	42,47,51,54	0
2	GLC	B	7	11/12	0.87	0.07	47,50,52,52	0
2	GLC	B	4	11/12	0.88	0.08	37,46,49,53	0
2	GLC	B	6	11/12	0.90	0.06	32,35,41,44	0
2	GLC	B	5	11/12	0.93	0.06	25,27,29,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	503	5/5	0.91	0.09	52,52,55,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.