



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:20 AM UTC

PDB ID : 2BF8 / pdb_00002bf8
Title : Crystal structure of SUMO modified ubiquitin conjugating enzyme E2- 25K
Authors : Pichler, A.; Knipscheer, P.; Oberhofer, E.; Van Dijk, W.J.; Korner, R.; Vellaard Olsen, J.; Jentsch, S.; Melchior, F.; Sixma, T.K.
Deposited on : 2004-12-06
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	 3% 77% 18% . .
2	B	77	 16% 71% 25% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1947 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUITIN-CONJUGATING ENZYME E2-25 KDA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	154	1237	793	210	231	3	0	1	0

- Molecule 2 is a protein called UBIQUITIN-LIKE PROTEIN SMT3C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	77	627	395	108	120	4	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	70	Total	O	0	0
			70	70		
3	B	13	Total	O	0	0
			13	13		

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	58.39Å 58.39Å 162.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	55.05 – 2.30 54.96 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (55.05-2.30) 99.6 (54.96-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.210 , 0.279 0.213 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	54.4	Xtrriage
Anisotropy	0.266	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 69.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1947	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	2/1266 (0.2%)	1.09	0/1721
2	B	0.83	0/637	1.00	0/851
All	All	1.02	2/1903 (0.1%)	1.06	0/2572

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	62	ILE	CA-CB	5.48	1.59	1.53
1	A	97	LYS	CA-C	5.12	1.59	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1237	0	1236	18	1
2	B	627	0	626	8	1
3	A	70	0	0	1	0
3	B	13	0	0	0	0
All	All	1947	0	1862	25	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:64:PHE:HB3	2:B:88:ILE:HD11	1.45	0.96
1:A:31:LEU:HD12	3:A:2025:HOH:O	2.00	0.60
1:A:147:LEU:O	1:A:151:VAL:HG23	2.04	0.57
1:A:20:GLU:OE2	1:A:24:LYS:HD2	2.11	0.51
1:A:25:ASN:HD22	1:A:25:ASN:N	2.09	0.50
1:A:21:GLU:HG2	1:A:111:LEU:HD21	1.93	0.50
2:B:59:MET:O	2:B:60:ASN:HB2	2.11	0.50
1:A:94:ASP:HB3	1:A:99:GLN:HE21	1.77	0.49
1:A:33:ASP:OD2	1:A:35:ASN:HB2	2.12	0.49
1:A:25:ASN:N	1:A:25:ASN:ND2	2.60	0.49
2:B:27:ILE:HA	2:B:32:SER:O	2.12	0.48
1:A:122:PRO:HB2	1:A:135:LYS:CD	2.45	0.46
2:B:62:LEU:HD13	2:B:90:VAL:HG21	1.98	0.46
1:A:93:LEU:HD21	1:A:112:SER:OG	2.16	0.45
1:A:122:PRO:HB2	1:A:135:LYS:HD3	1.98	0.45
2:B:44:LEU:HD13	2:B:71:ILE:CG2	2.46	0.45
1:A:26:GLN:HB3	1:A:114:GLN:HE22	1.82	0.45
2:B:40:MET:SD	2:B:77:PRO:HB2	2.58	0.44
1:A:43:ILE:HD12	1:A:117:LEU:HD22	2.00	0.43
1:A:3:ASN:ND2	2:B:92:GLN:OE1	2.52	0.43
1:A:50:PRO:HB3	1:A:146:ARG:HG3	2.00	0.42
1:A:94:ASP:CB	1:A:99:GLN:HE21	2.32	0.42
2:B:59:MET:O	2:B:60:ASN:CB	2.68	0.42
1:A:125:PRO:HB3	1:A:131:ALA:CB	2.50	0.41
1:A:79:ILE:CD1	1:A:117:LEU:HD11	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ALA:O	2:B:54:ARG:NH1[7_556]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/159 (96%)	150 (98%)	3 (2%)	0	100	100
2	B	75/77 (97%)	68 (91%)	6 (8%)	1 (1%)	9	10
All	All	228/236 (97%)	218 (96%)	9 (4%)	1 (0%)	30	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	31	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/133 (99%)	126 (96%)	6 (4%)	24	37
2	B	70/70 (100%)	60 (86%)	10 (14%)	3	3
All	All	202/203 (100%)	186 (92%)	16 (8%)	11	15

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	20	GLU
1	A	25	ASN
1	A	116	LEU
1	A	117	LEU
1	A	129	VAL
2	B	22	ILE
2	B	27	ILE
2	B	38	VAL
2	B	42	THR
2	B	47	LEU
2	B	76	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	79	GLU
2	B	88	ILE
2	B	92	GLN
2	B	94	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	25	ASN
1	A	99	GLN
1	A	114	GLN
1	A	136	GLN
2	B	55	GLN
2	B	60	ASN
2	B	92	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/159 (96%)	0.78	5 (3%) 50 52	28, 58, 63, 65	1 (0%)
2	B	77/77 (100%)	1.20	12 (15%) 5 6	53, 58, 62, 65	0
All	All	231/236 (97%)	0.92	17 (7%) 20 22	28, 58, 63, 65	1 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	77	PRO	3.3
2	B	56	GLY	2.7
1	A	31	LEU	2.7
1	A	120	ALA	2.7
2	B	38	VAL	2.7
1	A	68[A]	PHE	2.6
2	B	88	ILE	2.6
2	B	57	VAL	2.5
1	A	32	VAL	2.4
2	B	54	ARG	2.3
2	B	47	LEU	2.3
2	B	22	ILE	2.3
2	B	76	THR	2.2
1	A	98	ASP	2.2
2	B	79	GLU	2.1
2	B	87	VAL	2.1
2	B	44	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.