



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 09:31 PM UTC

PDB ID : 7BFI / pdb_00007bfi
Title : A double-histidine mutant of HSP47 slows down client release at low pH
Authors : Oecal, S.; Baumann, U.
Deposited on : 2021-01-03
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	 2% 81% 13% • 5%
1	B	393	 2% 79% 14% 6%
1	C	393	 3% 84% 10% • •
1	D	393	 2% 80% 13% 7%
2	E	17	 82% 6% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	17	 76% 12% 12%
2	G	17	 6% 76% 6% 18%
2	H	17	 76% 12% 12%
2	I	17	 76% 12% 12%
2	J	17	 82% 6% 12%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 12313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Collagen-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	372	2923	1856	511	544	12	0	0	0
1	C	376	2956	1877	515	552	12	0	0	0
1	D	367	2878	1829	502	535	12	0	0	0
1	B	369	2901	1844	508	537	12	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	initiating methionine	UNP E2RHY7
A	273	ASN	HIS	engineered mutation	UNP E2RHY7
A	274	ASN	HIS	engineered mutation	UNP E2RHY7
A	419	LEU	-	expression tag	UNP E2RHY7
A	420	GLU	-	expression tag	UNP E2RHY7
A	421	HIS	-	expression tag	UNP E2RHY7
A	422	HIS	-	expression tag	UNP E2RHY7
A	423	HIS	-	expression tag	UNP E2RHY7
A	424	HIS	-	expression tag	UNP E2RHY7
A	425	HIS	-	expression tag	UNP E2RHY7
A	426	HIS	-	expression tag	UNP E2RHY7
C	34	MET	-	initiating methionine	UNP E2RHY7
C	273	ASN	HIS	engineered mutation	UNP E2RHY7
C	274	ASN	HIS	engineered mutation	UNP E2RHY7
C	419	LEU	-	expression tag	UNP E2RHY7
C	420	GLU	-	expression tag	UNP E2RHY7
C	421	HIS	-	expression tag	UNP E2RHY7
C	422	HIS	-	expression tag	UNP E2RHY7
C	423	HIS	-	expression tag	UNP E2RHY7
C	424	HIS	-	expression tag	UNP E2RHY7
C	425	HIS	-	expression tag	UNP E2RHY7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	426	HIS	-	expression tag	UNP E2RHY7
D	34	MET	-	initiating methionine	UNP E2RHY7
D	273	ASN	HIS	engineered mutation	UNP E2RHY7
D	274	ASN	HIS	engineered mutation	UNP E2RHY7
D	419	LEU	-	expression tag	UNP E2RHY7
D	420	GLU	-	expression tag	UNP E2RHY7
D	421	HIS	-	expression tag	UNP E2RHY7
D	422	HIS	-	expression tag	UNP E2RHY7
D	423	HIS	-	expression tag	UNP E2RHY7
D	424	HIS	-	expression tag	UNP E2RHY7
D	425	HIS	-	expression tag	UNP E2RHY7
D	426	HIS	-	expression tag	UNP E2RHY7
B	34	MET	-	initiating methionine	UNP E2RHY7
B	273	ASN	HIS	engineered mutation	UNP E2RHY7
B	274	ASN	HIS	engineered mutation	UNP E2RHY7
B	419	LEU	-	expression tag	UNP E2RHY7
B	420	GLU	-	expression tag	UNP E2RHY7
B	421	HIS	-	expression tag	UNP E2RHY7
B	422	HIS	-	expression tag	UNP E2RHY7
B	423	HIS	-	expression tag	UNP E2RHY7
B	424	HIS	-	expression tag	UNP E2RHY7
B	425	HIS	-	expression tag	UNP E2RHY7
B	426	HIS	-	expression tag	UNP E2RHY7

- Molecule 2 is a protein called 15R8 collagen model peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	F	15	Total	C	N	O	0	0	0
			93	61	17	15			
2	H	15	Total	C	N	O	0	0	0
			93	61	17	15			
2	I	15	Total	C	N	O	0	0	0
			93	61	17	15			
2	J	15	Total	C	N	O	0	0	0
			93	61	17	15			
2	E	15	Total	C	N	O	0	0	0
			93	61	17	15			
2	G	14	Total	C	N	O	0	0	0
			86	56	16	14			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	28	Total O 28 28	0	0
3	C	18	Total O 18 18	0	0
3	D	27	Total O 27 27	0	0
3	B	28	Total O 28 28	0	0
3	H	1	Total O 1 1	0	0
3	I	1	Total O 1 1	0	0
3	J	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Collagen-binding protein




- Molecule 1: Collagen-binding protein




- Molecule 1: Collagen-binding protein




- Molecule 1: Collagen-binding protein




- Molecule 2: 15R8 collagen model peptide




- Molecule 2: 15R8 collagen model peptide




- Molecule 2: 15R8 collagen model peptide





- Molecule 2: 15R8 collagen model peptide



- Molecule 2: 15R8 collagen model peptide



- Molecule 2: 15R8 collagen model peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.70Å 77.41Å 122.34Å 90.00° 96.60° 90.00°	Depositor
Resolution (Å)	78.18 – 2.44 78.18 – 2.44	Depositor EDS
% Data completeness (in resolution range)	96.8 (78.18-2.44) 96.8 (78.18-2.44)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.19_4092	Depositor
R, R_{free}	0.217 , 0.248 (Not available) , 0.223	Depositor DCC
R_{free} test set	2172 reflections (3.97%)	wwPDB-VP
Wilson B-factor (Å ²)	52.5	Xtrriage
Anisotropy	0.556	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12313	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/2980	0.35	0/4019
1	B	0.12	0/2958	0.33	0/3989
1	C	0.12	0/3015	0.32	0/4069
1	D	0.13	0/2935	0.31	0/3960
2	E	0.96	1/99 (1.0%)	0.31	0/141
2	F	1.00	1/99 (1.0%)	0.32	0/141
2	G	1.05	1/91 (1.1%)	0.33	0/129
2	H	1.01	1/99 (1.0%)	0.30	0/141
2	I	1.03	1/99 (1.0%)	0.27	0/141
2	J	1.00	1/99 (1.0%)	0.32	0/141
All	All	0.25	6/12474 (0.0%)	0.33	0/16871

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	0	ACE	C-N	-10.16	1.33	1.43
2	H	0	ACE	C-N	-9.99	1.33	1.43
2	G	0	ACE	C-N	-9.93	1.33	1.43
2	F	0	ACE	C-N	-9.88	1.33	1.43
2	J	0	ACE	C-N	-9.88	1.33	1.43

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2923	0	2955	29	0
1	B	2901	0	2943	35	0
1	C	2956	0	2980	23	1
1	D	2878	0	2908	33	1
2	E	93	0	91	0	0
2	F	93	0	91	1	0
2	G	86	0	84	0	0
2	H	93	0	91	1	0
2	I	93	0	91	1	0
2	J	93	0	91	0	0
3	A	28	0	0	1	0
3	B	28	0	0	0	0
3	C	18	0	0	0	0
3	D	27	0	0	2	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
All	All	12313	0	12325	116	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 116 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:LYS:NZ	1:A:182:ASP:OD2	2.10	0.83
1:D:126:VAL:HG12	1:D:209:HIS:H	1.59	0.68
1:D:125:ASN:HB2	1:D:261:ALA:HB3	1.77	0.67
1:A:95:ALA:HA	1:A:100:GLU:OE2	1.96	0.66
1:D:299:LYS:NZ	3:D:501:HOH:O	2.27	0.66

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:378:SER:OG	1:D:144:GLU:OE1[2_654]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/393 (94%)	357 (97%)	10 (3%)	1 (0%)	36	44
1	B	365/393 (93%)	355 (97%)	10 (3%)	0	100	100
1	C	374/393 (95%)	364 (97%)	10 (3%)	0	100	100
1	D	363/393 (92%)	353 (97%)	10 (3%)	0	100	100
2	E	13/17 (76%)	13 (100%)	0	0	100	100
2	F	13/17 (76%)	13 (100%)	0	0	100	100
2	G	12/17 (71%)	12 (100%)	0	0	100	100
2	H	13/17 (76%)	13 (100%)	0	0	100	100
2	I	13/17 (76%)	13 (100%)	0	0	100	100
2	J	13/17 (76%)	13 (100%)	0	0	100	100
All	All	1547/1674 (92%)	1506 (97%)	40 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	124	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/338 (94%)	312 (98%)	5 (2%)	55	68
1	B	315/338 (93%)	312 (99%)	3 (1%)	68	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	320/338 (95%)	316 (99%)	4 (1%)	61	72
1	D	312/338 (92%)	311 (100%)	1 (0%)	86	91
2	E	10/10 (100%)	10 (100%)	0	100	100
2	F	10/10 (100%)	10 (100%)	0	100	100
2	G	9/10 (90%)	9 (100%)	0	100	100
2	H	10/10 (100%)	10 (100%)	0	100	100
2	I	10/10 (100%)	10 (100%)	0	100	100
2	J	10/10 (100%)	10 (100%)	0	100	100
All	All	1323/1412 (94%)	1310 (99%)	13 (1%)	68	77

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	125	ASN
1	C	375	GLU
1	B	264	LEU
1	B	161	ILE
1	B	171	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	216	HIS
1	B	289	GLN
1	D	216	HIS
1	D	254	GLN
1	D	289	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/393 (94%)	0.14	5 (1%) 75 77	38, 56, 85, 129	0
1	B	369/393 (93%)	0.15	7 (1%) 66 67	33, 57, 92, 129	0
1	C	376/393 (95%)	0.26	10 (2%) 56 55	42, 65, 106, 187	0
1	D	367/393 (93%)	0.31	8 (2%) 62 63	39, 64, 111, 154	0
2	E	14/17 (82%)	0.06	0 100 100	41, 48, 76, 84	0
2	F	14/17 (82%)	-0.01	0 100 100	44, 53, 77, 86	0
2	G	13/17 (76%)	-0.18	1 (7%) 19 16	40, 46, 75, 81	0
2	H	14/17 (82%)	0.09	0 100 100	44, 49, 72, 89	0
2	I	14/17 (82%)	-0.02	0 100 100	46, 54, 73, 76	0
2	J	14/17 (82%)	-0.24	0 100 100	43, 50, 77, 83	0
All	All	1567/1674 (93%)	0.20	31 (1%) 65 66	33, 60, 100, 187	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	346	LEU	3.9
1	C	370	ILE	3.5
1	B	346	LEU	3.3
1	B	345	ASP	3.1
1	A	125	ASN	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.