



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:43 AM UTC

PDB ID : 8BFI / pdb_00008bfi
Title : human CNOT1-CNOT10-CNOT11 module
Authors : Basquin, J.; Ozgur, S.; Conti, E.
Deposited on : 2022-10-26
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	778	
2	B	718	
3	C	455	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT transcription complex subunit 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	602	4496	2878	780	817	10	11	0	0	0

There are 94 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-90	MSE	-	initiating methionine	UNP A5YKK6
A	-89	HIS	-	expression tag	UNP A5YKK6
A	-88	HIS	-	expression tag	UNP A5YKK6
A	-87	HIS	-	expression tag	UNP A5YKK6
A	-86	HIS	-	expression tag	UNP A5YKK6
A	-85	HIS	-	expression tag	UNP A5YKK6
A	-84	HIS	-	expression tag	UNP A5YKK6
A	-83	GLY	-	expression tag	UNP A5YKK6
A	-82	MSE	-	expression tag	UNP A5YKK6
A	-81	GLU	-	expression tag	UNP A5YKK6
A	-80	GLY	-	expression tag	UNP A5YKK6
A	-79	GLU	-	expression tag	UNP A5YKK6
A	-78	TYR	-	expression tag	UNP A5YKK6
A	-77	ILE	-	expression tag	UNP A5YKK6
A	-76	LYS	-	expression tag	UNP A5YKK6
A	-75	LEU	-	expression tag	UNP A5YKK6
A	-74	LYS	-	expression tag	UNP A5YKK6
A	-73	VAL	-	expression tag	UNP A5YKK6
A	-72	ILE	-	expression tag	UNP A5YKK6
A	-71	GLY	-	expression tag	UNP A5YKK6
A	-70	GLN	-	expression tag	UNP A5YKK6
A	-69	ASP	-	expression tag	UNP A5YKK6
A	-68	SER	-	expression tag	UNP A5YKK6
A	-67	SER	-	expression tag	UNP A5YKK6
A	-66	GLU	-	expression tag	UNP A5YKK6
A	-65	ILE	-	expression tag	UNP A5YKK6
A	-64	HIS	-	expression tag	UNP A5YKK6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-63	PHE	-	expression tag	UNP A5YKK6
A	-62	LYS	-	expression tag	UNP A5YKK6
A	-61	VAL	-	expression tag	UNP A5YKK6
A	-60	LYS	-	expression tag	UNP A5YKK6
A	-59	MSE	-	expression tag	UNP A5YKK6
A	-58	THR	-	expression tag	UNP A5YKK6
A	-57	THR	-	expression tag	UNP A5YKK6
A	-56	HIS	-	expression tag	UNP A5YKK6
A	-55	LEU	-	expression tag	UNP A5YKK6
A	-54	LYS	-	expression tag	UNP A5YKK6
A	-53	LYS	-	expression tag	UNP A5YKK6
A	-52	LEU	-	expression tag	UNP A5YKK6
A	-51	LYS	-	expression tag	UNP A5YKK6
A	-50	GLU	-	expression tag	UNP A5YKK6
A	-49	SER	-	expression tag	UNP A5YKK6
A	-48	TYR	-	expression tag	UNP A5YKK6
A	-47	CYS	-	expression tag	UNP A5YKK6
A	-46	GLN	-	expression tag	UNP A5YKK6
A	-45	ARG	-	expression tag	UNP A5YKK6
A	-44	GLN	-	expression tag	UNP A5YKK6
A	-43	GLY	-	expression tag	UNP A5YKK6
A	-42	VAL	-	expression tag	UNP A5YKK6
A	-41	PRO	-	expression tag	UNP A5YKK6
A	-40	MSE	-	expression tag	UNP A5YKK6
A	-39	ASN	-	expression tag	UNP A5YKK6
A	-38	SER	-	expression tag	UNP A5YKK6
A	-37	LEU	-	expression tag	UNP A5YKK6
A	-36	ARG	-	expression tag	UNP A5YKK6
A	-35	PHE	-	expression tag	UNP A5YKK6
A	-34	LEU	-	expression tag	UNP A5YKK6
A	-33	PHE	-	expression tag	UNP A5YKK6
A	-32	GLU	-	expression tag	UNP A5YKK6
A	-31	GLY	-	expression tag	UNP A5YKK6
A	-30	GLN	-	expression tag	UNP A5YKK6
A	-29	ARG	-	expression tag	UNP A5YKK6
A	-28	ILE	-	expression tag	UNP A5YKK6
A	-27	ALA	-	expression tag	UNP A5YKK6
A	-26	ASP	-	expression tag	UNP A5YKK6
A	-25	ASN	-	expression tag	UNP A5YKK6
A	-24	HIS	-	expression tag	UNP A5YKK6
A	-23	THR	-	expression tag	UNP A5YKK6
A	-22	PRO	-	expression tag	UNP A5YKK6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	LYS	-	expression tag	UNP A5YKK6
A	-20	GLU	-	expression tag	UNP A5YKK6
A	-19	LEU	-	expression tag	UNP A5YKK6
A	-18	GLY	-	expression tag	UNP A5YKK6
A	-17	MSE	-	expression tag	UNP A5YKK6
A	-16	GLU	-	expression tag	UNP A5YKK6
A	-15	GLU	-	expression tag	UNP A5YKK6
A	-14	GLU	-	expression tag	UNP A5YKK6
A	-13	ASP	-	expression tag	UNP A5YKK6
A	-12	VAL	-	expression tag	UNP A5YKK6
A	-11	ILE	-	expression tag	UNP A5YKK6
A	-10	GLU	-	expression tag	UNP A5YKK6
A	-9	VAL	-	expression tag	UNP A5YKK6
A	-8	TYR	-	expression tag	UNP A5YKK6
A	-7	GLN	-	expression tag	UNP A5YKK6
A	-6	GLU	-	expression tag	UNP A5YKK6
A	-5	GLN	-	expression tag	UNP A5YKK6
A	-4	THR	-	expression tag	UNP A5YKK6
A	-3	GLY	-	expression tag	UNP A5YKK6
A	-2	GLY	-	expression tag	UNP A5YKK6
A	-1	ARG	-	expression tag	UNP A5YKK6
A	0	SER	-	expression tag	UNP A5YKK6
A	381	VAL	ILE	conflict	UNP A5YKK6
A	649	VAL	ALA	conflict	UNP A5YKK6
A	681	THR	MET	conflict	UNP A5YKK6

- Molecule 2 is a protein called CCR4-NOT transcription complex subunit 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	574	4437	2818	767	824	28	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP Q9H9A5
B	-2	PRO	-	expression tag	UNP Q9H9A5
B	-1	ASP	-	expression tag	UNP Q9H9A5
B	0	SER	-	expression tag	UNP Q9H9A5

- Molecule 3 is a protein called CCR4-NOT transcription complex subunit 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	410	2957	1900	499	543	15	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	56	GLY	-	expression tag	UNP Q9UKZ1
C	57	PRO	-	expression tag	UNP Q9UKZ1
C	58	ASP	-	expression tag	UNP Q9UKZ1
C	59	ARG	-	expression tag	UNP Q9UKZ1
C	60	SER	-	expression tag	UNP Q9UKZ1
C	507	GLU	LYS	conflict	UNP Q9UKZ1

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.76Å 110.64Å 128.58Å 90.00° 103.39° 90.00°	Depositor
Resolution (Å)	45.77 – 3.00 45.77 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.77-3.00) 99.9 (45.77-3.00)	Depositor EDS
R_{merge}	0.32	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.262 , 0.283 (Not available) , 0.261	Depositor DCC
R_{free} test set	2293 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	79.5	Xtrriage
Anisotropy	0.274	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 67.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	11890	wwPDB-VP
Average B, all atoms (Å ²)	96.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/4577	0.45	4/6205 (0.1%)
2	B	0.10	0/4513	0.32	0/6107
3	C	0.12	0/3024	0.34	0/4136
All	All	0.11	0/12114	0.38	4/16448 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
3	C	0	1
All	All	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	LYS	CA-C-N	9.25	138.35	121.70
1	A	66	LYS	C-N-CA	9.25	138.35	121.70
1	A	513	ASN	CA-C-N	9.07	138.02	121.70
1	A	513	ASN	C-N-CA	9.07	138.02	121.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	189	LEU	Peptide
1	A	285	VAL	Peptide
1	A	368	ILE	Peptide
3	C	248	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4496	0	4271	71	0
2	B	4437	0	4406	65	0
3	C	2957	0	2801	37	0
All	All	11890	0	11478	148	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 148 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:VAL:HG12	1:A:390:PHE:H	1.34	0.93
3:C:459:LEU:HD23	3:C:464:ILE:HD12	1.68	0.75
1:A:251:MSE:HE1	2:B:361:CYS:HB3	1.68	0.74
3:C:415:THR:HG22	3:C:464:ILE:HG21	1.69	0.73
1:A:251:MSE:HE2	2:B:313:ILE:HD11	1.72	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	588/778 (76%)	576 (98%)	7 (1%)	5 (1%)	14 48
2	B	560/718 (78%)	550 (98%)	6 (1%)	4 (1%)	18 53
3	C	402/455 (88%)	395 (98%)	7 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1550/1951 (79%)	1521 (98%)	20 (1%)	9 (1%)	21	56

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	340	VAL
1	A	341	LEU
2	B	141	GLU
1	A	389	VAL
1	A	193	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	446/660 (68%)	446 (100%)	0	100	100
2	B	472/607 (78%)	472 (100%)	0	100	100
3	C	295/406 (73%)	295 (100%)	0	100	100
All	All	1213/1673 (72%)	1213 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	302	ASN
3	C	373	ASN
2	B	77	GLN
2	B	92	ASN
2	B	109	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	591/778 (75%)	0.63	53 (8%) 15 8	39, 83, 220, 262	0
2	B	574/718 (79%)	0.02	15 (2%) 57 34	30, 59, 110, 139	0
3	C	410/455 (90%)	1.49	102 (24%) 2 1	60, 142, 220, 246	0
All	All	1575/1951 (80%)	0.63	170 (10%) 11 6	30, 84, 214, 262	0

The worst 5 of 170 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	361	GLY	4.9
3	C	295	GLU	4.8
3	C	465	ILE	4.8
3	C	127	ALA	4.7
1	A	569	ALA	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.