



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:13 AM UTC

PDB ID : 4BGA / pdb_00004bga
Title : Nucleotide-bound open form of a putative sugar kinase MK0840 from
Methanopyrus kandleri
Authors : Schacherl, M.; Baumann, U.
Deposited on : 2013-03-24
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	
1	B	325	
1	C	325	
1	D	325	
2	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PREDICTED MOLECULAR CHAPERONE DISTANTLY RELATED TO HSP70-F OLD METALLOPROTEASES.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2390	1500	416	464	10			
1	B	314	Total	C	N	O	S	0	1	0
			2396	1503	417	465	11			
1	C	322	Total	C	N	O	S	0	1	0
			2447	1531	429	476	11			
1	D	314	Total	C	N	O	S	0	1	0
			2396	1503	417	465	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	GLY	-	expression tag	UNP Q8TX37
A	35	SER	-	expression tag	UNP Q8TX37
A	36	HIS	-	expression tag	UNP Q8TX37
B	34	GLY	-	expression tag	UNP Q8TX37
B	35	SER	-	expression tag	UNP Q8TX37
B	36	HIS	-	expression tag	UNP Q8TX37
C	34	GLY	-	expression tag	UNP Q8TX37
C	35	SER	-	expression tag	UNP Q8TX37
C	36	HIS	-	expression tag	UNP Q8TX37
D	34	GLY	-	expression tag	UNP Q8TX37
D	35	SER	-	expression tag	UNP Q8TX37
D	36	HIS	-	expression tag	UNP Q8TX37

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	K	0	0
			2	2		
3	B	2	Total	K	0	0
			2	2		
3	C	2	Total	K	0	0
			2	2		
3	D	2	Total	K	0	0
			2	2		

- Molecule 4 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	B	1	Total	C	O	0	0
			12	6	6		
4	C	1	Total	C	O	0	0
			12	6	6		
4	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
5	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		
6	D	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	72	Total	O	0	0
			72	72		
7	B	61	Total	O	0	0
			61	61		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	51	Total	O	0	0
			51	51		
7	D	38	Total	O	0	0
			38	38		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PREDICTED MOLECULAR CHAPERONE DISTANTLY RELATED TO HSP70-F OLD METALLOPROTEASES




- Molecule 1: PREDICTED MOLECULAR CHAPERONE DISTANTLY RELATED TO HSP70-F OLD METALLOPROTEASES




- Molecule 1: PREDICTED MOLECULAR CHAPERONE DISTANTLY RELATED TO HSP70-F OLD METALLOPROTEASES




- Molecule 1: PREDICTED MOLECULAR CHAPERONE DISTANTLY RELATED TO HSP70-F OLD METALLOPROTEASES





- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



- Molecule 2: beta-D-fructofuranose-(2-1)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.57Å 111.67Å 87.88Å 90.00° 89.55° 90.00°	Depositor
Resolution (Å)	73.57 – 2.60 73.57 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (73.57-2.60) 100.0 (73.57-2.60)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.48Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.154 , 0.208 (Not available) , 0.182	Depositor DCC
R_{free} test set	2188 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10064	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FRU, MG, ADP, GLC, K, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2431	0.85	1/3304 (0.0%)
1	B	0.44	0/2437	0.83	1/3312 (0.0%)
1	C	0.42	0/2488	0.88	4/3377 (0.1%)
1	D	0.44	0/2437	0.83	0/3312
All	All	0.43	0/9793	0.85	6/13305 (0.0%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	248	PRO	N-CA-C	6.73	118.92	110.70
1	C	204	LYS	N-CA-C	5.79	118.40	110.24
1	B	248	PRO	N-CA-C	5.60	117.54	110.70
1	C	114	ASP	N-CA-C	5.34	117.10	111.28
1	C	321	ALA	N-CA-C	-5.25	103.76	110.53
1	A	248	PRO	N-CA-C	5.02	116.82	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2390	0	2375	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2396	0	2379	26	0
1	C	2447	0	2431	32	0
1	D	2396	0	2379	26	0
2	E	23	0	21	2	0
2	F	23	0	21	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	12	0	12	0	0
4	B	12	0	12	1	0
4	C	12	0	12	2	0
4	D	12	0	12	0	0
5	A	27	0	12	1	0
5	B	27	0	12	0	0
5	C	27	0	12	1	0
5	D	27	0	12	1	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	72	0	0	0	0
7	B	61	0	0	0	0
7	C	51	0	0	1	0
7	D	38	0	0	0	0
All	All	10064	0	9702	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:GLU:OE2	1:C:101:ARG:NH1	2.12	0.82
1:A:179:ARG:NH2	1:A:341:GLU:OE1	2.15	0.80
1:B:70:VAL:HA	1:B:73:LYS:HE2	1.65	0.78
1:C:257:ASP:OD1	1:C:260:ARG:NH2	2.22	0.73
1:C:351:GLY:H	1:C:358:THR:HG21	1.55	0.70
1:B:233:GLY:HA3	1:C:202:LEU:HA	1.76	0.66
1:D:308:ARG:NH1	1:D:311:GLU:OE1	2.29	0.65
1:C:239:ARG:O	1:C:241:ARG:NH1	2.29	0.62
1:C:63:ILE:HG13	1:C:324:GLU:HG3	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:GLU:HB2	2:E:2:FRU:H5	1.82	0.61
1:B:250:GLU:HG2	1:B:253:ARG:HH21	1.66	0.61
1:B:231:VAL:HG11	1:B:250:GLU:HB3	1.84	0.60
1:C:85:ARG:NH1	7:C:2011:HOH:O	2.35	0.60
1:C:204:LYS:N	1:C:204:LYS:HD3	2.17	0.60
1:C:208:GLU:O	1:C:211:LYS:HG2	2.02	0.59
1:C:169:MET:HG3	4:C:1361:BGC:H6C1	1.85	0.59
1:B:174:PRO:HG3	1:B:184:ASP:HB3	1.86	0.57
1:B:63:ILE:HG13	1:B:324:GLU:HG3	1.87	0.56
1:C:204:LYS:HD3	1:C:204:LYS:H	1.71	0.55
1:D:223:ARG:O	1:D:227:ARG:HG2	2.08	0.54
1:A:54:VAL:HG23	1:A:64:VAL:HG11	1.88	0.54
1:C:68:GLU:HG2	2:E:1:GLC:O6	2.07	0.54
1:C:231:VAL:HG11	1:C:250:GLU:HB3	1.89	0.54
1:B:252:GLU:OE2	1:B:256:ARG:NH2	2.42	0.53
1:C:202:LEU:HD11	1:C:212:LYS:HE3	1.90	0.53
1:A:252:GLU:O	1:A:256:ARG:HB2	2.08	0.53
1:D:80:LEU:O	1:D:84:ARG:HB2	2.08	0.53
1:D:112:ILE:HG13	1:D:113:VAL:HG23	1.91	0.53
1:C:149:GLY:O	1:C:329:GLY:HA3	2.09	0.52
1:B:37:LEU:HA	1:B:57:GLU:OE2	2.09	0.52
1:C:252:GLU:O	1:C:256:ARG:HB2	2.09	0.52
1:A:197:ALA:HA	1:A:259:ILE:HG21	1.92	0.52
1:D:300:VAL:HG13	1:D:306:LYS:HG3	1.92	0.51
1:A:195:ASP:OD2	5:A:1354:ADP:O3'	2.27	0.51
1:B:76:CYS:SG	1:D:84:ARG:NH1	2.83	0.51
1:D:149:GLY:O	1:D:329:GLY:HA3	2.11	0.51
1:A:149:GLY:O	1:A:329:GLY:HA3	2.10	0.51
1:A:212:LYS:HE2	1:A:267:GLU:HB3	1.93	0.51
1:C:162:VAL:HG23	1:C:290:MET:HB2	1.92	0.50
1:B:112:ILE:HG13	1:B:113:VAL:HG23	1.93	0.50
1:B:149:GLY:O	1:B:329:GLY:HA3	2.11	0.50
1:D:195:ASP:OD2	5:D:1356:ADP:O3'	2.24	0.50
1:C:169:MET:CG	4:C:1361:BGC:H6C1	2.41	0.50
1:A:179:ARG:NH2	1:A:336:ARG:HH11	2.10	0.50
1:D:240:VAL:HG22	1:D:245:ALA:HB2	1.94	0.49
1:D:252:GLU:HG2	1:D:256:ARG:HD3	1.95	0.48
1:C:63:ILE:CG1	1:C:324:GLU:HG3	2.43	0.48
1:C:169:MET:HE3	1:C:187:VAL:HG12	1.95	0.48
1:D:212:LYS:HE3	1:D:267:GLU:HB3	1.96	0.48
1:C:195:ASP:OD2	5:C:1364:ADP:O3'	2.30	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:37:LEU:HA	1:D:57:GLU:OE2	2.14	0.47
1:D:178:GLY:O	1:D:336:ARG:NH2	2.47	0.47
1:B:63:ILE:CG1	1:B:324:GLU:HG3	2.44	0.47
1:A:310:SER:HA	1:A:317:VAL:HG21	1.96	0.47
1:A:174:PRO:HG2	1:A:181:ASP:HB3	1.96	0.47
1:C:221:ASN:HD22	1:C:224:ARG:NH2	2.13	0.47
1:D:160:ASN:HA	1:D:175:ILE:O	2.15	0.47
1:A:37:LEU:HA	1:A:57:GLU:OE2	2.15	0.46
1:B:123:MET:HG3	1:D:139:PRO:HB2	1.97	0.46
1:C:37:LEU:HA	1:C:57:GLU:OE2	2.15	0.46
1:D:341:GLU:HG2	1:D:342:PRO:HD2	1.98	0.46
1:A:194:LEU:HD23	1:A:273:LEU:HB3	1.97	0.46
1:C:351:GLY:H	1:C:358:THR:CG2	2.26	0.46
1:C:50:THR:OG1	1:C:94:ARG:NH2	2.50	0.45
1:B:233:GLY:CA	1:C:202:LEU:HA	2.46	0.45
1:A:152:TYR:OH	1:A:179:ARG:NH1	2.49	0.45
1:C:178:GLY:O	1:C:336:ARG:NH2	2.46	0.45
1:D:252:GLU:O	1:D:256:ARG:HB2	2.17	0.44
1:D:131:PRO:HB3	1:D:349:GLU:HG2	1.98	0.44
1:A:87:LEU:HD11	1:A:120:LEU:HG	1.99	0.44
1:B:80:LEU:HD23	1:D:113:VAL:HG11	1.99	0.44
1:B:74:ILE:HB	1:B:116:LEU:HD13	2.00	0.44
1:A:112:ILE:HG13	1:A:113:VAL:HG23	2.00	0.43
1:A:44:GLN:O	1:A:50:THR:HA	2.19	0.42
1:A:290:MET:HE3	1:A:290:MET:HB2	1.91	0.42
1:C:282:VAL:HG21	1:C:312:LEU:HB2	2.01	0.42
1:A:131:PRO:HB3	1:A:349:GLU:HG2	2.02	0.42
1:D:216:GLY:HA3	1:D:267:GLU:O	2.20	0.42
1:A:234:LYS:HA	1:A:235:PRO:HD3	1.89	0.41
1:B:346:LEU:HD21	1:D:122:PRO:HG3	2.02	0.41
1:A:314:GLU:H	1:A:314:GLU:HG3	1.67	0.41
1:C:148:VAL:HA	1:C:182:PHE:CZ	2.55	0.41
1:D:150:ALA:HB3	1:D:151:PRO:HD3	2.02	0.41
1:A:198:ALA:O	1:A:202:LEU:HB2	2.20	0.41
1:B:313:TRP:CD1	1:B:315:GLY:H	2.38	0.41
1:C:307:ARG:HG2	1:C:311:GLU:OE2	2.20	0.41
1:D:179:ARG:HE	1:D:179:ARG:HB3	1.58	0.41
1:B:79:THR:HA	1:D:79:THR:HA	2.02	0.41
1:B:80:LEU:HB3	1:D:76:CYS:HA	2.03	0.41
1:B:187:VAL:HG11	4:B:1353:BGC:H6C2	2.02	0.41
1:B:44:GLN:O	1:B:50:THR:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:250:GLU:HG2	1:C:253:ARG:NH2	2.36	0.41
1:A:169:MET:HE3	1:A:187:VAL:HG12	2.01	0.41
1:B:336:ARG:HD3	1:B:341:GLU:CD	2.46	0.41
1:D:162:VAL:HG23	1:D:290:MET:HB2	2.03	0.41
1:A:122:PRO:O	1:A:126:GLU:HB2	2.21	0.40
1:C:265:PRO:HB2	1:C:267:GLU:OE1	2.21	0.40
1:B:209:GLY:HA3	1:B:211:LYS:NZ	2.36	0.40
1:B:154:VAL:HG21	1:B:163:HIS:CD2	2.56	0.40
1:B:174:PRO:HG2	1:B:181:ASP:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/325 (93%)	300 (99%)	3 (1%)	0	100	100
1	B	313/325 (96%)	309 (99%)	4 (1%)	0	100	100
1	C	321/325 (99%)	315 (98%)	6 (2%)	0	100	100
1	D	313/325 (96%)	307 (98%)	6 (2%)	0	100	100
All	All	1250/1300 (96%)	1231 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/259 (98%)	243 (96%)	10 (4%)	28	55
1	B	254/259 (98%)	251 (99%)	3 (1%)	63	83
1	C	258/259 (100%)	253 (98%)	5 (2%)	50	75
1	D	254/259 (98%)	252 (99%)	2 (1%)	73	88
All	All	1019/1036 (98%)	999 (98%)	20 (2%)	48	74

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	VAL
1	A	202	LEU
1	A	223	ARG
1	A	231	VAL
1	A	232	ASP
1	A	234	LYS
1	A	283	ILE
1	A	290	MET
1	A	300	VAL
1	A	314	GLU
1	B	153	CYS
1	B	179	ARG
1	B	184	ASP
1	C	62	GLU
1	C	201	LEU
1	C	202	LEU
1	C	204	LYS
1	C	253	ARG
1	D	64	VAL
1	D	169	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	221	ASN
1	C	221	ASN
1	D	251	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	E	1	2	11,11,12	0.54	0	15,15,17	0.92	1 (6%)
2	FRU	E	2	2	11,12,12	0.54	0	10,18,18	0.72	0
2	GLC	F	1	2	11,11,12	0.65	0	15,15,17	0.47	0
2	FRU	F	2	2	11,12,12	0.52	0	10,18,18	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	0/2/19/22	0/1/1/1
2	FRU	E	2	2	-	5/5/24/24	0/1/1/1
2	GLC	F	1	2	-	0/2/19/22	0/1/1/1
2	FRU	F	2	2	-	2/5/24/24	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	GLC	C1-O5-C5	2.73	115.84	112.19

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	FRU	O1-C1-C2-C3
2	E	2	FRU	O1-C1-C2-O2
2	E	2	FRU	O1-C1-C2-O5
2	F	2	FRU	C4-C5-C6-O6
2	F	2	FRU	O5-C5-C6-O6
2	E	2	FRU	O5-C5-C6-O6
2	E	2	FRU	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1	GLC	1	0
2	E	2	FRU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 11 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ADP	A	1354	-	28,29,29	1.41	6 (21%)	43,45,45	1.76	8 (18%)
5	ADP	B	1355	6	28,29,29	1.36	5 (17%)	43,45,45	1.86	11 (25%)
4	BGC	A	1353	-	12,12,12	1.49	1 (8%)	17,17,17	1.26	2 (11%)
5	ADP	D	1356	6	28,29,29	1.42	4 (14%)	43,45,45	1.90	10 (23%)
4	BGC	B	1353	-	12,12,12	1.54	3 (25%)	17,17,17	1.34	4 (23%)
4	BGC	C	1361	-	12,12,12	1.53	1 (8%)	17,17,17	1.17	1 (5%)
5	ADP	C	1364	6	28,29,29	1.41	5 (17%)	43,45,45	1.84	10 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BGC	D	1353	-	12,12,12	1.54	2 (16%)	17,17,17	1.38	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	A	1354	-	-	1/16/32/32	0/3/3/3
5	ADP	B	1355	6	-	0/16/32/32	0/3/3/3
4	BGC	A	1353	-	-	2/2/22/22	0/1/1/1
5	ADP	D	1356	6	-	5/16/32/32	0/3/3/3
4	BGC	B	1353	-	-	2/2/22/22	0/1/1/1
4	BGC	C	1361	-	-	1/2/22/22	0/1/1/1
5	ADP	C	1364	6	-	3/16/32/32	0/3/3/3
4	BGC	D	1353	-	-	2/2/22/22	0/1/1/1

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	1356	ADP	C5-C4	4.77	1.47	1.39
5	C	1364	ADP	C5-C4	4.63	1.47	1.39
5	A	1354	ADP	C5-C4	4.59	1.47	1.39
5	B	1355	ADP	C5-C4	4.52	1.47	1.39
4	C	1361	BGC	C4-C3	-3.92	1.42	1.52
4	D	1353	BGC	C4-C3	-3.91	1.42	1.52
4	B	1353	BGC	C4-C3	-3.84	1.42	1.52
4	A	1353	BGC	C4-C3	-3.71	1.42	1.52
5	D	1356	ADP	C5-C6	2.78	1.48	1.41
5	C	1364	ADP	C5-C6	2.70	1.48	1.41
5	A	1354	ADP	C5-C6	2.59	1.48	1.41
5	B	1355	ADP	C5-C6	2.57	1.48	1.41
5	D	1356	ADP	C8-N7	2.39	1.36	1.31
5	B	1355	ADP	C5-N7	-2.38	1.34	1.39
5	A	1354	ADP	C5-N7	-2.32	1.34	1.39
5	C	1364	ADP	C5-N7	-2.31	1.34	1.39
5	D	1356	ADP	C5-N7	-2.24	1.35	1.39
5	B	1355	ADP	C8-N7	2.20	1.35	1.31
4	B	1353	BGC	C3-C2	-2.20	1.46	1.52
5	C	1364	ADP	PA-O3A	2.14	1.61	1.59
5	A	1354	ADP	C4-N9	-2.11	1.33	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1364	ADP	C8-N7	2.10	1.35	1.31
4	B	1353	BGC	O5-C5	2.09	1.49	1.44
5	A	1354	ADP	C8-N7	2.08	1.35	1.31
5	B	1355	ADP	C4-N9	-2.07	1.33	1.37
5	A	1354	ADP	PA-O3A	2.06	1.61	1.59
4	D	1353	BGC	O5-C5	2.03	1.49	1.44

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1364	ADP	C5-C4-N3	-5.83	118.68	126.72
5	D	1356	ADP	C5-C4-N3	-5.83	118.69	126.72
5	B	1355	ADP	C5-C4-N3	-5.68	118.90	126.72
5	A	1354	ADP	C5-C4-N3	-5.46	119.20	126.72
5	C	1364	ADP	N3-C4-N9	4.65	135.07	127.17
5	D	1356	ADP	N3-C4-N9	4.61	135.01	127.17
5	A	1354	ADP	N3-C4-N9	4.52	134.86	127.17
5	B	1355	ADP	N3-C4-N9	4.50	134.83	127.17
5	D	1356	ADP	C2-N3-C4	3.72	120.92	111.83
5	C	1364	ADP	C2-N3-C4	3.68	120.81	111.83
5	B	1355	ADP	C2-N3-C4	3.67	120.79	111.83
5	C	1364	ADP	C4-C5-N7	-3.63	106.43	110.58
5	A	1354	ADP	C2-N3-C4	3.62	120.67	111.83
5	D	1356	ADP	C4-C5-N7	-3.54	106.53	110.58
5	A	1354	ADP	N3-C2-N1	-3.39	123.45	128.58
5	B	1355	ADP	C4-C5-N7	-3.37	106.73	110.58
5	B	1355	ADP	N3-C2-N1	-3.31	123.58	128.58
5	C	1364	ADP	N3-C2-N1	-3.24	123.68	128.58
4	D	1353	BGC	O5-C1-C2	3.19	115.91	110.30
5	D	1356	ADP	N3-C2-N1	-3.18	123.77	128.58
5	D	1356	ADP	C2'-C1'-N9	-3.14	105.51	113.30
5	A	1354	ADP	C4-C5-N7	-3.08	107.06	110.58
5	B	1355	ADP	C2'-C1'-N9	-3.02	105.80	113.30
5	A	1354	ADP	C4-N9-C8	2.82	108.70	105.74
5	D	1356	ADP	C4-N9-C8	2.78	108.65	105.74
5	C	1364	ADP	C4-N9-C8	2.73	108.60	105.74
5	C	1364	ADP	C5-N7-C8	2.68	107.66	103.45
5	B	1355	ADP	C4-N9-C8	2.67	108.54	105.74
5	D	1356	ADP	C5-N7-C8	2.60	107.54	103.45
4	B	1353	BGC	O6-C6-C5	2.54	120.00	111.33
4	A	1353	BGC	O5-C5-C6	2.44	112.48	106.44
5	B	1355	ADP	C5-N7-C8	2.41	107.23	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1353	BGC	O5-C5-C6	2.40	112.38	106.44
4	C	1361	BGC	O5-C1-C2	2.35	114.43	110.30
4	B	1353	BGC	O5-C1-C2	2.34	114.41	110.30
5	A	1354	ADP	C5-N7-C8	2.26	107.00	103.45
5	C	1364	ADP	C2'-C1'-N9	-2.22	107.79	113.30
5	D	1356	ADP	C6-C5-N7	2.17	136.27	132.09
4	A	1353	BGC	O5-C1-C2	2.17	114.11	110.30
4	D	1353	BGC	O5-C5-C6	2.17	111.81	106.44
5	B	1355	ADP	C6-C5-N7	2.15	136.24	132.09
5	B	1355	ADP	O3B-PB-O2B	2.11	115.70	107.80
5	B	1355	ADP	O2A-PA-O1A	2.09	122.17	112.44
5	D	1356	ADP	N9-C8-N7	-2.09	110.97	113.94
4	B	1353	BGC	O5-C5-C4	2.03	113.37	109.70
5	A	1354	ADP	C6-C5-N7	2.03	136.01	132.09
5	C	1364	ADP	C6-C5-N7	2.03	136.00	132.09
5	C	1364	ADP	N9-C8-N7	-2.03	111.06	113.94

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	1364	ADP	C5'-O5'-PA-O3A
5	D	1356	ADP	C5'-O5'-PA-O1A
5	D	1356	ADP	O4'-C4'-C5'-O5'
4	D	1353	BGC	O5-C5-C6-O6
5	D	1356	ADP	C3'-C4'-C5'-O5'
4	D	1353	BGC	C4-C5-C6-O6
4	A	1353	BGC	C4-C5-C6-O6
4	A	1353	BGC	O5-C5-C6-O6
4	B	1353	BGC	C4-C5-C6-O6
4	B	1353	BGC	O5-C5-C6-O6
5	C	1364	ADP	C5'-O5'-PA-O1A
5	C	1364	ADP	C5'-O5'-PA-O2A
5	D	1356	ADP	C5'-O5'-PA-O2A
5	D	1356	ADP	C5'-O5'-PA-O3A
4	C	1361	BGC	C4-C5-C6-O6
5	A	1354	ADP	PB-O3A-PA-O1A



There are no ring outliers.


5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1354	ADP	1	0
5	D	1356	ADP	1	0
4	B	1353	BGC	1	0
4	C	1361	BGC	2	0
5	C	1364	ADP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/325 (96%)	-0.70	1 (0%) 90 88	16, 27, 50, 58	0
1	B	314/325 (96%)	-0.73	2 (0%) 85 83	12, 27, 51, 63	1 (0%)
1	C	322/325 (99%)	-0.43	0 100 100	12, 32, 65, 82	1 (0%)
1	D	314/325 (96%)	-0.57	2 (0%) 85 83	14, 30, 55, 71	1 (0%)
All	All	1264/1300 (97%)	-0.61	5 (0%) 88 86	12, 29, 56, 82	3 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	350	GLY	3.1
1	A	350	GLY	2.8
1	B	233	GLY	2.6
1	D	350	GLY	2.3
1	D	224	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	E	1	11/12	0.90	0.09	36,48,56,56	0
2	GLC	F	1	11/12	0.90	0.09	36,52,56,60	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FRU	E	2	12/12	0.91	0.09	43,49,57,66	0
2	FRU	F	2	12/12	0.94	0.07	43,45,55,58	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BGC	A	1353	12/12	0.86	0.12	36,46,53,53	0
4	BGC	D	1353	12/12	0.86	0.12	46,55,70,77	0
4	BGC	C	1361	12/12	0.87	0.12	41,61,68,70	0
4	BGC	B	1353	12/12	0.87	0.11	42,50,59,60	0
6	MG	C	1363	1/1	0.88	0.18	55,55,55,55	0
5	ADP	C	1364	27/27	0.89	0.10	32,45,98,109	0
6	MG	D	1355	1/1	0.90	0.23	52,52,52,52	0
5	ADP	B	1355	27/27	0.92	0.08	22,28,72,99	0
5	ADP	D	1356	27/27	0.92	0.09	30,43,75,90	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	ADP	A	1354	27/27	0.93	0.07	20,30,83,85	0
6	MG	B	1354	1/1	0.94	0.13	50,50,50,50	0
3	K	D	1352	1/1	0.95	0.06	46,46,46,46	0
3	K	A	1351	1/1	0.97	0.05	48,48,48,48	0
3	K	C	1359	1/1	0.97	0.05	47,47,47,47	0
3	K	A	1352	1/1	0.99	0.02	27,27,27,27	0
3	K	C	1360	1/1	0.99	0.02	28,28,28,28	0
3	K	D	1351	1/1	0.99	0.03	29,29,29,29	0
3	K	B	1351	1/1	0.99	0.04	33,33,33,33	0
3	K	B	1352	1/1	0.99	0.04	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ADP B 1355:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP D 1356:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADP A 1354:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.