



wwPDB EM Validation Summary Report ⓘ

Mar 23, 2026 – 06:18 AM UTC

PDB ID : 7BG7 / pdb_00007bg7
EMDB ID : EMD-12172
Title : HRV14 in complex with its receptor ICAM-1
Authors : Hrebik, D.; Fuzik, T.; Plevka, P.
Deposited on : 2021-01-06
Resolution : 2.40 Å (reported)
Based on initial model : 4RHV

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	5628 (1.90 - 2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	293	
2	2	262	
3	3	236	
4	4	68	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	B	453	17% . 81%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	1	274	Total	C	N	O	S	0	0
			2179	1378	377	416	8		

- Molecule 2 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	2	250	Total	C	N	O	S	0	0
			1911	1214	322	363	12		

- Molecule 3 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	3	236	Total	C	N	O	S	0	0
			1849	1184	305	353	7		

- Molecule 4 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	4	41	Total	C	N	O	S	0	0
			307	195	47	63	2		

- Molecule 5 is a protein called Intercellular adhesion molecule 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	B	84	Total	C	N	O	S	0	0
			641	403	106	127	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		AltConf
6	1	152	Total	O	0
			152	152	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
6	2	118	Total 118	O 118	0
6	3	129	Total 129	O 129	0
6	4	13	Total 13	O 13	0
6	B	33	Total 33	O 33	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Genome polyprotein




- Molecule 2: Genome polyprotein



- Molecule 3: Genome polyprotein



- Molecule 4: Genome polyprotein



- Molecule 5: Intercellular adhesion molecule 1



THR	TYR	GLY	TYR	ALA	ALA	LEU	GLU	Q1
	LEU	PRO	VAL	HIS	SER	PRO	LYS	L11
	CYS	ARG	ARG	VAL	SER	THR	LEU	T23
	ARG	LEU	ALA	LYS	THR	PRO	ARG	S24
	ARG	GLU	THR	VAL	ALA	GLN	GLU	C25
	SER	ARG	THR	VAL	GLU	LEU	PRO	D26
	THR	ASP	LEU	LEU	ASP	VAL	ALA	L31
	GLN	CYS	ASN	GLY	GLY	SER	VAL	E41
	GLY	PRO	VAL	GLY	THR	PRO	GLY	V57
	GLU	GLY	VAL	VAL	THR	ARG	GLU	C69
THR	GLU	GLY	GLY	GLY	GLY	SER	VAL	W84
	VAL	ASN	PRO	PRO	THR	ARG	PRO	THR
	ASN	SER	ILE	VAL	GLN	THR	THR	PRO
	VAL	SER	ALA	ARG	LEU	THR	VAL	GLU
	VAL	GLN	GLN	GLY	GLY	VAL	VAL	ARG
	SER	THR	LEU	ASN	VAL	ARG	VAL	VAL
	PRO	PRO	LEU	GLN	CYS	ARG	GLU	GLU
	ARG	MET	LEU	LEU	SER	ASP	LEU	LEU
	TYR	GLN	GLN	LYS	GLN	HIS	HIS	ALA
	GLU	ALA	ALA	THR	THR	GLY	GLY	PRO
THR	TRP	ALA	PRO	PRO	LEU	LEU	ALA	PRO
	GLY	GLY	GLU	GLN	PHE	ASN	ASN	SER
	ASN	ASN	ASP	THR	PRO	PHE	THR	TRP
	PRO	PRO	ASN	VAL	VAL	GLN	GLN	PRO
	LEU	LEU	GLY	THR	SER	CYS	CYS	PRO
	PRO	PRO	ARG	ILE	GLU	ARG	VAL	VAL
	GLU	GLU	SER	TYR	ALA	THR	GLY	GLY
	LEU	LEU	PHE	SER	GLN	GLU	LYS	LYS
	LYS	LYS	SER	PHE	VAL	ASN	ASN	ASN
	CYS	CYS	CYS	PRO	HIS	VAL	LEU	LEU
THR	LEU	LEU	SER	ALA	ALA	LEU	LEU	THR
	LYS	LYS	ALA	PRO	PRO	ALA	ARG	LEU
	ASP	ASP	THR	THR	LEU	GLY	PRO	CYS
	GLY	GLY	GLU	VAL	GLY	ASP	GLN	GLN
	THR	PHE	THR	ILE	ILE	GLY	GLY	GLN
	PRO	PRO	ALA	THR	ARG	GLU	GLU	VAL
	LEU	LEU	GLY	LYS	ASN	PHE	THR	GLY
	PRO	PRO	GLN	PRO	ASN	ASN	GLU	ALA
	ILE	ILE	LEU	GLU	PRO	THR	PRO	PRO
	GLY	GLY	GLU	VAL	THR	ASN	ALA	ALA
THR	THR	THR	THR	THR	THR	THR	THR	THR
	ARG	ARG	ARG	VAL	ASP	GLN	GLN	VAL
	ASP	ASP	LEU	VAL	PHE	GLN	GLN	LEU
	LEU	LEU	ARG	VAL	SER	THR	THR	LEU
	GLY	GLY	VAL	GLN	LYS	THR	THR	ARG
	GLY	GLY	THR	THR	GLY	GLY	GLY	GLY
	GLU	GLU	THR	THR	GLU	GLY	GLY	GLY
	VAL	VAL	THR	THR	GLU	THR	THR	THR
	VAL	VAL	THR	THR	VAL	THR	THR	THR
	VAL	VAL	THR	THR	VAL	THR	THR	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	12085	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	83.6	Depositor
Minimum defocus (nm)	420	Depositor
Maximum defocus (nm)	3562	Depositor
Magnification	75000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.667	Depositor
Minimum map value	-0.356	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.041	Depositor
Recommended contour level	0.055	Depositor
Map size (Å)	403.93997, 403.93997, 403.93997	wwPDB
Map dimensions	380, 380, 380	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.063, 1.063, 1.063	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.27	0/2237	0.42	0/3043
2	2	0.25	0/1959	0.38	0/2679
3	3	0.27	0/1898	0.41	0/2597
4	4	0.55	1/313 (0.3%)	0.90	2/423 (0.5%)
5	B	0.22	0/654	0.36	0/891
All	All	0.28	1/7061 (0.0%)	0.43	2/9633 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	4	59	LEU	CA-C	5.99	1.60	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	4	61	LEU	N-CA-C	-6.71	105.07	112.72
4	4	62	LYS	CA-C-O	-5.36	115.62	121.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2179	0	2115	30	0
2	2	1911	0	1893	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	3	1849	0	1833	24	0
4	4	307	0	304	5	0
5	B	641	0	647	6	0
6	1	152	0	0	7	0
6	2	118	0	0	8	0
6	3	129	0	0	6	0
6	4	13	0	0	0	0
6	B	33	0	0	3	0
All	All	7332	0	6792	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 71 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:282:ARG:NH2	3:3:82:GLY:O	2.03	0.92
5:B:41:GLU:OE1	6:B:501:HOH:O	1.93	0.87
5:B:1:GLN:N	5:B:24:SER:OG	2.09	0.85
1:1:91:ASP:OD1	6:1:301:HOH:O	1.94	0.84
3:3:41:HIS:HB3	4:4:44:SER:OG	1.79	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	272/293 (93%)	266 (98%)	6 (2%)	0	100	100
2	2	248/262 (95%)	237 (96%)	11 (4%)	0	100	100
3	3	234/236 (99%)	224 (96%)	10 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	4	39/68 (57%)	38 (97%)	1 (3%)	0	100	100
5	B	82/453 (18%)	81 (99%)	1 (1%)	0	100	100
All	All	875/1312 (67%)	846 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	240/256 (94%)	240 (100%)	0	100	100
2	2	219/229 (96%)	219 (100%)	0	100	100
3	3	209/209 (100%)	208 (100%)	1 (0%)	81	91
4	4	35/57 (61%)	34 (97%)	1 (3%)	37	60
5	B	78/398 (20%)	78 (100%)	0	100	100
All	All	781/1149 (68%)	779 (100%)	2 (0%)	84	93

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	3	101	VAL
4	4	60	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	2	111	GLN
2	2	217	ASN
3	3	204	GLN
2	2	262	GLN
1	1	169	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12172. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 190




Y Index: 190




Z Index: 190


6.2.2 Raw map



X Index: 273



Y Index: 273




Z Index: 273


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 206




Y Index: 174




Z Index: 206


6.3.2 Raw map



X Index: 249



Y Index: 257




Z Index: 255


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X



Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.055. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_12172_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 3382 nm³; this corresponds to an approximate mass of 3055 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.417 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.417 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.40	-	-
Author-provided FSC curve	2.39	2.75	2.43
Unmasked-calculated*	2.49	2.88	2.53


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12172 and PDB model 7BG7. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlays


9.1.1 Map-model overlay [i](#)



X




Y




Z


9.1.2 Map-model assembly overlay [i](#)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.055 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.055).

9.4 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.055) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9340	0.6960
1	0.9390	0.6970
2	0.9570	0.7090
3	0.9610	0.7110
4	0.7210	0.6030
B	0.8860	0.6610

