



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:36 AM UTC

PDB ID : 2BHY / pdb_00002bhy
Title : Crystal structure of Deinococcus radiodurans maltooligosyltrehalose trehalohydrolase in complex with trehalose
Authors : Timmins, J.; Leiros, H.-K.S.; Leonard, G.; Leiros, I.; McSweeney, S.
Deposited on : 2005-01-20
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	602	88% 7% ••
2	B	2	100%
2	C	2	50% 50%
2	D	2	50% 50%
2	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	
2	G	2	
2	H	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GLC	A	705	X	-	-	-

2 Entry composition


There are 7 unique types of molecules in this entry. The entry contains 6219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MALTOOLIGOSYLTREHALOSE TREHALOHYDROLASE.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	580	Total	C	N	O	S	Se	35	48	0
			5047	3192	902	942	2	9			

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			
2	E	2	Total	C	O	0	0	0
			23	12	11			
2	F	2	Total	C	O	0	0	0
			23	12	11			
2	G	2	Total	C	O	0	0	0
			23	12	11			
2	H	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			16	8	2	6		
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is alpha-D-glucopyranose (CCD ID: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			12	6	6		
5	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	958	Total	O	0	0
			958	958		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MALTOOLIGOSYLTREHALOSE TREHALOHYDROLASE



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain F:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain G:  50% 50%



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose

Chain H:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.03Å 66.56Å 153.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.70 – 1.50 76.57 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (76.70-1.50) 99.0 (76.57-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.128 , 0.152 (Not available) , 0.173	Depositor DCC
R_{free} test set	2431 reflections (2.48%)	wwPDB-VP
Wilson B-factor (Å ²)	10.2	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 64.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6219	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BME, GLC, MG, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.53	17/5177 (0.3%)	1.31	19/7028 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	503[A]	GLU	CD-OE1	77.81	2.73	1.25
1	A	503[B]	GLU	CD-OE1	77.81	2.73	1.25
1	A	503[A]	GLU	CD-OE2	76.44	2.70	1.25
1	A	503[B]	GLU	CD-OE2	76.44	2.70	1.25
1	A	466	GLU	CD-OE2	62.19	2.43	1.25
1	A	503[A]	GLU	CG-CD	28.57	2.23	1.52
1	A	503[B]	GLU	CG-CD	28.57	2.23	1.52
1	A	22[A]	ARG	CD-NE	-22.91	1.14	1.46
1	A	22[B]	ARG	CD-NE	-22.91	1.14	1.46
1	A	580	ARG	CZ-NH1	21.13	1.62	1.32
1	A	579	ARG	CD-NE	12.48	1.63	1.46
1	A	46	ARG	CG-CD	11.82	1.88	1.52
1	A	482	ASP	CG-OD1	-9.39	1.07	1.25
1	A	570	GLU	CD-OE2	-8.27	1.09	1.25
1	A	243[A]	MSE	SE-CE	-5.30	1.79	1.95
1	A	243[B]	MSE	SE-CE	-5.30	1.79	1.95
1	A	196	MSE	SE-CE	-5.04	1.80	1.95

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	462	GLN	OE1-CD-NE2	-38.09	84.51	122.60
1	A	503[A]	GLU	OE1-CD-OE2	-37.18	33.66	122.90
1	A	503[B]	GLU	OE1-CD-OE2	-37.18	33.66	122.90
1	A	503[A]	GLU	CG-CD-OE2	-27.94	54.14	118.40
1	A	503[B]	GLU	CG-CD-OE2	-27.94	54.14	118.40
1	A	503[A]	GLU	CG-CD-OE1	-27.00	56.29	118.40
1	A	503[B]	GLU	CG-CD-OE1	-27.00	56.29	118.40
1	A	462	GLN	CG-CD-NE2	23.28	151.32	116.40
1	A	466	GLU	CG-CD-OE2	-9.32	96.97	118.40
1	A	22[A]	ARG	CD-NE-CZ	8.91	136.87	124.40
1	A	22[B]	ARG	CD-NE-CZ	8.91	136.87	124.40
1	A	570	GLU	OE1-CD-OE2	-8.79	101.81	122.90
1	A	579	ARG	CG-CD-NE	8.18	130.00	112.00
1	A	466	GLU	OE1-CD-OE2	7.47	140.83	122.90
1	A	579	ARG	CD-NE-CZ	7.15	134.41	124.40
1	A	22[A]	ARG	CB-CG-CD	-6.17	97.11	111.30
1	A	22[B]	ARG	CB-CG-CD	-6.17	97.11	111.30
1	A	459	GLU	CB-CG-CD	5.80	122.47	112.60
1	A	570	GLU	CG-CD-OE2	-5.34	106.12	118.40

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	22[A]	ARG	Sidechain
1	A	466	GLU	Sidechain
1	A	503[A]	GLU	Sidechain
1	A	512[A]	ARG	Sidechain
1	A	570	GLU	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5047	0	4768	69	1
2	B	23	0	21	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	23	0	21	0	0
2	D	23	0	21	0	0
2	E	23	0	21	0	1
2	F	23	0	21	0	0
2	G	23	0	21	0	0
2	H	23	0	21	1	0
3	A	24	0	36	9	0
4	A	4	0	6	2	0
5	A	24	0	24	0	0
6	A	1	0	0	0	0
7	A	958	0	0	19	3
All	All	6219	0	4981	74	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22[A]:ARG:CB	1:A:22[A]:ARG:CG	1.74	1.57
1:A:297[A]:HIS:HE1	7:A:837:HOH:O	1.20	1.18
1:A:503[A]:GLU:OE2	1:A:503[A]:GLU:HG3	1.47	1.14
1:A:503[A]:GLU:CD	1:A:503[A]:GLU:CG	2.23	1.12
1:A:503[B]:GLU:OE1	1:A:503[B]:GLU:HG2	1.52	1.06
1:A:22[A]:ARG:CB	1:A:22[A]:ARG:CD	2.34	1.06
1:A:297[A]:HIS:CE1	7:A:837:HOH:O	1.99	1.03
1:A:395:CYS:SG	4:A:703:BME:S2	2.36	1.00
1:A:22[B]:ARG:CZ	1:A:22[B]:ARG:HD3	1.94	0.97
1:A:22[B]:ARG:CZ	1:A:22[B]:ARG:CD	2.48	0.92
1:A:155:GLU:HG3	7:A:814:HOH:O	1.73	0.87
1:A:503[A]:GLU:HG3	1:A:503[A]:GLU:OE1	1.74	0.87
1:A:503[B]:GLU:CD	1:A:503[B]:GLU:CG	2.50	0.84
1:A:241[A]:TRP:NE1	7:A:805:HOH:O	2.09	0.84
1:A:401:GLN:HE22	2:B:1:GLC:H62	1.44	0.83
1:A:22[B]:ARG:CD	1:A:22[B]:ARG:CB	2.57	0.82
1:A:416[A]:GLY:O	3:A:701[A]:TRS:O2	1.97	0.81
1:A:456:HIS:H	1:A:487:GLN:HE22	1.28	0.78
1:A:287:HIS:HD2	1:A:289:LEU:H	1.30	0.77
1:A:22[A]:ARG:CG	1:A:22[A]:ARG:CA	2.63	0.76
1:A:503[B]:GLU:CD	1:A:503[B]:GLU:HB2	2.12	0.75
1:A:503[B]:GLU:CD	1:A:503[B]:GLU:CB	2.61	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:503[B]:GLU:CG	1:A:503[B]:GLU:OE2	2.38	0.72
1:A:503[A]:GLU:CG	1:A:503[A]:GLU:OE1	2.38	0.71
1:A:338:LEU:O	3:A:701[A]:TRS:H21	1.91	0.70
1:A:338:LEU:O	3:A:701[B]:TRS:H21	1.91	0.69
1:A:243[B]:MSE:SE	7:A:993:HOH:O	2.62	0.68
1:A:388:GLU:H	1:A:391:ASN:HD22	1.43	0.66
7:A:908:HOH:O	2:B:2:GLC:H62	1.95	0.66
3:A:702:TRS:H22	7:A:976:HOH:O	1.96	0.66
1:A:503[B]:GLU:OE1	1:A:503[B]:GLU:CG	2.37	0.65
7:A:908:HOH:O	2:B:2:GLC:C6	2.44	0.65
1:A:239[B]:SER:OG	7:A:807:HOH:O	2.15	0.65
1:A:216:ASN:OD1	1:A:217:HIS:HD2	1.79	0.65
1:A:503[B]:GLU:HG2	1:A:503[B]:GLU:OE2	1.98	0.64
1:A:503[A]:GLU:OE2	1:A:503[A]:GLU:CG	2.38	0.62
1:A:515:ARG:NE	7:A:814:HOH:O	2.32	0.61
1:A:22[A]:ARG:CD	1:A:22[A]:ARG:HB2	2.27	0.61
1:A:287:HIS:CD2	1:A:289:LEU:H	2.16	0.61
1:A:414:SER:O	3:A:702:TRS:H21	2.00	0.61
1:A:503[B]:GLU:OE1	1:A:503[B]:GLU:OE2	2.20	0.60
1:A:503[A]:GLU:OE2	1:A:503[A]:GLU:OE1	2.20	0.59
1:A:241[A]:TRP:CD1	7:A:805:HOH:O	2.54	0.59
1:A:581[A]:LEU:C	1:A:581[A]:LEU:HD23	2.30	0.57
1:A:401:GLN:HE22	2:B:1:GLC:C6	2.15	0.56
1:A:161[B]:ILE:CG2	1:A:209:VAL:HG13	2.37	0.55
1:A:388:GLU:H	1:A:391:ASN:ND2	2.06	0.54
1:A:204[A]:ARG:CD	7:A:1338:HOH:O	2.55	0.53
1:A:22[B]:ARG:CZ	1:A:22[B]:ARG:HD2	2.38	0.52
1:A:161[B]:ILE:HG23	1:A:209:VAL:HG13	1.93	0.51
1:A:338:LEU:O	3:A:701[B]:TRS:C2	2.58	0.50
1:A:236[B]:ARG:HH21	1:A:237[B]:PHE:HE2	1.58	0.50
1:A:236[B]:ARG:NH2	1:A:237[B]:PHE:CE2	2.81	0.49
1:A:22[A]:ARG:CG	1:A:22[A]:ARG:C	2.85	0.48
1:A:412:HIS:O	3:A:702:TRS:O2	2.30	0.48
1:A:236[B]:ARG:NH2	1:A:237[B]:PHE:HE2	2.10	0.48
1:A:397:GLN:NE2	1:A:411:LEU:H	2.11	0.48
1:A:78:ARG:CZ	1:A:97:PRO:HG2	2.44	0.47
1:A:338:LEU:O	3:A:701[A]:TRS:C2	2.59	0.46
1:A:220:PRO:HB3	1:A:241[A]:TRP:CE3	2.50	0.46
1:A:412:HIS:H	1:A:412:HIS:CD2	2.33	0.46
1:A:243[B]:MSE:SE	7:A:1435:HOH:O	2.83	0.45
1:A:293:ALA:O	1:A:297[A]:HIS:HD2	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297[A]:HIS:HE1	7:A:1463:HOH:O	2.01	0.43
1:A:539:HIS:HD2	7:A:1430:HOH:O	2.02	0.43
1:A:293:ALA:O	1:A:297[A]:HIS:CD2	2.72	0.42
7:A:1738:HOH:O	2:H:2:GLC:H61	2.18	0.42
3:A:702:TRS:N	7:A:812:HOH:O	2.29	0.42
1:A:22[A]:ARG:C	1:A:22[A]:ARG:HG3	2.44	0.42
1:A:216:ASN:ND2	1:A:276:ALA:HB3	2.35	0.42
1:A:22[B]:ARG:HD3	1:A:22[B]:ARG:CB	2.46	0.41
1:A:287:HIS:HE1	7:A:1392:HOH:O	2.03	0.41
1:A:159:THR:HB	7:A:824:HOH:O	2.21	0.41
1:A:395:CYS:CB	4:A:703:BME:S2	3.09	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1244:HOH:O	7:A:1457:HOH:O[3_755]	1.95	0.25
1:A:46:ARG:NE	2:E:1:GLC:O2[4_455]	2.07	0.13
7:A:1211:HOH:O	7:A:1356:HOH:O[3_755]	2.08	0.12
7:A:1026:HOH:O	7:A:1400:HOH:O[4_555]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	629/602 (104%)	626 (100%)	3 (0%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	516/475 (109%)	512 (99%)	4 (1%)	73	54

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22[A]	ARG
1	A	22[B]	ARG
1	A	356[A]	LEU
1	A	356[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	HIS
1	A	217	HIS
1	A	287	HIS
1	A	321	HIS
1	A	342	GLN
1	A	382	HIS
1	A	391	ASN
1	A	397	GLN
1	A	401	GLN
1	A	412	HIS
1	A	487	GLN
1	A	539	HIS
1	A	563	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	11,11,12	0.66	0	15,15,17	3.52	4 (26%)
2	GLC	B	2	2	12,12,12	0.66	0	17,17,17	1.58	3 (17%)
2	GLC	C	1	2	11,11,12	0.82	0	15,15,17	1.24	2 (13%)
2	GLC	C	2	2	12,12,12	0.88	0	17,17,17	0.86	0
2	GLC	D	1	2	11,11,12	0.81	0	15,15,17	0.58	0
2	GLC	D	2	2	12,12,12	0.74	1 (8%)	17,17,17	0.74	1 (5%)
2	GLC	E	1	2	11,11,12	0.80	0	15,15,17	0.57	0
2	GLC	E	2	2	12,12,12	0.76	0	17,17,17	0.66	0
2	GLC	F	1	2	11,11,12	0.68	0	15,15,17	1.06	1 (6%)
2	GLC	F	2	2	12,12,12	0.95	0	17,17,17	0.65	0
2	GLC	G	1	2	11,11,12	0.86	0	15,15,17	0.74	0
2	GLC	G	2	2	12,12,12	0.87	1 (8%)	17,17,17	0.49	0
2	GLC	H	1	2	11,11,12	0.63	0	15,15,17	0.71	0
2	GLC	H	2	2	12,12,12	1.06	1 (8%)	17,17,17	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/22/22	0/1/1/1
2	GLC	C	1	2	-	0/2/19/22	0/1/1/1
2	GLC	C	2	2	-	0/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/19/22	0/1/1/1
2	GLC	D	2	2	-	0/2/22/22	0/1/1/1
2	GLC	E	1	2	-	0/2/19/22	0/1/1/1
2	GLC	E	2	2	-	1/2/22/22	0/1/1/1
2	GLC	F	1	2	-	0/2/19/22	0/1/1/1
2	GLC	F	2	2	-	0/2/22/22	0/1/1/1
2	GLC	G	1	2	-	0/2/19/22	0/1/1/1
2	GLC	G	2	2	-	0/2/22/22	0/1/1/1
2	GLC	H	1	2	-	2/2/19/22	0/1/1/1
2	GLC	H	2	2	-	2/2/22/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	2	GLC	O1-C1	2.20	1.46	1.39
2	G	2	GLC	O1-C1	2.17	1.46	1.39
2	D	2	GLC	O1-C1	2.06	1.46	1.39

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-O5-C5	-11.26	97.10	112.19
2	B	1	GLC	C1-C2-C3	-6.16	100.67	109.64
2	B	2	GLC	O1-C1-C2	-5.20	93.89	108.98
2	B	1	GLC	C2-C3-C4	-2.92	105.73	110.86
2	D	2	GLC	C6-C5-C4	-2.73	106.32	113.02
2	C	1	GLC	C1-C2-C3	2.55	113.35	109.64
2	F	1	GLC	C1-O5-C5	2.47	115.50	112.19
2	C	1	GLC	O5-C5-C4	-2.39	105.02	110.83
2	B	2	GLC	O5-C5-C6	-2.23	100.91	106.44
2	B	2	GLC	O5-C1-C2	2.04	113.89	110.30
2	B	1	GLC	C6-C5-C4	-2.01	108.08	113.02

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	1	GLC	O5-C5-C6-O6
2	H	1	GLC	C4-C5-C6-O6
2	H	2	GLC	O5-C5-C6-O6
2	H	2	GLC	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	E	2	GLC	O5-C5-C6-O6



There are no ring outliers.



4 monomers are involved in 6 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	2	GLC	1	0
2	B	1	GLC	2	0
2	E	1	GLC	0	1
2	B	2	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	A	702	-	7,7,7	0.58	0	9,9,9	0.78	0
5	GLC	A	704	-	12,12,12	0.57	0	17,17,17	0.93	0
5	GLC	A	705	-	12,12,12	0.56	0	17,17,17	2.35	2 (11%)
3	TRS	A	701[A]	-	7,7,7	0.39	0	9,9,9	1.13	1 (11%)
4	BME	A	703	-	3,3,3	0.12	0	2,2,2	0.31	0
3	TRS	A	701[B]	-	7,7,7	0.92	1 (14%)	9,9,9	0.78	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	702	-	-	3/9/9/9	-
5	GLC	A	704	-	-	0/2/22/22	0/1/1/1
5	GLC	A	705	-	1/1/5/5	2/2/22/22	0/1/1/1
3	TRS	A	701[A]	-	-	6/9/9/9	-
4	BME	A	703	-	-	0/1/1/1	-
3	TRS	A	701[B]	-	-	6/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701[B]	TRS	O1-C1	2.28	1.49	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	705	GLC	O5-C1-C2	8.56	125.35	110.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701[A]	TRS	O1-C1-C	-3.26	101.78	110.88
5	A	705	GLC	O1-C1-O5	2.66	118.33	110.41
3	A	701[B]	TRS	O1-C1-C	2.16	116.90	110.88

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	705	GLC	C1

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701[A]	TRS	C2-C-C1-O1
3	A	701[A]	TRS	C3-C-C1-O1
3	A	701[A]	TRS	N-C-C1-O1
3	A	701[A]	TRS	N-C-C3-O3
3	A	701[B]	TRS	C2-C-C1-O1
3	A	701[B]	TRS	N-C-C1-O1
3	A	701[B]	TRS	N-C-C3-O3
3	A	702	TRS	C1-C-C3-O3
3	A	702	TRS	C2-C-C3-O3
3	A	702	TRS	N-C-C3-O3
5	A	705	GLC	O5-C5-C6-O6
5	A	705	GLC	C4-C5-C6-O6
3	A	701[A]	TRS	C2-C-C3-O3
3	A	701[B]	TRS	C2-C-C3-O3
3	A	701[B]	TRS	C3-C-C1-O1
3	A	701[A]	TRS	C1-C-C3-O3
3	A	701[B]	TRS	C1-C-C3-O3

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	TRS	4	0
3	A	701[A]	TRS	3	0
4	A	703	BME	2	0
3	A	701[B]	TRS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	572/602 (95%)	-0.47	9 (1%) 70 74	3, 9, 18, 36	72 (12%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	22[A]	ARG	4.3
1	A	482	ASP	3.7
1	A	240[A]	ALA	3.5
1	A	241[A]	TRP	3.1
1	A	236[A]	ARG	3.0
1	A	14	SER	2.7
1	A	204[A]	ARG	2.1
1	A	118	ASP	2.1
1	A	472	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	1	11/12	0.54	0.21	37,40,42,42	11
2	GLC	H	1	11/12	0.69	0.15	37,40,43,43	0
2	GLC	G	2	12/12	0.87	0.10	21,28,30,31	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	B	2	12/12	0.87	0.11	21,26,29,33	12
2	GLC	G	1	11/12	0.89	0.09	26,28,29,30	0
2	GLC	H	2	12/12	0.89	0.10	15,26,32,33	1
2	GLC	F	2	12/12	0.90	0.09	21,25,28,32	1
2	GLC	E	1	11/12	0.91	0.08	23,25,26,27	0
2	GLC	D	2	12/12	0.91	0.09	14,26,29,29	0
2	GLC	F	1	11/12	0.92	0.08	19,22,24,27	3
2	GLC	E	2	12/12	0.93	0.08	14,22,26,27	0
2	GLC	D	1	11/12	0.94	0.07	15,22,25,29	0
2	GLC	C	1	11/12	0.98	0.04	6,7,9,9	0
2	GLC	C	2	12/12	0.98	0.04	7,11,15,16	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TRS	A	701[A]	8/8	0.60	0.19	31,32,32,32	8
3	TRS	A	701[B]	8/8	0.60	0.19	32,32,32,32	8
3	TRS	A	702	8/8	0.62	0.23	31,31,31,32	4
5	GLC	A	704	12/12	0.65	0.16	60,60,61,61	3
5	GLC	A	705	12/12	0.81	0.17	55,56,56,57	2
4	BME	A	703	4/4	0.97	0.09	7,9,12,12	2
6	MG	A	706	1/1	1.00	0.06	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.