



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 23, 2026 – 10:26 AM UTC

PDB ID : 3BH4 / pdb_00003bh4
Title : High resolution crystal structure of Bacillus amyloliquefaciens alpha-amylase
Authors : Alikhajeh, J.; Khajeh, K.; Ranjbar, B.; Naderi-Manesh, H.; Lin, Y.H.; Liu, M.Y.; Chen, C.J.
Deposited on : 2007-11-28
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	6% 86% 13%
1	B	483	6% 87% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8435 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	483	Total	C	N	O	S	4	0	0
			3889	2466	664	753	6			
1	B	483	Total	C	N	O	S	4	0	0
			3889	2466	664	753	6			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		
2	B	4	Total	Ca	0	0
			4	4		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	312	Total	O	0	0
			312	312		
4	B	335	Total	O	0	0
			335	335		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-amylase



• Molecule 1: Alpha-amylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.76Å 148.68Å 76.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 30.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	98.3 (30.00-1.40) 99.1 (30.00-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.73 (at 1.40Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.206 , 0.219 0.205 , 0.218	Depositor DCC
R_{free} test set	19548 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.449	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 37.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8435	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/4003	0.85	10/5435 (0.2%)
1	B	0.34	0/4003	0.85	16/5435 (0.3%)
All	All	0.34	0/8006	0.85	26/10870 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	187	VAL	N-CA-C	-6.93	100.08	108.53
1	B	187	VAL	N-CA-C	-6.81	98.77	108.65
1	A	205	ASP	N-CA-C	-6.63	105.01	113.23
1	B	130	GLU	N-CA-C	-6.57	101.68	110.55
1	B	14	THR	N-CA-C	-6.54	101.34	109.64
1	B	9	TYR	N-CA-C	6.30	118.59	110.65
1	B	205	ASP	N-CA-C	-6.13	105.63	113.23
1	B	186	GLU	N-CA-C	-6.01	103.51	111.02
1	A	9	TYR	N-CA-C	5.75	118.67	110.68
1	B	264	TRP	N-CA-C	5.73	118.75	107.98
1	B	74	THR	N-CA-C	-5.55	101.25	109.86
1	A	130	GLU	N-CA-C	-5.55	103.06	110.55
1	A	356	GLU	N-CA-C	5.53	118.24	111.82
1	A	416	GLY	N-CA-C	-5.51	105.74	112.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	356	GLU	N-CA-C	5.44	118.13	111.82
1	B	56	GLY	N-CA-C	-5.43	101.26	112.34
1	B	334	GLN	N-CA-C	5.40	118.42	110.59
1	A	14	THR	N-CA-C	-5.39	101.72	109.48
1	A	264	TRP	N-CA-C	5.31	117.99	108.17
1	B	409	VAL	N-CA-C	5.23	115.89	107.73
1	B	230	ARG	N-CA-C	-5.19	99.95	108.41
1	A	74	THR	N-CA-C	-5.17	101.84	109.86
1	B	416	GLY	N-CA-C	-5.10	106.26	112.68
1	B	419	SER	N-CA-C	-5.09	107.72	114.04
1	A	57	PRO	N-CA-C	5.03	118.39	110.80
1	B	335	SER	N-CA-C	5.03	118.19	111.75

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	196	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3889	0	3630	44	0
1	B	3889	0	3630	46	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	312	0	0	2	0
4	B	335	0	0	2	1
All	All	8435	0	7260	90	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:ASN:HA	1:B:237:ILE:HD11	1.41	0.99
1:B:185:TRP:CG	1:B:186:GLU:H	1.90	0.90
1:B:103:ASN:CA	1:B:237:ILE:HD11	2.08	0.83
1:A:450:TRP:CE2	1:A:463:ILE:HD11	2.13	0.83
1:A:31:LEU:HD22	1:A:36:ILE:HD12	1.61	0.83
1:A:100:VAL:HB	1:A:231:ILE:HD13	1.63	0.80
1:A:448:GLU:HB2	1:A:463:ILE:HD12	1.65	0.78
1:B:103:ASN:HA	1:B:237:ILE:CD1	2.15	0.76
1:A:450:TRP:NE1	1:A:463:ILE:HD11	2.01	0.76
1:B:453:ILE:HD13	1:B:453:ILE:O	1.86	0.74
1:A:31:LEU:HD22	1:A:36:ILE:CD1	2.16	0.74
1:B:186:GLU:HG3	1:B:239:PHE:CD2	2.25	0.72
1:B:185:TRP:CD2	1:B:186:GLU:N	2.57	0.71
1:B:339:THR:HG21	1:B:378:ILE:HD11	1.71	0.71
1:B:420:ALA:HB3	1:B:423:SER:HB3	1.70	0.71
1:A:305:MET:SD	1:A:408:ASP:O	2.50	0.70
1:B:378:ILE:HD12	4:B:667:HOH:O	1.93	0.69
1:B:185:TRP:CG	1:B:186:GLU:N	2.55	0.67
1:B:121:PRO:HA	1:B:173:ILE:HD12	1.77	0.66
1:B:308:LEU:HD13	1:B:308:LEU:O	1.96	0.66
1:A:134:ILE:CD1	1:A:136:ALA:HB2	2.27	0.65
1:B:103:ASN:CB	1:B:237:ILE:HD11	2.29	0.63
1:B:140:PHE:HB2	1:B:154:TRP:HB2	1.81	0.62
1:B:387:PRO:HB3	1:B:453:ILE:HD13	1.82	0.62
1:B:339:THR:CG2	1:B:378:ILE:HD11	2.29	0.61
1:A:120:ASN:HD22	1:A:126:GLN:NE2	1.98	0.61
1:A:29:GLU:HA	1:A:92:ARG:HD3	1.84	0.60
1:A:6:LEU:CD2	1:A:361:GLN:HE21	2.15	0.59
1:B:186:GLU:HA	1:B:186:GLU:OE1	2.02	0.58
1:A:139:ASP:OD1	1:A:141:ARG:HG3	2.04	0.58
1:A:134:ILE:HD11	1:A:136:ALA:HB2	1.85	0.57
1:A:163:ASP:HA	1:A:172:ARG:O	2.04	0.57
1:B:387:PRO:O	1:B:453:ILE:HD12	2.04	0.57
1:B:6:LEU:CD2	1:B:361:GLN:HE21	2.18	0.57
1:B:178:GLY:O	1:B:181:LYS:HG2	2.06	0.56
1:A:140:PHE:HB2	1:A:154:TRP:HB2	1.89	0.55
1:A:31:LEU:CD2	1:A:36:ILE:HD12	2.35	0.54
1:B:121:PRO:HA	1:B:173:ILE:CD1	2.38	0.53
1:A:108:ALA:HB3	1:A:137:TRP:HB3	1.91	0.53
1:B:186:GLU:HB3	4:B:666:HOH:O	2.07	0.53
1:A:34:ILE:HD12	1:A:389:LEU:HD13	1.92	0.52
1:A:9:TYR:HB2	1:A:39:VAL:HG13	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:ASP:HA	1:B:172:ARG:O	2.10	0.51
1:A:100:VAL:CB	1:A:231:ILE:HD13	2.37	0.51
1:B:406:HIS:HB3	1:B:409:VAL:O	2.12	0.50
1:B:100:VAL:HG12	1:B:102:LEU:HG	1.93	0.50
1:B:479:SER:O	1:B:480:ILE:HD13	2.12	0.49
1:B:48:LEU:C	1:B:48:LEU:HD22	2.37	0.49
1:B:7:MET:HB2	1:B:36:ILE:HD12	1.95	0.48
1:B:40:TRP:HE3	1:B:361:GLN:NE2	2.11	0.48
1:A:185:TRP:O	1:A:186:GLU:CB	2.62	0.48
1:B:117:VAL:HG11	1:B:177:ARG:HH11	1.80	0.47
1:A:120:ASN:HD22	1:A:126:GLN:HE22	1.62	0.47
1:A:134:ILE:C	1:A:134:ILE:HD12	2.40	0.47
1:A:159:PHE:O	1:A:181:LYS:HE3	2.15	0.47
1:B:387:PRO:HB3	1:B:453:ILE:CD1	2.45	0.46
1:A:181:LYS:HA	4:A:566:HOH:O	2.16	0.46
1:A:173:ILE:HD12	1:A:173:ILE:N	2.31	0.45
1:B:427:ALA:CB	1:B:480:ILE:HD12	2.47	0.45
1:A:43:PRO:HB2	1:A:54:GLY:HA3	1.99	0.45
1:A:7:MET:SD	1:A:36:ILE:HD13	2.58	0.44
1:A:425:LEU:C	1:A:425:LEU:HD12	2.43	0.43
1:B:9:TYR:HB2	1:B:39:VAL:HG13	2.01	0.43
1:A:436:SER:HB3	1:A:472:HIS:ND1	2.33	0.43
1:A:445:ASN:HB3	1:A:463:ILE:HD13	1.99	0.43
1:A:40:TRP:C	1:A:40:TRP:CD1	2.97	0.43
1:A:185:TRP:CG	1:A:186:GLU:N	2.87	0.43
1:A:406:HIS:HB3	1:A:409:VAL:O	2.19	0.43
1:B:34:ILE:HD12	1:B:389:LEU:HD13	1.99	0.43
1:A:436:SER:HB3	1:A:472:HIS:CE1	2.54	0.43
1:B:34:ILE:HD12	1:B:389:LEU:CD1	2.49	0.43
1:B:108:ALA:HB3	1:B:137:TRP:HB3	2.01	0.43
1:B:185:TRP:HA	1:B:185:TRP:CE3	2.54	0.42
1:A:233:ALA:HB1	1:A:236:HIS:HD2	1.84	0.42
1:B:185:TRP:O	1:B:186:GLU:C	2.62	0.42
1:B:425:LEU:HD12	1:B:425:LEU:C	2.44	0.42
1:A:237:ILE:O	1:A:238:LYS:C	2.62	0.42
1:B:48:LEU:H	1:B:48:LEU:CD1	2.33	0.42
1:A:48:LEU:C	1:A:48:LEU:HD12	2.45	0.42
1:A:287:VAL:HB	1:A:288:PRO:HD3	2.02	0.42
1:B:40:TRP:CD1	1:B:40:TRP:C	2.97	0.42
1:A:42:PRO:O	1:A:43:PRO:C	2.64	0.41
1:A:163:ASP:HB3	1:A:196:TYR:OH	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:TYR:O	1:A:481:TYR:HA	2.20	0.41
1:B:237:ILE:O	1:B:238:LYS:C	2.63	0.41
1:A:60:LEU:HB3	1:A:216:TRP:CE2	2.56	0.41
1:B:453:ILE:HG22	1:B:480:ILE:HB	2.02	0.41
1:B:440:TYR:CZ	1:B:442:GLY:HA2	2.56	0.40
1:B:313:VAL:HG12	1:B:321:ALA:HB2	2.04	0.40
1:A:186:GLU:HB3	4:A:649:HOH:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:796:HOH:O	4:B:796:HOH:O[2_765]	1.58	0.62

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	481/483 (100%)	466 (97%)	14 (3%)	1 (0%)	43	19
1	B	481/483 (100%)	463 (96%)	17 (4%)	1 (0%)	43	19
All	All	962/966 (100%)	929 (97%)	31 (3%)	2 (0%)	43	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU
1	B	338	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/404 (100%)	403 (100%)	1 (0%)	87	71
1	B	404/404 (100%)	399 (99%)	5 (1%)	63	34
All	All	808/808 (100%)	802 (99%)	6 (1%)	76	52

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLU
1	B	11	GLU
1	B	40	TRP
1	B	48	LEU
1	B	308	LEU
1	B	453	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	26	ASN
1	A	93	ASN
1	A	126	GLN
1	A	299	GLN
1	A	341	GLN
1	A	361	GLN
1	B	8	GLN
1	B	26	ASN
1	B	93	ASN
1	B	247	GLN
1	B	251	GLN
1	B	299	GLN
1	B	341	GLN
1	B	361	GLN
1	B	384	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	483/483 (100%)	0.49	31 (6%)	25 26	8, 14, 24, 43	15 (3%)
1	B	483/483 (100%)	0.40	27 (5%)	30 32	7, 13, 22, 39	15 (3%)
All	All	966/966 (100%)	0.44	58 (6%)	27 29	7, 13, 24, 43	30 (3%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	185	TRP	18.6
1	B	186	GLU	9.6
1	A	186	GLU	8.5
1	B	421	ALA	7.4
1	B	420	ALA	6.8
1	B	418	SER	5.9
1	B	185	TRP	5.7
1	A	133	GLN	4.4
1	B	2	VAL	4.3
1	A	134	ILE	4.2
1	A	173	ILE	4.2
1	A	439	MET	4.2
1	B	179	GLU	4.1
1	B	419	SER	4.1
1	A	141	ARG	3.9
1	A	373	THR	3.8
1	A	182	ALA	3.4
1	B	453	ILE	3.3
1	A	463	ILE	3.2
1	B	22	LYS	3.2
1	A	305	MET	3.2
1	B	443	LEU	3.0
1	B	141	ARG	3.0
1	A	443	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	178	GLY	2.7
1	A	48	LEU	2.7
1	B	480	ILE	2.7
1	A	26	ASN	2.7
1	B	462	LYS	2.7
1	B	130	GLU	2.6
1	A	36	ILE	2.6
1	A	170	ILE	2.5
1	B	173	ILE	2.5
1	A	167	SER	2.5
1	A	136	ALA	2.5
1	B	385	ILE	2.5
1	B	388	ILE	2.5
1	A	121	PRO	2.5
1	A	163	ASP	2.4
1	B	182	ALA	2.4
1	A	93	ASN	2.4
1	A	265	GLN	2.4
1	A	177	ARG	2.4
1	A	92	ARG	2.3
1	B	373	THR	2.3
1	A	169	LYS	2.3
1	B	308	LEU	2.3
1	A	39	VAL	2.2
1	B	371	LYS	2.2
1	B	177	ARG	2.2
1	A	2	VAL	2.2
1	A	122	ALA	2.1
1	A	231	ILE	2.1
1	A	80	SER	2.1
1	B	48	LEU	2.0
1	B	484	LYS	2.0
1	B	167	SER	2.0
1	A	113	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	A	488	1/1	0.98	0.04	13,13,13,13	0
3	NA	A	489	1/1	0.98	0.13	12,12,12,12	0
3	NA	B	488	1/1	0.98	0.04	10,10,10,10	0
2	CA	B	1	1/1	0.99	0.03	10,10,10,10	0
2	CA	B	486	1/1	0.99	0.03	11,11,11,11	0
2	CA	B	487	1/1	0.99	0.04	12,12,12,12	0
2	CA	A	487	1/1	0.99	0.04	11,11,11,11	0
2	CA	A	486	1/1	0.99	0.02	11,11,11,11	0
2	CA	A	485	1/1	1.00	0.01	10,10,10,10	0
2	CA	B	485	1/1	1.00	0.02	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.