



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:15 AM UTC

PDB ID : 3BHU / pdb_00003bhu
Title : Structure of phosphorylated Thr160 CDK2/cyclin A in complex with the inhibitor meriolin 5
Authors : Echalier, A.; Bettayeb, K.; Ferandin, Y.; Lozach, O.; Clement, M.; Valette, A.; Liger, F.; Marquet, B.; Morris, J.C.; Endicott, J.A.; Joseph, B.; Meijer, L.
Deposited on : 2007-11-29
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	4% 79% 17% ...
1	C	300	5% 77% 11% .. 9%
2	B	262	3% 88% 11% .
2	D	262	2% 89% 8% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MHR	C	299	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9423 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	P	S	0	6	0
			2430	1576	413	431	1	9			
1	C	272	Total	C	N	O	P	S	0	3	0
			2198	1425	374	391	1	7			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P24941
A	0	SER	-	expression tag	UNP P24941
C	-1	GLY	-	expression tag	UNP P24941
C	0	SER	-	expression tag	UNP P24941

- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	262	Total	C	N	O	S		0	4	0
			2136	1384	347	395	10				
2	D	262	Total	C	N	O	S		0	2	0
			2123	1374	345	394	10				

- Molecule 3 is 4-(4-propoxy-1H-pyrrolo[2,3-b]pyridin-3-yl)pyrimidin-2-amine (CCD ID: MHR) (formula: C₁₄H₁₅N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			20	14	5	1		
3	C	1	Total	C	N	O	0	0
			20	14	5	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	176	Total	O	0	0
			176	176		
5	B	138	Total	O	0	0
			138	138		
5	C	90	Total	O	0	0
			90	90		
5	D	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cell division protein kinase 2




• Molecule 1: Cell division protein kinase 2




• Molecule 2: Cyclin-A2



• Molecule 2: Cyclin-A2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.15Å 133.96Å 147.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.30) 99.0 (20.00-2.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.194 , 0.240 0.190 , 0.233	Depositor DCC
R_{free} test set	3322 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.263	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 30.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9423	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, MHR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	5/2494 (0.2%)	0.92	5/3380 (0.1%)
1	C	0.68	2/2245 (0.1%)	0.93	8/3037 (0.3%)
2	B	0.59	1/2199 (0.0%)	0.89	0/2989
2	D	0.55	0/2179	0.98	8/2963 (0.3%)
All	All	0.65	8/9117 (0.1%)	0.93	21/12369 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	63	ILE	C-O	-6.33	1.17	1.24
1	A	55	LEU	C-O	-6.23	1.16	1.24
1	A	65	LYS	C-O	-5.73	1.17	1.24
1	C	63	ILE	C-O	-5.66	1.18	1.24
1	A	56	LYS	C-O	-5.52	1.17	1.24
1	A	164	VAL	C-O	-5.52	1.17	1.24
1	C	64	VAL	C-O	-5.42	1.18	1.24
2	B	172	VAL	C-O	-5.00	1.18	1.24

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	174	GLU	N-CA-C	13.99	130.86	111.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	175	VAL	N-CA-C	-11.13	79.82	111.00
1	C	163	VAL	N-CA-C	7.63	125.21	109.34
2	D	174	GLU	CA-C-N	7.19	134.65	121.70
2	D	174	GLU	C-N-CA	7.19	134.65	121.70
1	A	253	PRO	N-CA-C	6.80	119.00	110.70
2	D	431	ASN	N-CA-C	-6.37	103.46	110.91
1	C	253	PRO	N-CA-C	6.31	118.40	110.70
1	C	65	LYS	N-CA-C	5.88	118.31	108.96
1	A	65	LYS	N-CA-C	5.84	118.25	108.96
2	D	347	TYR	N-CA-C	5.84	119.67	112.54
1	C	15	TYR	N-CA-C	-5.53	99.02	110.80
1	A	163	VAL	N-CA-C	-5.51	95.58	111.00
2	D	175	VAL	CA-C-N	5.39	124.84	119.24
2	D	175	VAL	C-N-CA	5.39	124.84	119.24
1	C	1	MET	N-CA-C	-5.35	105.97	112.88
1	C	62	ASN	N-CA-C	5.29	118.75	112.72
1	A	162	GLU	CA-C-N	5.15	130.96	121.70
1	A	162	GLU	C-N-CA	5.15	130.96	121.70
1	C	-1	GLY	CA-C-N	-5.12	115.50	122.93
1	C	-1	GLY	C-N-CA	-5.12	115.50	122.93

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	15	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2430	0	2480	52	0
1	C	2198	0	2258	32	0
2	B	2136	0	2160	18	0
2	D	2123	0	2142	18	0
3	A	20	0	15	2	0
3	C	20	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	176	0	0	7	0
5	B	138	0	0	2	0
5	C	90	0	0	2	0
5	D	90	0	0	3	0
All	All	9423	0	9070	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177[A]:CYS:SG	1:A:233:MET:HE3	1.61	1.38
1:A:15:TYR:CB	1:A:16:GLY:HA3	1.70	1.20
1:A:15:TYR:HB2	1:A:16:GLY:CA	1.72	1.18
1:A:162:GLU:HB2	1:A:163:VAL:HA	1.27	1.12
1:C:15:TYR:HE1	1:C:35:ILE:CD1	1.81	0.93
1:A:161:HIS:O	1:A:162:GLU:C	2.17	0.85
1:A:177[A]:CYS:SG	1:A:233:MET:CE	2.58	0.84
1:A:162:GLU:HB2	1:A:163:VAL:CA	2.07	0.84
1:C:71:HIS:HD2	2:D:296:HIS:HE1	1.24	0.82
1:C:15:TYR:CE1	1:C:35:ILE:CD1	2.64	0.80
1:A:177[A]:CYS:HG	1:A:233:MET:HE3	1.48	0.77
1:C:15:TYR:HE1	1:C:35:ILE:HD13	1.48	0.77
2:B:183:HIS:HB2	2:B:317:GLN:HE22	1.51	0.75
1:C:15:TYR:CE1	1:C:35:ILE:HD11	2.22	0.74
1:A:0:SER:O	1:A:1:MET:HB2	1.88	0.73
1:C:278:LYS:HE3	2:D:177:ASP:O	1.89	0.72
1:C:15:TYR:HE1	1:C:35:ILE:HD11	1.55	0.70
1:A:161:HIS:O	1:A:162:GLU:O	2.10	0.70
2:D:171:SER:O	2:D:176:PRO:HD3	1.92	0.69
2:D:344:ALA:HB1	2:D:348:LEU:HD22	1.77	0.67
1:C:162[B]:GLU:HG3	1:C:163:VAL:N	2.10	0.66
1:C:71:HIS:HD2	2:D:296:HIS:CE1	2.12	0.66
1:A:14:THR:HA	1:A:15:TYR:C	2.21	0.65
1:A:15:TYR:HB3	1:A:33:LYS:HE2	1.80	0.64
2:D:175:VAL:O	2:D:175:VAL:HG13	1.98	0.64
2:D:216:ASP:HB2	2:D:406:GLN:HG3	1.80	0.63
1:C:60:HIS:CD2	1:C:62:ASN:H	2.17	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:LYS:HA	1:A:91:MET:HE2	1.81	0.62
1:C:71:HIS:CD2	2:D:296:HIS:HE1	2.12	0.61
1:A:60:HIS:CD2	1:A:62:ASN:H	2.20	0.60
1:C:60:HIS:HD2	1:C:62:ASN:H	1.49	0.60
2:B:344:ALA:HB1	2:B:348:LEU:HD22	1.85	0.59
1:C:268:HIS:HD2	1:C:270:ASP:H	1.50	0.59
2:D:274[A]:GLU:HG2	2:D:277:GLU:HG3	1.83	0.58
1:A:175:LEU:HB2	1:A:233:MET:HE2	1.84	0.58
1:C:41:THR:HG22	1:C:42[A]:GLU:HG2	1.86	0.58
1:A:60:HIS:HD2	1:A:62:ASN:H	1.52	0.57
1:C:160:TPO:HG23	1:C:162[B]:GLU:HG2	1.87	0.57
1:A:133:LEU:HD11	1:A:192:ILE:HD13	1.85	0.57
1:A:160:TPO:HG23	1:A:162:GLU:HG2	1.86	0.56
2:D:216:ASP:CB	2:D:406:GLN:HG3	2.36	0.55
1:A:95:ALA:O	1:A:96:LEU:HB2	2.06	0.55
2:B:179[A]:HIS:HE1	2:B:320:LEU:HD12	1.73	0.54
2:D:274[A]:GLU:HG2	2:D:277:GLU:CG	2.39	0.53
1:A:161:HIS:HD2	5:A:374:HOH:O	1.91	0.52
2:B:210:MET:HE1	2:B:250:ARG:CB	2.40	0.52
1:A:12:GLU:HG2	1:A:13:GLY:H	1.75	0.52
1:A:162:GLU:CB	1:A:163:VAL:HA	2.20	0.52
1:C:65:LYS:HD3	1:C:67:LEU:HD23	1.93	0.51
1:A:16:GLY:HA2	1:A:34:LYS:O	2.11	0.51
2:B:289:LYS:NZ	2:B:293:ARG:HE	2.08	0.50
2:B:395:HIS:HE1	2:B:427:PRO:O	1.95	0.49
1:C:161:HIS:CE1	5:C:336:HOH:O	2.64	0.49
2:D:175:VAL:O	2:D:175:VAL:CG1	2.61	0.48
1:A:15:TYR:HB2	1:A:16:GLY:HA3	0.77	0.48
1:A:60:HIS:HE1	5:A:466:HOH:O	1.97	0.48
1:C:119:HIS:HD2	5:D:443:HOH:O	1.95	0.48
1:A:176:GLY:O	1:A:234:PRO:HG2	2.14	0.48
2:B:225:TYR:HE2	2:B:281:ILE:HG21	1.78	0.48
2:B:361:HIS:CD2	2:B:391:LEU:HD21	2.50	0.47
1:A:2[A]:GLU:HG2	1:C:73:GLU:OE2	2.14	0.47
2:B:210:MET:HE1	2:B:250:ARG:HB2	1.95	0.47
1:A:34:LYS:NZ	1:A:75:LYS:HE2	2.29	0.47
2:D:227:LEU:HD12	2:D:261:MET:HE3	1.97	0.47
1:A:103:LEU:HD21	1:A:294:PRO:HA	1.97	0.47
1:A:260:ARG:HD3	5:A:405:HOH:O	2.13	0.47
1:C:71:HIS:CD2	2:D:296:HIS:CE1	2.97	0.47
1:A:175:LEU:CB	1:A:233:MET:HE2	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:218:THR:HG23	1:C:251:VAL:HG21	1.97	0.46
1:C:155:PRO:HB3	2:D:174:GLU:HB3	1.97	0.46
2:B:388:LYS:HB3	2:B:389:PRO:HD3	1.98	0.45
2:B:183:HIS:HD2	5:B:490:HOH:O	1.99	0.45
1:C:15:TYR:CD1	1:C:35:ILE:HG12	2.52	0.45
1:A:98:GLY:HA2	1:A:199:ARG:NE	2.32	0.45
1:C:38:ASP:HB3	1:C:41:THR:HB	1.98	0.45
1:C:41:THR:HG22	1:C:42[B]:GLU:HG3	1.98	0.45
2:B:296:HIS:NE2	2:B:300[A]:LYS:HE2	2.32	0.45
1:A:22[B]:ARG:NH1	5:A:404:HOH:O	2.49	0.44
1:A:83:LEU:O	3:A:299:MHR:H3	2.16	0.44
2:B:210:MET:HE2	2:B:210:MET:HB3	1.80	0.44
1:A:230:VAL:HA	1:A:233:MET:SD	2.58	0.43
1:A:0:SER:O	1:A:1:MET:CB	2.63	0.43
2:B:196:LYS:CG	2:B:199:TYR:HB3	2.48	0.43
1:A:-1:GLY:HA3	1:A:0:SER:HB2	2.00	0.43
1:A:119:HIS:HD2	5:B:550:HOH:O	2.01	0.43
2:B:196:LYS:HG3	2:B:199:TYR:HB3	1.99	0.43
2:D:372:TRP:HB3	2:D:384:LEU:HD11	2.00	0.43
1:A:98:GLY:HA2	1:A:199:ARG:HE	1.83	0.43
1:C:160:TPO:CG2	1:C:162[B]:GLU:HG2	2.49	0.43
1:C:148:LEU:HG	5:C:309:HOH:O	2.17	0.42
1:A:84:HIS:HB2	1:A:298:LEU:HD23	2.00	0.42
2:D:254:GLN:NE2	5:D:433:HOH:O	2.51	0.42
3:A:299:MHR:H14	3:A:299:MHR:O1	2.19	0.42
1:A:217:ARG:HG2	1:A:243:TRP:CD2	2.54	0.42
1:C:15:TYR:CE1	1:C:35:ILE:HD13	2.37	0.42
1:A:160:TPO:CG2	1:A:162:GLU:HG2	2.49	0.42
1:A:161:HIS:HE1	5:A:384:HOH:O	2.01	0.42
2:B:417:LYS:HE3	2:B:418:TYR:OH	2.20	0.42
1:A:15:TYR:CD1	1:A:15:TYR:N	2.88	0.41
1:C:130:PRO:HD2	1:C:131:GLN:OE1	2.20	0.41
1:A:88:LYS:HA	1:A:91:MET:CE	2.50	0.41
1:A:162:GLU:HG3	1:A:163:VAL:CG1	2.51	0.41
1:C:133:LEU:HD11	1:C:192:ILE:HD13	2.01	0.41
1:A:65:LYS:HD2	5:A:469:HOH:O	2.20	0.41
1:A:237:LYS:HE3	5:A:426:HOH:O	2.21	0.41
1:C:295:HIS:CD2	1:C:295:HIS:H	2.37	0.41
1:A:12:GLU:HG2	1:A:13:GLY:N	2.36	0.41
1:A:230:VAL:HG13	1:A:231:THR:N	2.36	0.41
1:C:51:GLU:O	1:C:55:LEU:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:200:MET:HB2	5:D:448:HOH:O	2.21	0.41
2:B:199:TYR:CE2	2:B:348:LEU:HD21	2.56	0.40
1:A:71:HIS:CD2	1:A:76:LEU:HD13	2.57	0.40
1:A:119:HIS:HE1	1:A:185:ASP:OD2	2.04	0.40
2:B:275:VAL:HG11	2:B:292:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/300 (100%)	289 (97%)	5 (2%)	5 (2%)	7	7
1	C	270/300 (90%)	256 (95%)	11 (4%)	3 (1%)	11	13
2	B	264/262 (101%)	262 (99%)	2 (1%)	0	100	100
2	D	262/262 (100%)	257 (98%)	5 (2%)	0	100	100
All	All	1095/1124 (97%)	1064 (97%)	23 (2%)	8 (1%)	18	23

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	GLU
1	A	1	MET
1	A	164	VAL
1	C	164	VAL
1	A	15	TYR
1	A	96	LEU
1	C	15	TYR
1	C	163	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/263 (102%)	253 (95%)	14 (5%)	21	31
1	C	240/263 (91%)	230 (96%)	10 (4%)	26	40
2	B	239/235 (102%)	232 (97%)	7 (3%)	37	55
2	D	237/235 (101%)	224 (94%)	13 (6%)	19	28
All	All	983/996 (99%)	939 (96%)	44 (4%)	25	37

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	TYR
1	A	64	VAL
1	A	75	LYS
1	A	78	LEU
1	A	97	THR
1	A	101	LEU
1	A	143	LEU
1	A	148	LEU
1	A	162	GLU
1	A	163	VAL
1	A	206[A]	ASP
1	A	206[B]	ASP
1	A	297	ARG
1	A	298	LEU
2	B	171	SER
2	B	196	LYS
2	B	232	LEU
2	B	292	LEU
2	B	348	LEU
2	B	384	LEU
2	B	432	VAL
1	C	0	SER
1	C	34	LYS
1	C	36	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	55	LEU
1	C	73	GLU
1	C	78	LEU
1	C	101	LEU
1	C	131	GLN
1	C	148	LEU
1	C	150	ARG
2	D	171	SER
2	D	174	GLU
2	D	196	LYS
2	D	232	LEU
2	D	281	ILE
2	D	283	ASP
2	D	292	LEU
2	D	296	HIS
2	D	312	ASN
2	D	348	LEU
2	D	403	GLN
2	D	406	GLN
2	D	432	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	62	ASN
1	A	71	HIS
1	A	119	HIS
1	A	161	HIS
1	A	246	GLN
1	A	268	HIS
2	B	183	HIS
2	B	228	GLN
2	B	254	GLN
2	B	313	GLN
2	B	317	GLN
2	B	395	HIS
2	B	403	GLN
2	B	431	ASN
1	C	59	ASN
1	C	60	HIS
1	C	62	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	71	HIS
1	C	113	GLN
1	C	119	HIS
1	C	265	GLN
1	C	268	HIS
1	C	287	GLN
1	C	295	HIS
2	D	173	ASN
2	D	183	HIS
2	D	228	GLN
2	D	254	GLN
2	D	296	HIS
2	D	313	GLN
2	D	396	GLN
2	D	419	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	C	160	1	8,10,11	2.32	3 (37%)	10,14,16	1.35	2 (20%)
1	TPO	A	160	1	8,10,11	0.80	0	10,14,16	1.08	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	C	160	1	-	1/9/11/13	-
1	TPO	A	160	1	-	0/9/11/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	160	TPO	P-O3P	-3.86	1.40	1.54
1	C	160	TPO	P-O2P	-3.50	1.41	1.54
1	C	160	TPO	P-O1P	-3.20	1.40	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	160	TPO	O3P-P-O2P	2.96	118.91	107.80
1	C	160	TPO	O3P-P-OG1	-2.19	97.33	105.85
1	A	160	TPO	O3P-P-O2P	2.05	115.51	107.80

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	160	TPO	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	160	TPO	2	0
1	A	160	TPO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MHR	C	299	-	22,22,22	3.45	3 (13%)	28,30,30	3.21	17 (60%)
3	MHR	A	299	-	22,22,22	3.42	3 (13%)	28,30,30	3.02	16 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MHR	C	299	-	-	6/8/8/8	0/3/3/3
3	MHR	A	299	-	-	4/8/8/8	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	299	MHR	C8-C9	14.37	1.51	1.37
3	A	299	MHR	C8-C9	14.32	1.51	1.37
3	C	299	MHR	C10-C1	5.15	1.49	1.40
3	A	299	MHR	C10-C1	4.87	1.49	1.40
3	A	299	MHR	C10-C4	4.80	1.47	1.41
3	C	299	MHR	C10-C4	4.76	1.47	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	299	MHR	C11-N5-C12	6.91	121.47	116.37
3	C	299	MHR	C11-C9-C8	-6.74	115.49	127.41
3	C	299	MHR	C11-N5-C12	5.79	120.64	116.37
3	C	299	MHR	C4-N2-C8	5.43	111.04	108.48
3	A	299	MHR	N7-C12-N5	-5.22	119.89	125.70
3	C	299	MHR	C13-N7-C12	5.21	121.39	116.21
3	C	299	MHR	N7-C12-N5	-4.94	120.20	125.70
3	A	299	MHR	C4-N2-C8	4.71	110.70	108.48
3	A	299	MHR	C13-N7-C12	4.59	120.78	116.21
3	A	299	MHR	C11-C9-C8	-4.58	119.31	127.41
3	A	299	MHR	C10-C4-N1	-4.49	121.86	126.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	299	MHR	C10-C9-C11	4.11	132.38	126.40
3	C	299	MHR	C10-C4-N1	-3.99	122.38	126.64
3	A	299	MHR	C3-C2-C1	3.59	122.05	118.07
3	C	299	MHR	C14-C13-N7	-3.58	119.58	123.97
3	C	299	MHR	C3-C2-C1	3.41	121.85	118.07
3	C	299	MHR	C10-C9-C8	-3.38	103.49	105.68
3	A	299	MHR	N6-C12-N7	3.19	120.70	117.40
3	A	299	MHR	C14-C11-N5	-3.17	119.22	122.92
3	A	299	MHR	C14-C13-N7	-3.16	120.11	123.97
3	C	299	MHR	C9-C11-N5	3.14	121.53	116.40
3	C	299	MHR	C14-C11-N5	-3.02	119.39	122.92
3	A	299	MHR	C10-C9-C8	-2.97	103.76	105.68
3	A	299	MHR	C10-C9-C11	2.85	130.54	126.40
3	C	299	MHR	C13-C14-C11	2.78	118.73	117.02
3	A	299	MHR	C9-C8-N2	-2.72	108.71	110.46
3	C	299	MHR	N6-C12-N7	2.58	120.06	117.40
3	C	299	MHR	C9-C8-N2	-2.50	108.85	110.46
3	C	299	MHR	O1-C1-C10	2.37	123.99	117.66
3	A	299	MHR	C13-C14-C11	2.20	118.37	117.02
3	A	299	MHR	O1-C1-C10	2.19	123.52	117.66
3	A	299	MHR	C2-C1-C10	-2.18	115.72	120.14
3	C	299	MHR	C2-C1-C10	-2.10	115.88	120.14

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	299	MHR	N5-C11-C9-C8
3	C	299	MHR	O1-C15-C16-C17
3	A	299	MHR	N5-C11-C9-C10
3	C	299	MHR	C2-C1-O1-C15
3	C	299	MHR	C10-C1-O1-C15
3	C	299	MHR	N5-C11-C9-C8
3	A	299	MHR	C14-C11-C9-C8
3	C	299	MHR	C14-C11-C9-C8
3	A	299	MHR	C16-C15-O1-C1
3	C	299	MHR	N5-C11-C9-C10


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	299	MHR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/300 (99%)	-0.02	12 (4%) 42 44	2, 7, 23, 32	6 (2%)
1	C	271/300 (90%)	0.11	16 (5%) 28 30	2, 9, 34, 60	3 (1%)
2	B	262/262 (100%)	-0.20	7 (2%) 56 58	2, 8, 18, 30	4 (1%)
2	D	262/262 (100%)	0.11	6 (2%) 61 63	4, 9, 22, 46	2 (0%)
All	All	1092/1124 (97%)	0.00	41 (3%) 44 46	2, 8, 25, 60	15 (1%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	251	VAL	6.1
2	D	432	VAL	5.3
2	D	171	SER	4.6
1	A	-1	GLY	4.4
2	D	428	GLU	4.4
1	C	221	THR	4.0
1	C	248	PHE	3.8
1	A	162	GLU	3.6
1	A	14	THR	3.6
1	C	295	HIS	3.6
1	C	-1	GLY	3.5
1	A	0	SER	3.4
2	B	171	SER	3.4
2	B	431	ASN	3.3
2	B	284	ASP	3.2
1	C	249	SER	3.2
2	B	432	VAL	3.1
1	A	247	ASP	3.0
1	A	96	LEU	3.0
2	D	284	ASP	2.9
1	C	253	PRO	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	431	ASN	2.8
2	B	323	GLN	2.7
2	B	428	GLU	2.6
1	C	0	SER	2.6
1	A	41	THR	2.5
1	C	216	PHE	2.4
1	C	250	LYS	2.4
1	A	15	TYR	2.4
1	C	269	TYR	2.4
2	B	283[A]	ASP	2.3
1	C	2	GLU	2.3
1	C	72	THR	2.3
2	D	179	HIS	2.3
1	A	297	ARG	2.3
1	A	73	GLU	2.2
1	A	287	GLN	2.2
1	C	71	HIS	2.2
1	C	287	GLN	2.2
1	C	252	VAL	2.1
1	A	161	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	C	160	11/12	0.97	0.06	8,11,14,14	0
1	TPO	A	160	11/12	0.98	0.06	4,8,10,10	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	D	1	1/1	0.72	0.19	15,15,15,15	0
4	MG	B	1	1/1	0.81	0.10	19,19,19,19	0
3	MHR	C	299	20/20	0.93	0.11	3,5,11,12	0
3	MHR	A	299	20/20	0.95	0.08	2,2,2,3	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MHR C 299:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MHR A 299:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.