



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:32 PM UTC

PDB ID : 3BIY / pdb_00003biy
Title : Crystal structure of p300 histone acetyltransferase domain in complex with a bisubstrate inhibitor, Lys-CoA
Authors : Liu, X.; Wang, L.; Zhao, K.; Thompson, P.R.; Hwang, Y.; Marmorstein, R.; Cole, P.A.
Deposited on : 2007-12-02
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	9% 72% 11% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase p300.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	0	0
			2589	1658	446	471	14			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1637	ARG	LYS	engineered mutation	UNP Q09472
A	1652	GLY	MET	engineered mutation	UNP Q09472

- Molecule 2 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Br	0	0
			5	5		

- Molecule 3 is [(2R,3S,4R,5R)-5-(6-amino-9H-purin-9-yl)-4-hydroxy-3-(phosphonooxy)tetrahydrofuran-2-yl]methyl (3R,20R)-20-carbamoyl-3-hydroxy-2,2-dimethyl-4,8,14,22-tetraoxo-12-thia-5,9,15,21-tetraazatricos-1-yl dihydrogen diphosphate (CCD ID: 01K) (formula: C₃₁H₅₃N₁₀O₁₉P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			64	31	10	19	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	234	Total	O	0	0
			234	234		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone acetyltransferase p300



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	61.51Å 61.51Å 101.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.65 – 1.70 26.65 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.9 (26.65-1.70) 94.8 (26.65-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 1.70Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.182 , 0.213 0.179 , 0.211	Depositor DCC
R_{free} test set	2070 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	21.9	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.048 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2892	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BR, 01K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2661	0.82	2/3600 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1479	ILE	N-CA-C	-5.11	106.69	111.45
1	A	1607	ASN	N-CA-C	5.01	117.86	111.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2589	0	2537	28	0
2	A	5	0	0	1	0
3	A	64	0	49	4	0
4	A	234	0	0	4	0
All	All	2892	0	2586	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1419:ILE:HD12	1:A:1474:ALA:HB2	1.28	1.14
1:A:1307:ASN:HD21	1:A:1321:VAL:H	1.23	0.84
1:A:1419:ILE:CD1	1:A:1474:ALA:HB2	2.11	0.76
2:A:1525:BR:BR	4:A:1850:HOH:O	2.59	0.76
1:A:1363:PHE:HE1	1:A:1654:MET:HE1	1.61	0.66
1:A:1419:ILE:CD1	1:A:1470:MET:SD	2.86	0.63
1:A:1385:CYS:O	1:A:1390:GLN:HB3	1.98	0.63
1:A:1392:ARG:HH11	1:A:1434:HIS:CE1	2.17	0.62
1:A:1660:THR:O	1:A:1663:GLN:HG2	2.01	0.59
1:A:1419:ILE:HD11	1:A:1470:MET:SD	2.43	0.59
1:A:1392:ARG:HH11	1:A:1434:HIS:HE1	1.50	0.58
1:A:1419:ILE:HD13	1:A:1470:MET:SD	2.46	0.56
3:A:700:01K:O44	4:A:1885:HOH:O	2.18	0.55
1:A:1436:TRP:CZ3	3:A:700:01K:HB	2.43	0.53
1:A:1401:VAL:HG11	1:A:1641:PHE:HB3	1.92	0.51
1:A:1378:VAL:HG12	1:A:1395:ILE:HD12	1.94	0.49
1:A:1363:PHE:CE1	1:A:1654:MET:HE1	2.45	0.47
1:A:1377:HIS:HE1	4:A:1824:HOH:O	1.97	0.47
1:A:1363:PHE:HE1	1:A:1654:MET:CE	2.27	0.46
1:A:1392:ARG:HD3	1:A:1434:HIS:CE1	2.51	0.46
1:A:1383:SER:HB3	1:A:1612:ILE:HG12	1.97	0.46
1:A:1469:LYS:NZ	4:A:1894:HOH:O	2.50	0.44
1:A:1446:TYR:O	1:A:1627:ARG:HD2	2.18	0.44
1:A:1327:HIS:HE1	1:A:1329:SER:OG	2.00	0.44
3:A:700:01K:H63A	3:A:700:01K:C	2.48	0.44
3:A:700:01K:C	3:A:700:01K:C63	2.96	0.43
1:A:1329:SER:HA	1:A:1618:LEU:HD22	2.00	0.42
1:A:1495:LEU:C	1:A:1495:LEU:HD12	2.45	0.42
1:A:1449:HIS:HD2	1:A:1639:LEU:O	2.04	0.41
1:A:1328:ALA:HA	1:A:1357:THR:O	2.21	0.41
1:A:1503:TYR:CZ	1:A:1510:PRO:HB3	2.55	0.41
1:A:1287:LYS:HD2	1:A:1289:SER:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/380 (82%)	307 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/343 (83%)	278 (98%)	6 (2%)	47	30

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1402	HIS
1	A	1427	LYS
1	A	1495	LEU
1	A	1512	VAL
1	A	1517	ILE
1	A	1637	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1307	ASN
1	A	1313	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1314	ASN
1	A	1327	HIS
1	A	1377	HIS
1	A	1379	GLN
1	A	1402	HIS
1	A	1434	HIS
1	A	1481	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	01K	A	700	-	64,66,66	1.08	4 (6%)	89,95,95	1.58	17 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	01K	A	700	-	-	8/68/84/84	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	700	01K	P43-O44	3.71	1.62	1.50
3	A	700	01K	P2-O3	3.55	1.63	1.59
3	A	700	01K	C53-N52	-2.95	1.33	1.39
3	A	700	01K	C54-N50	-2.34	1.32	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	700	01K	C19-S20-C21	-4.34	94.56	101.72
3	A	700	01K	C53-C54-N55	-4.33	120.76	126.72
3	A	700	01K	N57-C56-N55	-4.16	122.28	128.58
3	A	700	01K	CA-N-C61	3.55	131.54	121.58
3	A	700	01K	N50-C51-N52	-3.36	109.17	113.94
3	A	700	01K	O42-P43-O44	-3.34	97.44	109.33
3	A	700	01K	N55-C54-N50	2.85	132.02	127.17
3	A	700	01K	C63-C61-N	2.78	120.73	116.12
3	A	700	01K	C56-N55-C54	2.74	118.53	111.83
3	A	700	01K	C18-C19-S20	-2.67	103.44	114.39
3	A	700	01K	O46-P43-O42	2.55	115.79	105.85
3	A	700	01K	P43-O42-C41	-2.42	116.97	123.43
3	A	700	01K	O46-P43-O45	2.42	116.87	107.80
3	A	700	01K	C53-N52-C51	2.37	107.17	103.45
3	A	700	01K	C54-N50-C51	2.25	108.11	105.74
3	A	700	01K	C8-C7-C10	2.16	112.45	108.77
3	A	700	01K	C18-N17-C16	-2.01	119.08	122.82

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	700	01K	O-C-CA-CB
3	A	700	01K	N64-C-CA-CB
3	A	700	01K	C-CA-N-C61
3	A	700	01K	C63-C61-N-CA
3	A	700	01K	O62-C61-N-CA
3	A	700	01K	CA-CB-CG-CD
3	A	700	01K	C41-O42-P43-O46

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	700	01K	C18-C19-S20-C21

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	700	01K	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	317/380 (83%)	0.86	33 (10%)	11 12	17, 24, 34, 47	8 (2%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1517	ILE	4.5
1	A	1338	GLY	4.4
1	A	1519	GLU	3.3
1	A	1602	ALA	3.0
1	A	1608	SER	3.0
1	A	1607	ASN	2.9
1	A	1639	LEU	2.9
1	A	1513	LEU	2.9
1	A	1346	SER	2.8
1	A	1496	THR	2.7
1	A	1369	VAL	2.7
1	A	1511	ASN	2.7
1	A	1337	PRO	2.5
1	A	1664	ASP	2.4
1	A	1605	ALA	2.3
1	A	1344	VAL	2.3
1	A	1347	GLY	2.3
1	A	1405	ARG	2.3
1	A	1603	GLY	2.3
1	A	1516	SER	2.3
1	A	1334	GLU	2.2
1	A	1606	ALA	2.2
1	A	1660	THR	2.2
1	A	1582	GLN	2.2
1	A	1495	LEU	2.2
1	A	1613	VAL	2.1
1	A	1457	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1612	ILE	2.1
1	A	1626	GLY	2.1
1	A	1342	ARG	2.1
1	A	1662	SER	2.1
1	A	1288	PHE	2.1
1	A	1384	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BR	A	1523	1/1	0.90	0.12	43,43,43,43	1
2	BR	A	1524	1/1	0.91	0.10	46,46,46,46	1
2	BR	A	1522	1/1	0.96	0.10	37,37,37,37	1
2	BR	A	1525	1/1	0.97	0.08	45,45,45,45	1
3	01K	A	700	64/64	0.97	0.07	17,21,32,34	0
2	BR	A	1521	1/1	0.99	0.12	37,37,37,37	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 01K A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.