



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:24 AM UTC

PDB ID : 7BIL / pdb_00007bil
Title : Crystal structure of helicase Pif1 from *Thermus oshimai* in complex with oligo GGTGGTTGGTTGGTT
Authors : Dai, Y.X.; Chen, W.F.; Teng, F.Y.; Liu, N.N.; Hou, X.M.; Dou, S.X.; Rety, S.; Xi, X.G.
Deposited on : 2021-01-12
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	2% 75% 10% 14%
1	B	507	2% 76% 9% 14%
2	C	14	29% 29% 71%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PIF1 helicase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	0	0
			3504	2241	640	620	3			
1	B	436	Total	C	N	O	S	0	0	0
			3504	2241	640	620	3			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	ALA	VAL	conflict	UNP K7RJ88
A	46	SER	PRO	conflict	UNP K7RJ88
A	55	LYS	GLU	conflict	UNP K7RJ88
A	64	THR	ALA	conflict	UNP K7RJ88
A	162	ILE	MET	conflict	UNP K7RJ88
A	456	LEU	PRO	conflict	UNP K7RJ88
B	13	ALA	VAL	conflict	UNP K7RJ88
B	46	SER	PRO	conflict	UNP K7RJ88
B	55	LYS	GLU	conflict	UNP K7RJ88
B	64	THR	ALA	conflict	UNP K7RJ88
B	162	ILE	MET	conflict	UNP K7RJ88
B	456	LEU	PRO	conflict	UNP K7RJ88

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*TP*TP*TP*GP*GP*TP*TP*TP*GP*GP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	14	Total	C	N	O	P	0	0	0
			289	140	46	90	13			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
3	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0

- Molecule 5 is TETRAFLUOROALUMINATE ION (CCD ID: ALF) (formula: AlF_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	Al	F	0	0
			5	1	4		
5	B	1	Total	Al	F	0	0
			5	1	4		

- Molecule 6 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	K	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	87	Total	O	0	0
			87	87		
7	B	22	Total	O	0	0
			22	22		
7	C	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PIF1 helicase



• Molecule 1: PIF1 helicase



• Molecule 2: DNA (5'-D(*GP*GP*TP*TP*TP*GP*GP*TP*TP*TP*GP*GP*TP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	88.05Å 103.44Å 251.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.02 – 2.21 44.02 – 2.21	Depositor EDS
% Data completeness (in resolution range)	96.0 (44.02-2.21) 96.1 (44.02-2.21)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.22Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874+SVN, PHENIX 1.18.2_3874+SVN	Depositor
R, R_{free}	0.179 , 0.220 0.182 , 0.223	Depositor DCC
R_{free} test set	2768 reflections (4.81%)	wwPDB-VI
Wilson B-factor (Å ²)	50.4	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7474	wwPDB-VI
Average B, all atoms (Å ²)	70.0	wwPDB-VI

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, K, ALF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	1/3587 (0.0%)	0.95	14/4868 (0.3%)
1	B	0.40	0/3587	0.70	4/4868 (0.1%)
2	C	0.65	0/322	0.84	0/498
All	All	0.49	1/7496 (0.0%)	0.83	18/10234 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
All	All	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	332	PHE	CB-CG	5.18	1.62	1.50

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	332	PHE	O-C-N	-18.85	97.52	122.59
1	A	108	TYR	CA-C-N	15.90	153.00	125.02
1	A	108	TYR	C-N-CA	15.90	153.00	125.02
1	A	332	PHE	CA-C-O	-14.76	99.41	120.51
1	A	332	PHE	CA-C-N	13.33	145.69	121.70
1	A	332	PHE	C-N-CA	13.33	145.69	121.70
1	B	108	TYR	CA-C-N	11.91	145.99	125.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	TYR	C-N-CA	11.91	145.99	125.02
1	A	109	LYS	N-CA-C	11.61	131.36	113.72
1	A	109	LYS	CA-C-N	6.30	128.22	120.22
1	A	109	LYS	C-N-CA	6.30	128.22	120.22
1	B	403	TYR	N-CA-C	6.24	119.71	111.28
1	A	332	PHE	N-CA-C	6.18	123.97	110.80
1	A	109	LYS	N-CA-CB	-6.03	104.78	113.65
1	A	109	LYS	CB-CA-C	-5.97	99.01	110.65
1	B	404	ASP	CB-CA-C	-5.76	100.16	109.72
1	A	109	LYS	CA-C-O	-5.58	111.74	119.16
1	A	108	TYR	N-CA-C	5.53	118.37	111.24

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	TYR	Peptide
1	A	109	LYS	Peptide
1	A	332	PHE	Mainchain,Peptide
1	B	108	TYR	Peptide
1	B	109	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3504	0	3560	39	0
1	B	3504	0	3561	22	0
2	C	289	0	164	14	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	1	0	0	0	0
7	A	87	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	22	0	0	1	0
7	C	1	0	0	1	0
All	All	7474	0	7309	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:332:PHE:O	1:A:332:PHE:CD2	2.36	0.78
1:A:269:ARG:NH2	7:A:1101:HOH:O	2.17	0.77
1:B:105:GLN:O	1:B:109:LYS:HB3	1.86	0.75
1:A:397:GLU:H	2:C:6:DG:H22	1.38	0.72
1:A:237:GLN:O	1:A:500:LYS:NZ	2.28	0.64
1:A:332:PHE:O	1:A:332:PHE:CG	2.40	0.64
1:A:355:ARG:NH2	2:C:7:DG:N7	2.46	0.63
1:A:332:PHE:HD2	1:A:337:PHE:CE1	2.17	0.63
2:C:9:DT:H2'	2:C:9:DT:O2	1.99	0.63
2:C:6:DG:O5'	2:C:6:DG:H8	1.82	0.62
1:A:162:ILE:HA	1:A:165:MET:HE3	1.82	0.61
1:B:332:PHE:CD1	1:B:398:LYS:HB2	2.36	0.60
2:C:5:DT:O2	7:C:101:HOH:O	2.13	0.58
1:B:198:PRO:HG2	1:B:244:VAL:HB	1.86	0.58
1:B:108:TYR:HB3	1:B:111:ARG:HB2	1.86	0.58
1:A:332:PHE:HD2	1:A:337:PHE:HE1	1.52	0.58
1:A:303:ARG:HB3	1:A:424:ARG:NH2	2.19	0.57
1:A:448:ARG:NH2	2:C:2:DG:OP2	2.39	0.56
1:A:170:LEU:HD22	1:A:184:MET:HE1	1.87	0.56
1:A:131:HIS:HB3	2:C:6:DG:H5'	1.89	0.53
1:B:162:ILE:HG23	1:B:188:LEU:HD23	1.90	0.53
1:A:105:GLN:O	1:A:109:LYS:HB2	2.09	0.53
1:A:354:LEU:HD11	1:A:424:ARG:HG3	1.90	0.53
1:B:131:HIS:HB3	2:C:13:DT:H5''	1.90	0.53
1:A:378:LEU:HD21	1:A:421:VAL:HG11	1.91	0.52
1:A:92:PRO:HD2	1:A:95:THR:HG21	1.92	0.52
1:A:332:PHE:CE1	1:A:398:LYS:HB2	2.45	0.52
1:A:312:LYS:O	1:A:315:GLU:HG2	2.11	0.51
1:A:152:PRO:HB2	1:A:156:SER:OG	2.11	0.50
1:A:144:ARG:NH1	1:A:186:TRP:CE3	2.79	0.50
1:B:162:ILE:HA	1:B:165:MET:HE3	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:ILE:HD13	1:B:100:LEU:HD21	1.94	0.50
1:A:448:ARG:HH22	2:C:2:DG:P	2.34	0.50
1:B:307:ASP:O	1:B:311:LEU:HG	2.12	0.49
1:B:332:PHE:CE1	1:B:398:LYS:HB2	2.47	0.49
1:A:144:ARG:HD3	1:A:186:TRP:CE3	2.47	0.49
1:A:358:PRO:HB3	2:C:8:DT:H5'	1.95	0.49
1:A:394:PHE:CD1	1:A:422:PRO:HG3	2.48	0.49
1:A:332:PHE:CD1	1:A:398:LYS:HB2	2.49	0.47
1:A:145:HIS:CE1	1:A:147:GLU:HB2	2.49	0.47
1:A:357:ASP:HB2	1:A:362:TYR:CD2	2.51	0.46
1:B:84:HIS:HA	1:B:85:PRO:HD3	1.86	0.45
1:A:432:HIS:NE2	7:A:1105:HOH:O	2.36	0.45
1:B:332:PHE:HD2	1:B:337:PHE:CE1	2.35	0.45
1:A:307:ASP:O	1:A:311:LEU:HG	2.17	0.44
1:A:314:LEU:HD21	1:A:344:THR:HG22	1.99	0.44
1:B:332:PHE:HD2	1:B:337:PHE:HE1	1.66	0.44
1:B:213:LEU:HB2	1:B:435:GLN:OE1	2.18	0.43
1:B:68:LEU:HD12	1:B:68:LEU:H	1.83	0.43
1:B:241:TRP:CZ3	1:B:246:LEU:HG	2.53	0.43
1:A:135:ARG:HD3	1:B:448:ARG:HH12	1.82	0.43
1:B:410:ILE:HG22	7:B:1117:HOH:O	2.18	0.43
1:A:322:LEU:HD23	1:A:373:LEU:HB3	2.00	0.42
1:A:385:ASN:ND2	1:A:387:ARG:HD3	2.34	0.42
1:A:358:PRO:CB	2:C:8:DT:H5'	2.49	0.42
1:B:433:LYS:HE2	2:C:11:DG:OP1	2.19	0.42
1:B:300:THR:O	1:B:430:THR:HA	2.19	0.41
2:C:3:DT:H72	2:C:4:DT:H73	2.03	0.41
1:A:378:LEU:HG	1:A:391:ILE:HB	2.01	0.41
1:A:116:ALA:O	1:A:129:THR:HA	2.19	0.41
1:B:332:PHE:CE2	1:B:337:PHE:HD1	2.39	0.41
1:B:297:LEU:HD23	1:B:439:LEU:HD13	2.01	0.41
1:A:98:THR:O	1:A:101:LEU:HB3	2.21	0.41
1:A:397:GLU:H	2:C:6:DG:N2	2.11	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/507 (86%)	434 (100%)	0	0	100	100
1	B	434/507 (86%)	433 (100%)	1 (0%)	0	100	100
All	All	868/1014 (86%)	867 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/420 (87%)	357 (98%)	7 (2%)	50	64
1	B	364/420 (87%)	347 (95%)	17 (5%)	23	29
All	All	728/840 (87%)	704 (97%)	24 (3%)	33	44

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	VAL
1	A	109	LYS
1	A	387	ARG
1	A	421	VAL
1	A	446	LEU
1	A	481	LEU
1	A	485	GLU
1	B	74	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	79	VAL
1	B	81	GLN
1	B	101	LEU
1	B	167	VAL
1	B	246	LEU
1	B	262	LEU
1	B	313	ARG
1	B	328	VAL
1	B	366	ASP
1	B	372	ASP
1	B	376	GLU
1	B	400	VAL
1	B	402	THR
1	B	413	GLN
1	B	488	VAL
1	B	500	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	131	HIS
1	A	255	GLN
1	A	325	GLN
1	A	413	GLN
1	A	432	HIS
1	A	453	HIS
1	A	502	HIS
1	B	325	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ADP	A	1001	4	28,29,29	1.67	4 (14%)	43,45,45	1.89	11 (25%)
3	ADP	B	1001	4	28,29,29	1.53	4 (14%)	43,45,45	1.75	10 (23%)
5	ALF	B	1003	-	4,4,4	1.40	0	-		
5	ALF	A	1003	-	4,4,4	1.40	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1001	4	-	2/16/32/32	0/3/3/3
3	ADP	B	1001	4	-	2/16/32/32	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1001	ADP	C5-C4	5.30	1.48	1.39
3	A	1001	ADP	C5-C4	4.85	1.47	1.39
3	A	1001	ADP	PA-O3A	4.40	1.64	1.59
3	B	1001	ADP	C5-C6	2.89	1.49	1.41
3	A	1001	ADP	C5-C6	2.84	1.48	1.41
3	B	1001	ADP	C8-N7	2.66	1.36	1.31
3	A	1001	ADP	C8-N7	2.56	1.36	1.31
3	B	1001	ADP	PA-O3A	2.26	1.61	1.59

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	ADP	C5-C4-N3	-4.91	119.95	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1001	ADP	C5-C4-N3	-4.90	119.97	126.72
3	B	1001	ADP	N3-C4-N9	4.22	134.34	127.17
3	A	1001	ADP	N3-C4-N9	4.08	134.10	127.17
3	A	1001	ADP	N3-C2-N1	-3.81	122.81	128.58
3	A	1001	ADP	C2-N3-C4	3.69	120.83	111.83
3	A	1001	ADP	C4-N9-C8	3.38	109.28	105.74
3	A	1001	ADP	C4-C5-N7	-3.23	106.88	110.58
3	B	1001	ADP	C2-N3-C4	3.19	119.62	111.83
3	B	1001	ADP	N3-C2-N1	-3.16	123.81	128.58
3	B	1001	ADP	O3A-PB-O1B	-3.02	95.15	111.04
3	B	1001	ADP	C4-N9-C8	2.97	108.85	105.74
3	B	1001	ADP	C4-C5-N7	-2.81	107.37	110.58
3	A	1001	ADP	C6-C5-N7	2.78	137.45	132.09
3	A	1001	ADP	C5-N7-C8	2.78	107.82	103.45
3	A	1001	ADP	N9-C8-N7	-2.77	110.01	113.94
3	B	1001	ADP	C2-N1-C6	2.59	122.98	118.73
3	A	1001	ADP	C2-N1-C6	2.24	122.42	118.73
3	B	1001	ADP	O3B-PB-O1B	2.20	119.41	110.83
3	B	1001	ADP	C6-C5-N7	2.10	136.13	132.09
3	A	1001	ADP	O3B-PB-O2B	2.00	115.31	107.80

There are no chirality outliers.

All (4) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	A	1001	ADP	PA-O3A-PB-O2B
3	B	1001	ADP	PA-O3A-PB-O2B
3	B	1001	ADP	PA-O3A-PB-O3B
3	A	1001	ADP	PA-O3A-PB-O1B


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/507 (85%)	-0.03	12 (2%) 55 52	37, 58, 86, 158	0
1	B	436/507 (85%)	0.28	10 (2%) 61 59	52, 72, 105, 148	0
2	C	14/14 (100%)	1.24	4 (28%) 1 1	65, 88, 172, 184	0
All	All	886/1028 (86%)	0.14	26 (2%) 53 51	37, 66, 102, 184	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	1	DG	4.7
1	B	482	TRP	4.2
1	B	311	LEU	3.8
1	B	403	TYR	3.8
1	A	403	TYR	3.5
1	A	482	TRP	3.4
2	C	7	DG	3.4
1	A	311	LEU	3.3
2	C	14	DT	3.2
1	A	404	ASP	2.7
1	A	135	ARG	2.6
1	A	410	ILE	2.6
1	B	99	THR	2.6
1	B	410	ILE	2.6
1	A	109	LYS	2.5
1	B	481	LEU	2.5
1	A	332	PHE	2.3
1	B	404	ASP	2.3
1	A	500	LYS	2.2
1	A	407	ARG	2.2
2	C	8	DT	2.2
1	B	401	TYR	2.1
1	B	155	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	409	GLU	2.1
1	A	401	TYR	2.0
1	B	132	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	K	A	1004	1/1	0.88	0.14	97,97,97,97	0
3	ADP	B	1001	27/27	0.96	0.09	57,65,84,89	0
5	ALF	B	1003	5/5	0.97	0.05	52,53,57,57	0
3	ADP	A	1001	27/27	0.98	0.07	31,42,57,63	0
5	ALF	A	1003	5/5	0.98	0.06	36,36,43,43	0
4	MG	B	1002	1/1	0.99	0.02	62,62,62,62	0
4	MG	A	1002	1/1	0.99	0.03	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADP B 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ALF B 1003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ALF A 1003:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG B 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.