



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:08 PM UTC

PDB ID : 7BJT / pdb_00007bjt
Title : Structure-function analysis of a new PL17 oligoalginate lyase from the marine bacterium *Zobellia galactanivorans* DsijT
Authors : Czjzek, M.; Roret, T.; Jouanneau, D.; Le Duff, N.; Jeudy, A.
Deposited on : 2021-01-14
Resolution : 1.42 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	4% 93%
1	B	751	7% 93%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alginate lyase, family PL17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	727	Total	C	N	O	S	0	5	0
			5837	3721	984	1118	14			
1	B	727	Total	C	N	O	S	0	9	0
			5863	3736	988	1125	14			


- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		
3	B	2	Total	Mg	0	0
			2	2		

- Molecule 4 is D-MALATE (CCD ID: MLT) (formula: C₄H₆O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			9	4	5		
4	B	1	Total	C	O	0	0
			9	4	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	668	Total	O	0	0
			668	668		
6	B	658	Total	O	0	0
			658	658		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alginate lyase, family PL17



- Molecule 1: Alginate lyase, family PL17



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	163.39Å 163.39Å 166.73Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.70 – 1.42 43.70 – 1.42	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.70-1.42) 99.8 (43.70-1.42)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.42Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.161 , 0.170 0.165 , 0.174	Depositor DCC
R_{free} test set	23668 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.015 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13069	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MLT, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/5987 (0.0%)	0.68	1/8099 (0.0%)
1	B	0.45	0/6032	0.69	0/8160
All	All	0.46	1/12019 (0.0%)	0.69	1/16259 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	433	ASP	CA-CB	-6.86	1.48	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	711	ASN	N-CA-C	7.46	122.04	112.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5837	0	5664	16	0
1	B	5863	0	5697	16	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	9	0	4	0	0
4	B	9	0	4	0	0
5	A	6	0	8	0	0
5	B	12	0	16	0	0
6	A	668	0	0	0	0
6	B	658	0	0	3	0
All	All	13069	0	11393	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:585:LEU:HD12	1:B:585:LEU:HD12	1.53	0.91
1:B:453:ALA:H	1:B:470:ASN:HD22	1.27	0.83
1:A:453:ALA:H	1:A:470:ASN:HD22	1.24	0.80
1:B:240:ASN:ND2	6:B:901:HOH:O	2.18	0.72
1:B:609:ASN:HD21	1:B:692:ILE:H	1.40	0.70
1:A:609:ASN:HD21	1:A:692:ILE:H	1.41	0.67
1:B:71:ASP:OD2	1:B:82:TYR:OH	2.16	0.62
1:B:236:GLY:HA3	6:B:1427:HOH:O	2.00	0.60
1:B:275:GLN:O	1:B:279:MET:HG2	2.05	0.56
1:A:275:GLN:O	1:A:279:MET:HG2	2.06	0.55
1:B:279:MET:HA	1:B:279:MET:HE2	1.88	0.55
1:A:431:HIS:HB3	1:A:433:ASP:OD2	2.06	0.54
1:B:668:ILE:HG21	1:B:689:ILE:HD11	1.90	0.54
1:A:433:ASP:OD2	1:A:480:HIS:CE1	2.62	0.52
1:B:708:ALA:HB3	1:B:713:GLU:HB2	1.91	0.52
1:A:279:MET:HE2	1:A:279:MET:HA	1.93	0.50
1:A:486:ASN:HD21	1:A:560:ASN:HD22	1.61	0.48
1:A:668:ILE:HG21	1:A:689:ILE:HD11	1.96	0.48
1:A:283[A]:LEU:HD13	1:A:343:ALA:HA	1.95	0.47
1:A:555:GLY:HA2	1:A:662:ASN:HD22	1.78	0.47
1:A:433:ASP:OD2	1:A:480:HIS:NE2	2.49	0.46
1:B:437:PHE:CZ	1:B:447:LEU:HD23	2.53	0.43
1:A:453:ALA:H	1:A:470:ASN:ND2	2.05	0.43
1:A:283[B]:LEU:HD22	1:A:343:ALA:HA	2.01	0.43
1:A:446:ILE:O	1:A:447:LEU:HD12	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:668:ILE:CG2	1:B:689:ILE:HD11	2.50	0.42
1:B:555:GLY:HA2	1:B:662:ASN:HD22	1.85	0.42
1:B:203:TRP:CZ2	1:B:281:PRO:HD3	2.55	0.41
1:A:431:HIS:CD2	1:A:475:LYS:HG2	2.55	0.41
1:B:539[B]:LYS:NZ	6:B:904:HOH:O	2.36	0.41
1:B:368:LEU:O	1:B:373:GLY:HA3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	729/751 (97%)	706 (97%)	23 (3%)	0	100	100
1	B	735/751 (98%)	710 (97%)	24 (3%)	1 (0%)	48	22
All	All	1464/1502 (98%)	1416 (97%)	47 (3%)	1 (0%)	48	22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	711	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	622/643 (97%)	617 (99%)	5 (1%)	73	47
1	B	629/643 (98%)	625 (99%)	4 (1%)	78	56
All	All	1251/1286 (97%)	1242 (99%)	9 (1%)	76	52

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	LYS
1	A	195	ASN
1	A	452	LEU
1	A	583	LYS
1	A	629	ASN
1	B	452	LEU
1	B	604	GLN
1	B	669	LEU
1	B	735	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	HIS
1	A	144	GLN
1	A	150	ASN
1	A	205	ASN
1	A	301	HIS
1	A	465	ASN
1	A	470	ASN
1	A	476	GLN
1	A	485	GLN
1	A	500	GLN
1	A	560	ASN
1	A	609	ASN
1	A	611	GLN
1	A	662	ASN
1	A	684	ASN
1	A	686	ASN
1	B	95	GLN
1	B	144	GLN
1	B	150	ASN
1	B	205	ASN
1	B	222	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	250	GLN
1	B	275	GLN
1	B	353	HIS
1	B	389	GLN
1	B	470	ASN
1	B	476	GLN
1	B	485	GLN
1	B	560	ASN
1	B	571	GLN
1	B	609	ASN
1	B	628	ASN
1	B	662	ASN
1	B	686	ASN
1	B	727	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	802	-	5,5,5	0.33	0	5,5,5	0.32	0
5	GOL	B	804	-	5,5,5	0.65	0	5,5,5	0.54	0
4	MLT	B	803	-	8,8,8	1.73	2 (25%)	10,10,10	1.64	2 (20%)
4	MLT	A	803	-	8,8,8	1.10	0	10,10,10	1.19	1 (10%)
5	GOL	A	804	-	5,5,5	0.33	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	B	802	-	-	0/4/4/4	-
5	GOL	B	804	-	-	0/4/4/4	-
4	MLT	B	803	-	-	0/8/8/8	-
4	MLT	A	803	-	-	4/8/8/8	-
5	GOL	A	804	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	803	MLT	O5-C4	-3.53	1.19	1.30
4	B	803	MLT	O2-C1	-3.21	1.20	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	803	MLT	O2-C1-C2	3.21	119.54	112.74
4	A	803	MLT	O2-C1-C2	2.31	117.62	112.74
4	B	803	MLT	O1-C1-C2	-2.19	118.23	122.60

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	803	MLT	O1-C1-C2-C3
4	A	803	MLT	O2-C1-C2-C3
4	A	803	MLT	O1-C1-C2-O3
4	A	803	MLT	O2-C1-C2-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	727/751 (96%)	0.07	33 (4%)	38 40	9, 16, 27, 37	4 (0%)
1	B	727/751 (96%)	0.29	54 (7%)	20 22	7, 17, 28, 40	22 (3%)
All	All	1454/1502 (96%)	0.18	87 (5%)	27 30	7, 16, 28, 40	26 (1%)

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	712	THR	13.3
1	B	235	ILE	11.2
1	B	710	SER	9.4
1	B	711	ASN	9.2
1	A	711	ASN	7.9
1	B	713	GLU	7.7
1	B	242	GLY	6.5
1	A	710	SER	6.5
1	B	82	TYR	5.9
1	B	234	PRO	5.9
1	B	25	ALA	5.6
1	A	25	ALA	5.6
1	B	236	GLY	5.3
1	B	240	ASN	5.3
1	A	712	THR	4.9
1	A	583	LYS	4.5
1	A	604	GLN	4.5
1	B	604	GLN	4.4
1	B	413	ASN	4.2
1	B	709	LYS	4.2
1	A	413	ASN	4.1
1	A	433	ASP	4.1
1	A	606	LYS	4.1
1	A	629	ASN	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	239	ASP	4.0
1	B	629	ASN	4.0
1	B	230	ASP	3.9
1	A	740	ARG	3.8
1	B	233	LEU	3.7
1	B	736	GLY	3.7
1	A	557	ASN	3.6
1	A	47	ASN	3.6
1	B	229	GLU	3.6
1	B	734	ILE	3.6
1	B	47	ASN	3.5
1	B	557	ASN	3.5
1	B	243	GLY	3.3
1	B	591	ASP	3.2
1	B	237	ALA	3.2
1	B	39	GLU	3.1
1	B	574	GLN	3.1
1	B	731	ILE	3.1
1	B	606	LYS	3.1
1	B	740	ARG	2.9
1	A	731	ILE	2.9
1	A	735	LYS	2.9
1	B	585	LEU	2.9
1	A	628	ASN	2.9
1	A	530	GLU	2.9
1	B	689	ILE	2.8
1	A	689	ILE	2.7
1	A	669	LEU	2.6
1	B	241	ASP	2.6
1	A	713	GLU	2.6
1	B	352	ASN	2.6
1	A	729	GLU	2.5
1	B	583	LYS	2.5
1	A	585	LEU	2.5
1	B	669	LEU	2.5
1	A	694	LEU	2.5
1	B	737	LYS	2.4
1	B	735	LYS	2.4
1	B	732	ILE	2.4
1	B	530	GLU	2.4
1	A	737	LYS	2.4
1	A	352	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	748	ILE	2.4
1	B	729	GLU	2.4
1	B	733	GLU	2.4
1	A	591	ASP	2.3
1	B	699	LYS	2.3
1	B	198[A]	HIS	2.3
1	B	27	GLU	2.3
1	A	734	ILE	2.2
1	A	584	SER	2.2
1	B	581	THR	2.1
1	B	628	ASN	2.1
1	B	172	LYS	2.1
1	A	39	GLU	2.1
1	A	229	GLU	2.1
1	B	607	GLY	2.1
1	A	732	ILE	2.1
1	B	726	VAL	2.0
1	B	353	HIS	2.0
1	A	230	ASP	2.0
1	B	707	ASP	2.0
1	B	727	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	B	804	6/6	0.90	0.12	20,20,20,20	0
3	MG	B	801	1/1	0.91	0.15	34,34,34,34	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MLT	B	803	9/9	0.95	0.08	18,19,21,22	0
5	GOL	B	802	6/6	0.96	0.09	19,21,22,22	0
4	MLT	A	803	9/9	0.97	0.06	16,16,18,18	0
5	GOL	A	804	6/6	0.97	0.07	16,16,17,17	0
3	MG	A	802	1/1	0.98	0.10	22,22,22,22	0
3	MG	B	806	1/1	0.99	0.07	11,11,11,11	0
3	MG	A	805	1/1	0.99	0.07	12,12,12,12	0
2	CA	A	806	1/1	0.99	0.07	17,17,17,17	0
2	CA	B	805	1/1	1.00	0.06	8,8,8,8	0
2	CA	A	801	1/1	1.00	0.07	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.