



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:58 PM UTC

PDB ID : 8BJH / pdb_00008bjh
Title : chimera of the inactive ExoY Nucleotidyl Cyclase domain from *Vibrio nigrificans* MARTX toxin, with the double mutation K3528M and K3535I, fused to a proline-rich domain (PRD) and profilin, bound to Latrunculin B-ADP-Mg-actin
Authors : Teixeira-Nunes, M.; Renault, L.; Retailleau, P.
Deposited on : 2022-11-04
Resolution : 1.69 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	7% 93% 5%
2	B	591	14% 77% 7% 15%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 7980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Actin, alpha skeletal muscle, intermediate form.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	366	Total	C	N	O	S	0	13	0
			2948	1881	493	553	21			

- Molecule 2 is a protein called Putative Adenylate cyclase, Profilin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	502	Total	C	N	O	S	0	12	0
			3952	2500	655	781	16			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	451	GLY	-	expression tag	UNP A0A6N3LUE9
B	452	PRO	-	expression tag	UNP A0A6N3LUE9
B	453	GLY	-	expression tag	UNP A0A6N3LUE9
B	454	SER	-	expression tag	UNP A0A6N3LUE9
B	528	MET	LYS	engineered mutation	UNP A0A6N3LUE9
B	535	ILE	LYS	engineered mutation	UNP A0A6N3LUE9
B	897	PRO	-	linker	UNP A0A6N3LUE9
B	898	PRO	-	linker	UNP A0A6N3LUE9
B	899	PRO	-	linker	UNP A0A6N3LUE9
B	900	PRO	-	linker	UNP A0A6N3LUE9
B	901	GLY	-	linker	UNP A0A6N3LUE9
B	902	GLY	-	linker	UNP A0A6N3LUE9

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0

- Molecule 5 is LATRUNCULIN B (CCD ID: LAB) (formula: $C_{20}H_{29}NO_5S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			27	20	1	5	1		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		
6	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is HYDROGEN PEROXIDE (CCD ID: PEO) (formula: H_2O_2).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	O	0	0
			2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total O 2 2	0	0
8	A	1	Total O 2 2	0	0
8	A	1	Total O 2 2	0	0
8	A	1	Total O 2 2	0	0
8	B	1	Total O 2 2	0	0
8	B	1	Total O 2 2	0	0
8	B	1	Total O 2 2	0	0

- Molecule 9 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total O S 5 4 1	0	0
9	B	1	Total O S 5 4 1	0	0

- Molecule 10 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	415	Total	O	0	0
			415	415		
11	B	543	Total	O	0	0
			543	543		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Actin, alpha skeletal muscle, intermediate form



- Molecule 2: Putative Adenylate cyclase, Profilin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.74Å 62.83Å 93.58Å 90.00° 91.03° 90.00°	Depositor
Resolution (Å)	93.57 – 1.69 93.57 – 1.69	Depositor EDS
% Data completeness (in resolution range)	76.5 (93.57-1.69) 76.5 (93.57-1.69)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 1.69Å)	Xtriage
Refinement program	BUSTER 2.10.3 (20-MAY-2020)	Depositor
R, R_{free}	0.178 , 0.212 0.182 , 0.215	Depositor DCC
R_{free} test set	4009 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7980	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, TRS, PEO, MG, SO4, LAB, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	2/3053 (0.1%)	0.95	0/4133
2	B	0.70	1/4065 (0.0%)	0.94	4/5504 (0.1%)
All	All	0.72	3/7118 (0.0%)	0.94	4/9637 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	762	ALA	CA-C	10.47	1.58	1.52
1	A	176	MET	SD-CE	-6.52	1.63	1.79
1	A	257	CYS	CA-C	5.19	1.58	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	773	PHE	CA-CB-CG	-5.81	107.99	113.80
2	B	914	ALA	N-CA-C	5.74	117.21	111.07
2	B	622	ILE	N-CA-C	5.50	113.32	107.76
2	B	513	ASP	CA-CB-CG	5.37	117.97	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2948	0	2954	10	0
2	B	3952	0	3899	23	0
3	A	27	0	12	0	0
4	A	1	0	0	0	0
5	A	27	0	29	0	0
6	A	8	0	12	0	0
6	B	8	0	12	0	0
7	A	6	0	8	0	0
7	B	12	0	16	2	0
8	A	10	0	0	3	0
8	B	6	0	0	1	0
9	B	10	0	0	1	0
10	B	7	0	10	0	0
11	A	415	0	0	0	0
11	B	543	0	0	1	0
All	All	7980	0	6952	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:B:1207:PEO:O1	8:B:1207:PEO:O2	1.53	1.26
8:A:409:PEO:O1	8:A:409:PEO:O2	1.53	1.25
8:A:408:PEO:O1	8:A:408:PEO:O2	1.53	1.25
8:A:406:PEO:O1	8:A:406:PEO:O2	1.58	1.22
2:B:617:ASN:HB2	7:B:1205:GOL:H11	1.72	0.71
1:A:353:GLN:HA	1:A:356[B]:TRP:CD1	2.25	0.70
1:A:353:GLN:HA	1:A:356[B]:TRP:HD1	1.58	0.68
2:B:468:TYR:HE1	2:B:788:LYS:HZ3	1.44	0.66
1:A:372:ARG:HG3	2:B:1026:ASN:OD1	1.99	0.63
1:A:117:GLU:OE2	1:A:371[B]:HIS:NE2	2.25	0.61
2:B:468:TYR:HE1	2:B:788:LYS:NZ	2.05	0.54
1:A:244:ASP:OD1	1:A:246[A]:GLN:HB2	2.09	0.52
2:B:490[A]:GLU:HG2	9:B:1202:SO4:O3	2.11	0.51
2:B:602:LEU:HD21	2:B:626:LEU:HD12	1.93	0.49
2:B:617:ASN:HB2	7:B:1205:GOL:C1	2.40	0.49
1:A:39:ARG:HG2	1:A:66:THR:HG23	1.94	0.49
2:B:598:SER:HB3	11:B:1314:HOH:O	2.13	0.47
2:B:799[A]:GLU:H	2:B:799[A]:GLU:CD	2.24	0.46
2:B:577:ALA:HB1	2:B:653:ILE:HG23	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:TYR:HB2	1:A:72:GLU:HG3	1.97	0.45
2:B:838:PRO:HB2	2:B:839:LEU:HD23	1.98	0.45
2:B:527:SER:OG	2:B:752:HIS:NE2	2.45	0.45
2:B:631:VAL:HB	2:B:636:LEU:HD12	1.99	0.44
2:B:468:TYR:HB3	2:B:475:LEU:HB2	1.99	0.44
1:A:59:GLN:HE21	2:B:684:VAL:HG12	1.83	0.43
2:B:684:VAL:HG23	2:B:685:LYS:HD2	2.01	0.43
2:B:861:LEU:O	2:B:1019:GLY:HA2	2.19	0.42
2:B:1016:GLY:HA3	2:B:1025:ILE:HD11	2.02	0.42
2:B:764:MET:SD	2:B:768:PHE:HD2	2.43	0.41
2:B:837:SER:HA	2:B:838:PRO:HD3	1.99	0.41
1:A:352[B]:PHE:HD1	1:A:355:MET:SD	2.44	0.41
2:B:764:MET:SD	2:B:768:PHE:CD2	3.13	0.41
1:A:269:MET:HE2	1:A:269:MET:HB3	1.93	0.41
2:B:811:PHE:CD2	2:B:840:PHE:HZ	2.39	0.41
2:B:896:PRO:HA	2:B:897:PRO:HD3	1.97	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/375 (100%)	368 (98%)	7 (2%)	0	100	100
2	B	508/591 (86%)	484 (95%)	19 (4%)	5 (1%)	12	3
All	All	883/966 (91%)	852 (96%)	26 (3%)	5 (1%)	21	9

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	835	LEU
2	B	763	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	833	SER
2	B	759	ASN
2	B	838	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/318 (102%)	321 (99%)	3 (1%)	70	62
2	B	440/504 (87%)	430 (98%)	10 (2%)	44	27
All	All	764/822 (93%)	751 (98%)	13 (2%)	55	38

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180	LEU
1	A	246[A]	GLN
1	A	246[B]	GLN
2	B	513	ASP
2	B	526	SER
2	B	542	MET
2	B	560	ASP
2	B	657	VAL
2	B	788	LYS
2	B	828	ASN
2	B	839	LEU
2	B	960	PHE
2	B	1010	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	280	ASN
2	B	640	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	815	GLN
2	B	981	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	PEG	B	1206	-	6,6,6	0.24	0	5,5,5	0.04	0
6	TRS	B	1203	-	7,7,7	0.18	0	9,9,9	0.29	0
8	PEO	A	408	-	1,1,1	0.76	0	-		
8	PEO	A	407	-	1,1,1	0.57	0	-		
8	PEO	A	409	-	1,1,1	0.72	0	-		
6	TRS	A	404	-	7,7,7	0.25	0	9,9,9	0.33	0
8	PEO	A	406	-	1,1,1	1.08	0	-		
7	GOL	A	405	-	5,5,5	0.06	0	5,5,5	0.25	0
8	PEO	B	1208	-	1,1,1	0.60	0	-		
8	PEO	A	410	-	1,1,1	0.24	0	-		
7	GOL	B	1204	-	5,5,5	0.05	0	5,5,5	0.21	0
3	ADP	A	401	4	28,29,29	0.48	0	43,45,45	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	B	1205	-	5,5,5	0.09	0	5,5,5	0.23	0
8	PEO	B	1207	-	1,1,1	0.79	0	-		
5	LAB	A	403	-	28,29,29	0.88	1 (3%)	29,41,41	0.74	1 (3%)
9	SO4	B	1202	-	4,4,4	0.31	0	6,6,6	0.37	0
8	PEO	B	1209	-	1,1,1	0.37	0	-		
9	SO4	B	1201	-	4,4,4	0.27	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	PEG	B	1206	-	-	4/4/4/4	-
6	TRS	B	1203	-	-	0/9/9/9	-
6	TRS	A	404	-	-	0/9/9/9	-
7	GOL	A	405	-	-	0/4/4/4	-
7	GOL	B	1204	-	-	0/4/4/4	-
3	ADP	A	401	4	-	0/16/32/32	0/3/3/3
7	GOL	B	1205	-	-	0/4/4/4	-
5	LAB	A	403	-	-	4/21/49/49	0/2/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	403	LAB	C15-C16	4.43	1.62	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	403	LAB	C3-C2-C1	2.21	132.65	127.36

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	403	LAB	O3-C15-C16-C17
10	B	1206	PEG	O2-C3-C4-O4
10	B	1206	PEG	O1-C1-C2-O2
10	B	1206	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
10	B	1206	PEG	C4-C3-O2-C2
5	A	403	LAB	O2-C1-C2-C3
5	A	403	LAB	C11-C10-C9-C8
5	A	403	LAB	O1-C1-C2-C3


There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	408	PEO	1	0
8	A	409	PEO	1	0
8	A	406	PEO	1	0
7	B	1205	GOL	2	0
8	B	1207	PEO	1	0
9	B	1202	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	366/375 (97%)	0.14	25 (6%) 23 25	13, 25, 51, 90	13 (3%)
2	B	502/591 (84%)	0.78	81 (16%) 4 4	13, 31, 77, 93	12 (2%)
All	All	868/966 (89%)	0.51	106 (12%) 8 8	13, 28, 69, 93	25 (2%)

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	839	LEU	7.4
2	B	835	LEU	7.1
2	B	763	VAL	6.7
2	B	840	PHE	6.6
2	B	762	ALA	6.5
2	B	684	VAL	6.5
2	B	838	PRO	6.1
2	B	827	LEU	6.0
1	A	43	VAL	5.4
2	B	760	PRO	5.2
2	B	842	THR	5.2
2	B	755	ALA	5.1
2	B	995	GLY	5.1
2	B	765	ALA	5.0
2	B	902	GLY	4.7
2	B	834	GLY	4.6
2	B	831	TRP	4.5
2	B	761	TYR	4.5
2	B	836	ASP	4.5
2	B	841	THR	4.5
2	B	759	ASN	4.3
2	B	837	SER	4.2
1	A	38	PRO	4.0
1	A	42	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	40	HIS	3.9
2	B	787	GLY	3.8
2	B	897	PRO	3.8
2	B	758	ALA	3.8
2	B	764	MET	3.7
2	B	994	THR	3.7
1	A	67	LEU	3.6
2	B	896	PRO	3.6
2	B	833	SER	3.6
1	A	65	LEU	3.6
2	B	683	LYS	3.4
2	B	901	GLY	3.4
1	A	39	ARG	3.4
2	B	757	ASP	3.4
2	B	766	ASP	3.4
1	A	5	THR	3.4
2	B	685	LYS	3.4
2	B	996	GLY	3.2
2	B	682	ASP	3.2
1	A	66	THR	3.2
1	A	64	ILE	3.2
2	B	843	LYS	3.1
2	B	862	GLY	3.1
2	B	468	TYR	3.0
2	B	828	ASN	3.0
1	A	63	GLY	3.0
2	B	784	LEU	3.0
2	B	938	GLY	3.0
2	B	904	GLY	2.9
2	B	903	ALA	2.9
2	B	478	ILE	2.9
1	A	73	HIS	2.8
2	B	475	LEU	2.8
1	A	36	GLY	2.8
2	B	606	VAL	2.8
2	B	681	GLN	2.7
2	B	811	PHE	2.7
2	B	675	TYR	2.7
2	B	680	PRO	2.6
1	A	351	THR	2.6
2	B	633	ASP	2.6
2	B	729	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	826	SER	2.6
2	B	756	ASP	2.5
2	B	480	HIS	2.5
2	B	1041	TYR	2.5
2	B	900	PRO	2.5
2	B	895	SER	2.5
2	B	861	LEU	2.4
2	B	785	GLY	2.4
2	B	786	GLU	2.4
2	B	788	LYS	2.4
2	B	782	ASP	2.4
2	B	982	ASP	2.4
1	A	41	GLN	2.4
2	B	481	PRO	2.4
1	A	51	ASP	2.4
2	B	631	VAL	2.4
2	B	783	GLY	2.3
1	A	71	ILE	2.3
2	B	469	GLN	2.3
2	B	474	VAL	2.3
1	A	62	ARG	2.3
2	B	899	PRO	2.3
2	B	482	LYS	2.3
1	A	231	ALA	2.3
2	B	466	LYS	2.3
2	B	1008	ASP	2.3
2	B	467	THR	2.2
1	A	60	SER	2.2
2	B	676	GLY	2.2
1	A	37	ARG	2.2
2	B	928	ASP	2.2
2	B	845	LYS	2.2
2	B	829	ASP	2.2
1	A	270	GLU	2.2
2	B	908	TYR	2.2
2	B	604	LEU	2.1
2	B	832	ASN	2.1
1	A	56	ASP	2.1
1	A	350	SER	2.1
1	A	53	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	B	1204	6/6	0.69	0.21	70,70,70,71	0
7	GOL	B	1205	6/6	0.72	0.22	51,51,52,52	0
8	PEO	B	1208	2/2	0.75	0.17	64,64,64,64	0
10	PEG	B	1206	7/7	0.76	0.18	51,51,52,52	0
7	GOL	A	405	6/6	0.77	0.16	64,65,65,65	0
8	PEO	B	1207	2/2	0.78	0.27	48,48,48,48	0
9	SO4	B	1201	5/5	0.79	0.15	82,82,83,83	0
8	PEO	B	1209	2/2	0.81	0.20	66,66,66,66	0
6	TRS	B	1203	8/8	0.85	0.12	42,42,42,43	0
8	PEO	A	408	2/2	0.87	0.16	47,47,47,47	0
8	PEO	A	407	2/2	0.87	0.23	37,37,37,37	0
8	PEO	A	409	2/2	0.88	0.14	47,47,47,47	0
8	PEO	A	410	2/2	0.90	0.10	54,54,54,54	0
8	PEO	A	406	2/2	0.90	0.10	38,38,38,39	0
6	TRS	A	404	8/8	0.90	0.09	29,30,31,31	0
5	LAB	A	403	27/27	0.96	0.07	25,27,29,29	0
9	SO4	B	1202	5/5	0.97	0.06	33,34,34,35	0
4	MG	A	402	1/1	0.98	0.04	20,20,20,20	0
3	ADP	A	401	27/27	0.98	0.05	18,20,22,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LAB A 403:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.