



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 12:04 PM UTC

PDB ID : 2BK4 / pdb_00002bk4
Title : Human Monoamine Oxidase B: I199F mutant in complex with rasagiline
Authors : Binda, C.; Edmondson, D.E.; Mattevi, A.; Hubalek, F.; Khalil, A.; Li, M.; Castagnoli, N.
Deposited on : 2005-02-10
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	520	 90% 6% • •
1	B	520	 88% 7% • 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8594 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMINE OXIDASE [FLAVIN-CONTAINING] B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	0	0
			3974	2541	681	728	24			
1	B	494	Total	C	N	O	S	0	0	0
			3943	2522	676	721	24			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	PHE	ILE	engineered mutation	UNP P27338
B	199	PHE	ILE	engineered mutation	UNP P27338

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is (1R)-N-(prop-2-en-1-yl)-2,3-dihydro-1H-inden-1-amine (CCD ID: RAS) (formula: C₁₂H₁₅N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			13	12	1		
3	B	1	Total	C	N	0	0
			13	12	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	249	Total	O	0	0
			249	249		
4	B	296	Total	O	0	0
			296	296		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



• Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.03Å 223.52Å 86.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90	Depositor
% Data completeness (in resolution range)	97.1 (15.00-1.90)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.191 , 0.218	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	8594	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, RAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	1/4072 (0.0%)	0.76	0/5527
1	B	0.60	1/4041 (0.0%)	0.76	0/5484
All	All	0.61	2/8113 (0.0%)	0.76	0/11011

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	489	VAL	CA-CB	5.82	1.57	1.54
1	A	489	VAL	CA-CB	5.34	1.57	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3974	0	3965	29	0
1	B	3943	0	3935	31	0
2	A	53	0	29	3	0
2	B	53	0	29	2	0
3	A	13	0	14	2	0
3	B	13	0	14	2	0
4	A	249	0	0	3	2

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	296	0	0	5	0
All	All	8594	0	7986	52	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.49	0.95
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.51	0.92
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.32	0.77
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.34	0.76
1:A:353:ALA:HB1	1:B:280:MET:HE1	1.71	0.71
1:A:251:ASN:HD22	1:A:251:ASN:H	1.37	0.70
1:B:251:ASN:HD22	1:B:251:ASN:H	1.43	0.67
1:A:280:MET:HE1	1:B:353:ALA:HB1	1.79	0.64
1:B:271:LYS:HE2	4:B:2239:HOH:O	2.01	0.60
1:A:445:ARG:HD2	1:A:463:TRP:CH2	2.40	0.56
1:A:445:ARG:HD2	1:A:463:TRP:CZ2	2.41	0.56
1:A:47:ARG:HD3	4:A:2121:HOH:O	2.06	0.55
1:B:445:ARG:HD2	1:B:463:TRP:CH2	2.41	0.55
1:B:445:ARG:HD2	1:B:463:TRP:CZ2	2.41	0.55
1:B:47:ARG:HD3	4:B:2160:HOH:O	2.08	0.54
1:B:412:ARG:HG2	4:B:2249:HOH:O	2.10	0.51
1:A:410:TYR:OH	1:B:350:ARG:HD3	2.10	0.50
1:A:251:ASN:H	1:A:251:ASN:ND2	2.07	0.50
1:A:350:ARG:HD3	1:B:410:TYR:OH	2.11	0.50
1:A:445:ARG:HD3	4:A:2218:HOH:O	2.11	0.49
1:A:412:ARG:HD2	4:A:2202:HOH:O	2.12	0.48
1:A:451:LEU:HD23	1:A:454:MET:CE	2.44	0.48
1:A:58:GLY:HA2	2:A:600:FAD:C4X	2.44	0.48
1:A:321:GLU:H	1:A:321:GLU:CD	2.21	0.48
1:A:248:GLU:OE2	1:B:252:HIS:NE2	2.45	0.47
1:A:252:HIS:NE2	1:B:248:GLU:OE2	2.47	0.47
1:B:445:ARG:HD3	4:B:2267:HOH:O	2.13	0.47
1:B:387:ASN:O	1:B:390:GLU:HG2	2.15	0.47
1:B:451:LEU:HD23	1:B:454:MET:CE	2.45	0.46
1:B:122:MET:CE	1:B:185:PHE:HE2	2.29	0.45
1:B:410:TYR:O	1:B:413:VAL:HG13	2.17	0.45
1:A:270:MET:HE1	1:B:268:LEU:HD23	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:GLY:HA2	2:B:600:FAD:C4X	2.47	0.45
2:A:600:FAD:C4	3:A:601:RAS:H24	2.48	0.44
1:A:451:LEU:HD23	1:A:454:MET:HE1	1.99	0.44
1:B:251:ASN:H	1:B:251:ASN:ND2	2.11	0.44
1:B:321:GLU:H	1:B:321:GLU:CD	2.26	0.43
1:A:171:LEU:HD21	3:A:601:RAS:H21	2.01	0.43
1:B:280:MET:HB3	1:B:280:MET:HE3	1.84	0.43
1:A:389:CYS:CB	1:B:280:MET:HG3	2.49	0.42
1:A:434:GLY:O	2:A:600:FAD:H1'2	2.20	0.42
2:B:600:FAD:C4	3:B:601:RAS:H24	2.48	0.42
1:B:171:LEU:HD21	3:B:601:RAS:H21	2.02	0.42
1:A:265:PRO:HD2	1:A:268:LEU:HD12	2.02	0.42
1:B:163:GLN:HG2	4:B:2122:HOH:O	2.19	0.42
1:B:451:LEU:HD23	1:B:454:MET:HE1	2.01	0.41
1:A:268:LEU:HD23	1:B:270:MET:HE1	2.02	0.41
1:A:280:MET:HE3	1:A:280:MET:HB3	1.84	0.41
1:B:233:ARG:HG3	1:B:251:ASN:HD21	1.86	0.41
1:A:280:MET:HG3	1:B:389:CYS:CB	2.51	0.40
1:A:400:THR:HB	1:A:427:GLU:HG2	2.03	0.40
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.11	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2017:HOH:O	4:A:2017:HOH:O[3_655]	1.54	0.66
4:A:2092:HOH:O	4:A:2092:HOH:O[6_565]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/520 (96%)	487 (98%)	10 (2%)	0	100	100
1	B	492/520 (95%)	482 (98%)	10 (2%)	0	100	100
All	All	989/1040 (95%)	969 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/444 (96%)	418 (98%)	9 (2%)	47	44
1	B	424/444 (96%)	413 (97%)	11 (3%)	40	35
All	All	851/888 (96%)	831 (98%)	20 (2%)	43	38

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	LYS
1	A	209	LYS
1	A	251	ASN
1	A	280	MET
1	A	321	GLU
1	A	379	GLU
1	A	413	VAL
1	A	459	GLU
1	A	495	LEU
1	B	95	LYS
1	B	190	LYS
1	B	209	LYS
1	B	251	ASN
1	B	280	MET
1	B	321	GLU
1	B	379	GLU
1	B	397	CYS
1	B	413	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	459	GLU
1	B	495	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	ASN
1	A	251	ASN
1	B	117	ASN
1	B	251	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	RAS	B	601	2	13,14,14	0.72	0	15,18,18	1.51	3 (20%)
2	FAD	A	600	3,1	58,58,58	1.21	6 (10%)	85,89,89	1.61	14 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	RAS	A	601	2	13,14,14	0.69	0	15,18,18	1.34	2 (13%)
2	FAD	B	600	3,1	58,58,58	1.14	4 (6%)	85,89,89	1.66	16 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RAS	B	601	2	-	3/4/13/13	0/2/2/2
2	FAD	A	600	3,1	-	2/34/50/50	0/6/6/6
3	RAS	A	601	2	-	3/4/13/13	0/2/2/2
2	FAD	B	600	3,1	-	2/34/50/50	0/6/6/6

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	FAD	C4X-N5	4.29	1.40	1.30
2	B	600	FAD	C4X-N5	4.23	1.39	1.30
2	A	600	FAD	C10-N1	3.20	1.39	1.33
2	B	600	FAD	C10-N1	3.14	1.39	1.33
2	A	600	FAD	C2A-N1A	2.75	1.38	1.33
2	A	600	FAD	P-O3P	2.73	1.62	1.59
2	A	600	FAD	C2A-N3A	2.56	1.38	1.33
2	A	600	FAD	C8A-N7A	2.37	1.36	1.31
2	B	600	FAD	C2A-N1A	2.28	1.38	1.33
2	B	600	FAD	C2A-N3A	2.20	1.37	1.33

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	FAD	N3A-C2A-N1A	-5.98	119.53	128.58
2	B	600	FAD	N3A-C2A-N1A	-5.84	119.75	128.58
2	B	600	FAD	C9A-C5X-N5	-4.96	117.19	122.45
2	A	600	FAD	C9A-C5X-N5	-4.10	118.11	122.45
2	B	600	FAD	N9A-C8A-N7A	-4.05	108.19	113.94
2	B	600	FAD	C5A-C4A-N3A	-3.84	121.42	126.72
2	A	600	FAD	N9A-C8A-N7A	-3.82	108.52	113.94
2	A	600	FAD	C5A-C4A-N3A	-3.67	121.67	126.72
2	B	600	FAD	C2A-N3A-C4A	3.30	119.90	111.83
2	A	600	FAD	C5A-N7A-C8A	3.28	108.61	103.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	FAD	C5A-N7A-C8A	3.27	108.58	103.45
2	A	600	FAD	C2A-N3A-C4A	3.16	119.54	111.83
2	B	600	FAD	C9A-N10-C10	-3.13	115.98	120.75
2	A	600	FAD	C9A-N10-C10	-3.01	116.16	120.75
2	B	600	FAD	C4-N3-C2	-3.01	120.30	125.64
3	A	601	RAS	C11-N10-C9	2.95	119.96	113.86
2	B	600	FAD	C4X-C4-N3	2.89	120.61	113.25
2	A	600	FAD	C4-N3-C2	-2.84	120.60	125.64
2	A	600	FAD	C4X-C4-N3	2.77	120.32	113.25
3	B	601	RAS	C11-N10-C9	2.72	119.47	113.86
3	B	601	RAS	C6-C7-C8	-2.66	117.81	120.99
2	B	600	FAD	O4-C4-C4X	-2.61	119.65	126.53
2	B	600	FAD	N3A-C4A-N9A	2.54	131.49	127.17
2	A	600	FAD	C4A-C5A-N7A	-2.49	107.73	110.58
2	A	600	FAD	C5'-C4'-C3'	-2.43	107.64	112.22
2	B	600	FAD	C5'-C4'-C3'	-2.39	107.71	112.22
2	A	600	FAD	C10-C4X-N5	-2.33	120.06	124.81
2	B	600	FAD	C4A-N9A-C8A	2.33	108.18	105.74
2	B	600	FAD	C4A-C5A-N7A	-2.30	107.95	110.58
2	B	600	FAD	C10-C4X-N5	-2.30	120.12	124.81
2	A	600	FAD	O4-C4-C4X	-2.24	120.61	126.53
3	A	601	RAS	C6-C7-C8	-2.18	118.38	120.99
2	B	600	FAD	C4X-C10-N1	-2.11	119.41	124.59
2	A	600	FAD	N3A-C4A-N9A	2.10	130.74	127.17
3	B	601	RAS	C5-C4-C3	-2.07	117.84	120.88

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	601	RAS	C1-C9-N10-C11
3	A	601	RAS	C8-C9-N10-C11
3	B	601	RAS	C1-C9-N10-C11
3	B	601	RAS	C8-C9-N10-C11
3	A	601	RAS	N10-C11-C12-C13
3	B	601	RAS	N10-C11-C12-C13
2	B	600	FAD	PA-O3P-P-O5'
2	A	600	FAD	C2'-C1'-N10-C10
2	B	600	FAD	C2'-C1'-N10-C10
2	A	600	FAD	C2B-C1B-N9A-C8A


There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	601	RAS	2	0
2	A	600	FAD	3	0
3	A	601	RAS	2	0
2	B	600	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.