



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:38 AM UTC

PDB ID : 4BKE / pdb_00004bke
Title : Recombinant human serum albumin with palmitic acid. Synthetic cationic antimicrobial peptides bind with their hydrophobic parts to drug site II of human serum albumin
Authors : Sivertsen, A.; Isaksson, J.; Leiros, H.-K.S.; Svenson, J.; Svendsen, J.-S.; Brandsdal, B.-O.
Deposited on : 2013-04-24
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4793 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERUM ALBUMIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	183	2	0
			4648	2933	786	888	41			

- Molecule 2 is PALMITIC ACID (CCD ID: PLM) (formula: $C_{16}H_{32}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	11	2		
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	0	0
			17	15	2		
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	0	0
			17	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			18	16	2		
2	A	1	Total	C	O	2	0
			18	16	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total	O	0	0
			26	26		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SERUM ALBUMIN

Chain A:  76% 17% . .



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	189.37Å 39.00Å 96.30Å 90.00° 105.48° 90.00°	Depositor
Resolution (Å)	20.00 – 2.35 20.00 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.6 (20.00-2.35) 96.4 (20.00-2.35)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.35Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.198 , 0.254 0.194 , 0.249	Depositor DCC
R_{free} test set	2079 reflections (7.21%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 42.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4793	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.89	44/4738 (0.9%)	1.71	67/6390 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	396	GLU	CD-OE1	-38.83	0.51	1.25
1	A	372	LYS	CA-CB	-34.34	1.06	1.53
1	A	137	LYS	CD-CE	-31.59	0.57	1.52
1	A	519	LYS	CG-CD	-29.33	0.64	1.52
1	A	204	GLN	CB-CG	-29.07	0.65	1.52
1	A	205	LYS	CD-CE	-28.29	0.67	1.52
1	A	170	GLN	CG-CD	-26.48	0.85	1.52
1	A	244	GLU	CD-OE1	-25.62	0.76	1.25
1	A	396	GLU	CD-OE2	23.63	1.70	1.25
1	A	570	GLU	CA-CB	-23.38	1.16	1.53
1	A	500	LYS	CE-NZ	-22.55	0.81	1.49
1	A	41	LYS	CE-NZ	-21.93	0.83	1.49
1	A	396	GLU	CG-CD	-21.06	0.99	1.52
1	A	277	GLU	CG-CD	-20.92	0.99	1.52
1	A	389	LYS	CG-CD	-20.55	0.90	1.52
1	A	466	LYS	CD-CE	-18.18	0.97	1.52
1	A	240	LYS	CG-CD	-17.09	1.01	1.52
1	A	294	GLU	CG-CD	16.92	1.94	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	560	LYS	CB-CG	-16.80	1.02	1.52
1	A	538	LYS	CE-NZ	-16.25	1.00	1.49
1	A	184	GLU	CD-OE1	15.59	1.54	1.25
1	A	6	GLU	CD-OE1	15.29	1.54	1.25
1	A	536	LYS	CG-CD	-14.85	1.07	1.52
1	A	444	LYS	CB-CG	14.70	1.96	1.52
1	A	109	ASN	CB-CG	-14.45	1.16	1.52
1	A	313	LYS	CA-CB	-13.94	1.30	1.53
1	A	459	GLN	CB-CG	-13.72	1.11	1.52
1	A	276	LYS	CB-CG	-13.35	1.12	1.52
1	A	524	LYS	CG-CD	-12.89	1.13	1.52
1	A	244	GLU	CD-OE2	11.92	1.48	1.25
1	A	184	GLU	CD-OE2	-10.58	1.05	1.25
1	A	351	LYS	CD-CE	-8.54	1.26	1.52
1	A	170	GLN	CB-CG	8.16	1.76	1.52
1	A	466	LYS	CG-CD	-8.08	1.28	1.52
1	A	111	ASN	CB-CG	-7.90	1.32	1.52
1	A	301	ASP	CA-CB	7.58	1.64	1.53
1	A	545	LYS	CE-NZ	-7.19	1.27	1.49
1	A	390	GLN	CB-CG	-7.14	1.31	1.52
1	A	402	LYS	CE-NZ	-6.71	1.29	1.49
1	A	57	GLU	CB-CG	-6.48	1.33	1.52
1	A	82	GLU	CB-CG	5.99	1.70	1.52
1	A	3	HIS	CA-CB	5.90	1.65	1.53
1	A	518	GLU	CB-CG	-5.43	1.36	1.52
1	A	436	LYS	CG-CD	-5.11	1.37	1.52

All (67) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	519	LYS	CG-CD-CE	-35.66	29.29	111.30
1	A	277	GLU	CB-CG-CD	-27.99	65.02	112.60
1	A	109	ASN	CA-CB-CG	-27.49	85.11	112.60
1	A	536	LYS	CG-CD-CE	23.83	166.12	111.30
1	A	519	LYS	CB-CG-CD	-23.28	57.75	111.30
1	A	521	ARG	CB-CG-CD	-22.25	60.13	111.30
1	A	570	GLU	CB-CA-C	21.48	146.31	110.86
1	A	244	GLU	OE1-CD-OE2	21.31	174.04	122.90
1	A	570	GLU	N-CA-CB	-20.39	80.10	110.07
1	A	396	GLU	OE1-CD-OE2	20.00	170.90	122.90
1	A	317	LYS	CB-CG-CD	-18.78	68.11	111.30
1	A	444	LYS	CA-CB-CG	-18.62	76.86	114.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	277	GLU	CG-CD-OE2	-18.36	76.17	118.40
1	A	396	GLU	CG-CD-OE2	-18.17	76.61	118.40
1	A	244	GLU	CG-CD-OE2	-18.01	76.98	118.40
1	A	205	LYS	CG-CD-CE	17.93	152.54	111.30
1	A	109	ASN	CB-CG-ND2	17.35	142.42	116.40
1	A	277	GLU	CG-CD-OE1	17.14	157.82	118.40
1	A	137	LYS	CG-CD-CE	-15.72	75.14	111.30
1	A	389	LYS	CB-CG-CD	15.29	146.46	111.30
1	A	317	LYS	CG-CD-CE	-15.04	76.70	111.30
1	A	111	ASN	N-CA-CB	14.19	127.85	110.53
1	A	536	LYS	CB-CG-CD	14.07	143.66	111.30
1	A	538	LYS	CD-CE-NZ	-13.33	69.25	111.90
1	A	109	ASN	CB-CG-OD1	-13.27	94.26	120.80
1	A	570	GLU	CA-CB-CG	-12.41	89.29	114.10
1	A	372	LYS	CB-CA-C	12.26	130.87	111.13
1	A	3	HIS	CB-CA-C	-11.91	87.47	110.10
1	A	294	GLU	CB-CG-CD	-11.77	92.59	112.60
1	A	396	GLU	CB-CG-CD	11.70	132.49	112.60
1	A	6	GLU	OE1-CD-OE2	-11.19	96.04	122.90
1	A	372	LYS	N-CA-CB	-10.63	93.74	111.17
1	A	372	LYS	CA-CB-CG	10.61	135.32	114.10
1	A	170	GLN	CA-CB-CG	-10.54	93.03	114.10
1	A	560	LYS	CB-CG-CD	-10.34	87.51	111.30
1	A	276	LYS	CB-CG-CD	-10.12	88.03	111.30
1	A	481	LEU	N-CA-CB	-9.76	93.73	110.32
1	A	402	LYS	CD-CE-NZ	9.62	142.67	111.90
1	A	276	LYS	CA-CB-CG	-9.36	95.37	114.10
1	A	556	GLU	CB-CG-CD	9.15	128.16	112.60
1	A	378	LYS	CB-CG-CD	9.12	132.26	111.30
1	A	137	LYS	CD-CE-NZ	-9.06	82.92	111.90
1	A	466	LYS	CG-CD-CE	-9.06	90.47	111.30
1	A	498	VAL	CG1-CB-CG2	8.86	130.28	110.80
1	A	240	LYS	CB-CG-CD	8.75	131.41	111.30
1	A	524	LYS	CB-CG-CD	8.60	131.08	111.30
1	A	389	LYS	CG-CD-CE	8.24	130.24	111.30
1	A	184	GLU	CG-CD-OE2	8.13	137.09	118.40
1	A	170	GLN	CB-CG-CD	-8.04	98.93	112.60
1	A	436	LYS	CB-CG-CD	7.96	129.62	111.30
1	A	60	GLU	CA-CB-CG	7.80	129.69	114.10
1	A	466	LYS	CB-CG-CD	7.63	128.84	111.30
1	A	311	GLU	CB-CG-CD	-7.58	99.71	112.60
1	A	351	LYS	CG-CD-CE	7.30	128.10	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	311	GLU	CA-CB-CG	-7.11	99.88	114.10
1	A	301	ASP	CA-CB-CG	6.70	119.30	112.60
1	A	390	GLN	CA-CB-CG	6.64	127.39	114.10
1	A	536	LYS	CD-CE-NZ	-6.44	91.30	111.90
1	A	244	GLU	CG-CD-OE1	-6.41	103.66	118.40
1	A	168	CYS	N-CA-C	6.26	119.08	111.82
1	A	466	LYS	CD-CE-NZ	-6.03	92.60	111.90
1	A	3	HIS	CA-CB-CG	-5.78	108.02	113.80
1	A	178	LEU	N-CA-C	5.74	117.33	111.14
1	A	301	ASP	N-CA-CB	-5.27	104.44	110.35
1	A	360	CYS	N-CA-C	5.11	116.84	111.28
1	A	4	LYS	CA-CB-CG	5.05	124.19	114.10
1	A	262	LYS	CG-CD-CE	5.04	122.89	111.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	6	GLU	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4648	0	4561	33	0
2	A	119	0	194	3	0
3	A	26	0	0	3	0
All	All	4793	0	4755	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:GLU:HG3	1:A:384:PRO:HD3	1.75	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:VAL:HG22	1:A:256:ASP:HB3	1.77	0.65
1:A:272:SER:HB3	1:A:275:LEU:HD22	1.79	0.65
1:A:460:LEU:HD11	2:A:1004:PLM:HC1	1.77	0.64
1:A:108:ASP:HB3	1:A:148:TYR:CE2	2.39	0.58
1:A:519:LYS:O	1:A:523:ILE:HG12	2.07	0.54
1:A:216:VAL:CG2	1:A:235:VAL:HG21	2.38	0.53
1:A:408:LEU:HD13	1:A:427:SER:HB2	1.91	0.53
1:A:195:LYS:HG3	1:A:455:VAL:HG11	1.90	0.52
1:A:426:VAL:HG11	1:A:460:LEU:HD13	1.92	0.51
1:A:81:ARG:NH2	1:A:89:ASP:OD1	2.42	0.51
1:A:415:VAL:HG12	1:A:418:VAL:HG23	1.93	0.51
1:A:312:SER:HB3	1:A:315:VAL:HG23	1.92	0.51
1:A:7:VAL:HG22	3:A:2001:HOH:O	2.12	0.50
1:A:216:VAL:HG22	1:A:235:VAL:HG21	1.94	0.50
1:A:108:ASP:HB3	1:A:148:TYR:HE2	1.76	0.50
1:A:383:GLU:HG3	1:A:384:PRO:CD	2.42	0.49
1:A:408:LEU:HD13	1:A:427:SER:CB	2.42	0.49
1:A:101:CYS:O	1:A:105:HIS:HD2	1.96	0.49
1:A:576:VAL:HG12	1:A:580:GLN:HE21	1.78	0.48
1:A:507:PHE:CD1	2:A:1005:PLM:HB2	2.50	0.47
1:A:29[B]:GLN:HB2	1:A:147:PRO:HA	1.97	0.46
1:A:107:ASP:OD1	1:A:108:ASP:O	2.34	0.45
1:A:224:PRO:HD2	1:A:296:ASP:HB3	1.99	0.45
1:A:344:VAL:HG22	1:A:450:GLU:OE2	2.17	0.45
1:A:348:ARG:HG2	1:A:482:VAL:HG12	1.99	0.44
1:A:268:GLN:HG2	3:A:2015:HOH:O	2.17	0.44
1:A:119:GLU:HG3	3:A:2010:HOH:O	2.17	0.43
2:A:1003:PLM:H81	2:A:1004:PLM:H22	2.01	0.43
1:A:302:LEU:HA	1:A:303:PRO:HD3	1.88	0.42
1:A:307:ALA:HA	1:A:311:GLU:HB2	2.01	0.42
1:A:156:PHE:HZ	1:A:284:LEU:O	2.04	0.41
1:A:417:GLN:HB2	1:A:469:VAL:HG13	2.02	0.41
1:A:348:ARG:CG	1:A:482:VAL:HG12	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/609 (96%)	559 (96%)	21 (4%)	2 (0%)	36	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	561	ALA
1	A	322	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	511/533 (96%)	481 (94%)	30 (6%)	18	22

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	14	LEU
1	A	57	GLU
1	A	74	LEU
1	A	97	GLU
1	A	203	LEU
1	A	225	LYS
1	A	238	LEU
1	A	275	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	283	LEU
1	A	297	GLU
1	A	313	LYS
1	A	359	LYS
1	A	380	LEU
1	A	383	GLU
1	A	389	LYS
1	A	408	LEU
1	A	419	SER
1	A	430	LEU
1	A	451	ASP
1	A	457	LEU
1	A	469	VAL
1	A	496	THR
1	A	498	VAL
1	A	544	LEU
1	A	545	LYS
1	A	560	LYS
1	A	562	ASP
1	A	566	THR
1	A	573	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	HIS
1	A	61	ASN
1	A	99	ASN
1	A	105	HIS
1	A	146	HIS
1	A	242	HIS
1	A	318	ASN
1	A	397	GLN
1	A	580	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLM	A	1001	-	12,12,17	0.66	0	12,12,17	0.92	0
2	PLM	A	1007	-	17,17,17	3.80	1 (5%)	17,17,17	1.58	1 (5%)
2	PLM	A	1006	-	17,17,17	0.47	0	17,17,17	1.02	0
2	PLM	A	1003	-	16,16,17	0.55	0	16,16,17	1.06	1 (6%)
2	PLM	A	1004	-	17,17,17	0.51	0	17,17,17	1.03	1 (5%)
2	PLM	A	1002	-	17,17,17	0.56	0	17,17,17	0.91	1 (5%)
2	PLM	A	1005	-	16,16,17	0.53	0	16,16,17	0.92	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLM	A	1001	-	-	8/10/10/15	-
2	PLM	A	1007	-	-	12/15/15/15	-
2	PLM	A	1006	-	-	9/15/15/15	-
2	PLM	A	1003	-	-	8/14/14/15	-
2	PLM	A	1004	-	-	9/15/15/15	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLM	A	1002	-	-	12/15/15/15	-
2	PLM	A	1005	-	-	8/14/14/15	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1007	PLM	CF-CE	15.51	2.47	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1007	PLM	CF-CE-CD	-5.39	67.97	115.25
2	A	1003	PLM	C3-C2-C1	-2.39	108.27	114.51
2	A	1004	PLM	C3-C2-C1	-2.25	108.64	114.51
2	A	1005	PLM	O1-C1-C2	2.10	120.62	114.00
2	A	1002	PLM	O1-C1-C2	2.08	120.58	114.00

There are no chirality outliers.

All (66) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1007	PLM	C1-C2-C3-C4
2	A	1004	PLM	C1-C2-C3-C4
2	A	1006	PLM	CC-CD-CE-CF
2	A	1002	PLM	C2-C3-C4-C5
2	A	1006	PLM	C6-C7-C8-C9
2	A	1006	PLM	C5-C6-C7-C8
2	A	1002	PLM	C7-C8-C9-CA
2	A	1005	PLM	C9-CA-CB-CC
2	A	1002	PLM	C6-C7-C8-C9
2	A	1003	PLM	CA-CB-CC-CD
2	A	1001	PLM	C6-C7-C8-C9
2	A	1004	PLM	CC-CD-CE-CF
2	A	1001	PLM	C1-C2-C3-C4
2	A	1007	PLM	C5-C6-C7-C8
2	A	1002	PLM	C8-C9-CA-CB
2	A	1004	PLM	C2-C3-C4-C5
2	A	1003	PLM	C1-C2-C3-C4
2	A	1002	PLM	C5-C6-C7-C8
2	A	1007	PLM	CA-CB-CC-CD
2	A	1006	PLM	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1007	PLM	C2-C3-C4-C5
2	A	1001	PLM	C2-C3-C4-C5
2	A	1007	PLM	CB-CC-CD-CE
2	A	1005	PLM	C5-C6-C7-C8
2	A	1006	PLM	C9-CA-CB-CC
2	A	1003	PLM	CB-CC-CD-CE
2	A	1002	PLM	C9-CA-CB-CC
2	A	1005	PLM	C7-C8-C9-CA
2	A	1003	PLM	C9-CA-CB-CC
2	A	1001	PLM	C3-C4-C5-C6
2	A	1002	PLM	CA-CB-CC-CD
2	A	1007	PLM	C4-C5-C6-C7
2	A	1007	PLM	C9-CA-CB-CC
2	A	1007	PLM	C7-C8-C9-CA
2	A	1003	PLM	C6-C7-C8-C9
2	A	1007	PLM	C8-C9-CA-CB
2	A	1002	PLM	CC-CD-CE-CF
2	A	1006	PLM	C1-C2-C3-C4
2	A	1004	PLM	CB-CC-CD-CE
2	A	1004	PLM	C3-C4-C5-C6
2	A	1002	PLM	CB-CC-CD-CE
2	A	1006	PLM	C3-C4-C5-C6
2	A	1005	PLM	CA-CB-CC-CD
2	A	1004	PLM	CD-CE-CF-CG
2	A	1004	PLM	C8-C9-CA-CB
2	A	1002	PLM	C4-C5-C6-C7
2	A	1002	PLM	O2-C1-C2-C3
2	A	1002	PLM	O1-C1-C2-C3
2	A	1005	PLM	O2-C1-C2-C3
2	A	1003	PLM	O1-C1-C2-C3
2	A	1001	PLM	C4-C5-C6-C7
2	A	1003	PLM	O2-C1-C2-C3
2	A	1005	PLM	O1-C1-C2-C3
2	A	1001	PLM	O1-C1-C2-C3
2	A	1001	PLM	C5-C6-C7-C8
2	A	1004	PLM	O1-C1-C2-C3
2	A	1004	PLM	O2-C1-C2-C3
2	A	1007	PLM	O1-C1-C2-C3
2	A	1001	PLM	O2-C1-C2-C3
2	A	1006	PLM	O1-C1-C2-C3
2	A	1006	PLM	O2-C1-C2-C3
2	A	1007	PLM	O2-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1003	PLM	C4-C5-C6-C7
2	A	1005	PLM	C4-C5-C6-C7
2	A	1005	PLM	C6-C7-C8-C9
2	A	1007	PLM	CD-CE-CF-CG


There are no ring outliers.


3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1003	PLM	1	0
2	A	1004	PLM	2	0
2	A	1005	PLM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	582/609 (95%)	-0.19	2 (0%) 90 92	23, 56, 80, 94	77 (13%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	561	ALA	6.6
1	A	562	ASP	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PLM	A	1001	13/18	0.80	0.16	72,74,77,77	0
2	PLM	A	1007	18/18	0.82	0.17	73,79,88,89	2
2	PLM	A	1002	18/18	0.89	0.10	55,59,65,65	0
2	PLM	A	1006	18/18	0.91	0.10	56,59,72,72	0
2	PLM	A	1004	18/18	0.91	0.12	56,59,61,61	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PLM	A	1005	17/18	0.94	0.10	59,62,68,68	0
2	PLM	A	1003	17/18	0.95	0.11	49,56,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PLM A 1001:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLM A 1007:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PLM A 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PLM A 1006:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around PLM A 1004:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PLM A 1005:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.