



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:03 AM UTC

PDB ID : 1BLI / pdb_00001bli
Title : BACILLUS LICHENIFORMIS ALPHA-AMYLASE
Authors : Machius, M.; Declerck, N.; Huber, R.; Wiegand, G.
Deposited on : 1998-01-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	483	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-AMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	481	Total	C	N	O	S	0	0	0
			3905	2491	678	729	7			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	LEU	ARG	conflict	UNP P06278
A	190	PHE	ASN	engineered mutation	UNP P06278
A	264	SER	GLN	engineered mutation	UNP P06278
A	265	TYR	ASN	engineered mutation	UNP P06278
A	310	GLY	SER	conflict	UNP P06278
A	320	SER	ALA	conflict	UNP P06278
A	465	ALA	GLU	conflict	UNP P06278

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	221	Total	O	0	0
			221	221		

Note EDS was not executed.

Chain A: 84% 13% 3%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	91.30Å 91.30Å 137.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.90	Depositor
% Data completeness (in resolution range)	89.4 (30.00-1.90)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.154 , 0.185	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4130	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/4029 (0.0%)	0.98	13/5472 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	286	VAL	CA-CB	5.02	1.57	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	VAL	N-CA-C	-7.24	104.79	111.67
1	A	21	HIS	N-CA-C	6.37	119.03	111.33
1	A	14	TYR	N-CA-C	6.25	119.93	112.93
1	A	204	ASP	N-CA-C	-5.95	105.85	113.23
1	A	12	GLU	N-CA-C	-5.87	100.42	109.52
1	A	229	ARG	N-CA-C	-5.60	99.29	108.41
1	A	415	GLY	N-CA-C	-5.58	106.04	112.29
1	A	77	TYR	N-CA-C	5.50	119.70	112.89
1	A	198	TYR	CB-CA-C	-5.37	109.40	115.79
1	A	413	ARG	N-CA-C	-5.36	99.70	108.76
1	A	12	GLU	N-CA-CB	-5.32	102.30	111.55
1	A	131	GLY	N-CA-C	-5.19	104.66	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	373	SER	N-CA-C	-5.09	102.46	110.36

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	195	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3905	0	3645	36	0
2	A	3	0	0	0	0
3	A	1	0	0	0	0
4	A	221	0	0	2	0
All	All	4130	0	3645	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:389:LYS:HE2	1:A:393:GLN:HE22	1.29	0.96
1:A:389:LYS:HG3	1:A:393:GLN:NE2	1.87	0.89
1:A:389:LYS:HG3	1:A:393:GLN:HE21	1.38	0.88
1:A:458:GLU:H	1:A:458:GLU:CD	1.81	0.84
1:A:450:HIS:HB3	1:A:483:ARG:NH1	1.96	0.80
1:A:459:PRO:HB3	1:A:483:ARG:HH12	1.49	0.78
1:A:450:HIS:HB3	1:A:483:ARG:HH12	1.51	0.76
1:A:389:LYS:CE	1:A:393:GLN:HE22	2.03	0.71
1:A:9:GLN:HE22	1:A:326:ASN:HB2	1.58	0.67
1:A:303:ASP:HB3	1:A:306:LYS:HD2	1.77	0.66
1:A:389:LYS:HG2	1:A:452:ILE:HD11	1.85	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:459:PRO:HB3	1:A:483:ARG:NH1	2.16	0.58
1:A:389:LYS:HE2	1:A:393:GLN:NE2	2.10	0.58
1:A:35:HIS:CE1	1:A:385:GLU:HG2	2.40	0.56
1:A:250:GLU:HG3	1:A:251:LYS:N	2.19	0.55
1:A:381:LYS:HG2	1:A:382:HIS:N	2.21	0.54
1:A:24:ARG:NH1	4:A:2012:HOH:O	2.40	0.54
1:A:164:ASP:HA	1:A:173:ARG:O	2.09	0.53
1:A:424:LEU:C	1:A:424:LEU:HD12	2.33	0.53
1:A:458:GLU:CD	1:A:458:GLU:N	2.61	0.52
1:A:101:VAL:HB	1:A:230:LEU:HD12	1.93	0.49
1:A:68:HIS:HD2	4:A:1028:HOH:O	1.95	0.48
1:A:184:TRP:CD2	1:A:185:GLU:HA	2.48	0.48
1:A:34:GLU:HG2	1:A:35:HIS:CD2	2.48	0.47
1:A:205:HIS:CE1	1:A:207:ASP:HB2	2.50	0.46
1:A:233:VAL:HG11	1:A:260:ALA:HB1	1.98	0.45
1:A:435:ALA:HB2	1:A:471:HIS:CE1	2.52	0.44
1:A:176:LYS:HG3	1:A:182:TRP:CZ2	2.54	0.42
1:A:389:LYS:CG	1:A:393:GLN:NE2	2.71	0.42
1:A:69:GLN:HG2	1:A:145:GLY:HA3	2.02	0.42
1:A:41:TRP:CD1	1:A:41:TRP:C	2.97	0.41
1:A:399:GLN:HA	1:A:412:THR:O	2.20	0.41
1:A:184:TRP:CG	1:A:185:GLU:HA	2.55	0.41
1:A:327:HIS:O	1:A:333:GLN:HG3	2.20	0.41
1:A:442:ARG:HH11	1:A:443:GLN:HA	1.85	0.41
1:A:70:LYS:HA	1:A:70:LYS:HD2	1.77	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/483 (99%)	466 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	401/402 (100%)	375 (94%)	26 (6%)	15 8

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	23	LYS
1	A	28	ASP
1	A	34	GLU
1	A	41	TRP
1	A	46	TYR
1	A	70	LYS
1	A	88	LYS
1	A	134	LEU
1	A	142	HIS
1	A	176	LYS
1	A	230	LEU
1	A	250	GLU
1	A	251	LYS
1	A	255	GLU
1	A	315	LYS
1	A	372	ASP
1	A	381	LYS
1	A	383	LYS
1	A	437	ARG
1	A	442	ARG
1	A	450	HIS
1	A	456	ARG
1	A	457	SER
1	A	458	GLU
1	A	483	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	27	ASN
1	A	35	HIS
1	A	68	HIS
1	A	96	ASN
1	A	178	GLN
1	A	221	ASN
1	A	278	ASN
1	A	293	HIS
1	A	393	GLN
1	A	406	HIS
1	A	421	ASN
1	A	482	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.