



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 11:54 PM UTC

PDB ID : 4BLD / pdb\_00004bld  
Title : Crystal structure of a human Suppressor of fused (SUFU)-GLI3p complex  
Authors : Cherry, A.L.; Finta, C.; Karlstrom, M.; De Sanctis, D.; Toftgard, R.; Jovine, L.  
Deposited on : 2013-05-02  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	753	 85% 12% . .
1	B	753	 80% 15% 5%
1	C	753	 81% 15% .
1	D	753	 84% 13% .
2	E	17	 47% 53%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	17	 47% 53%
2	G	17	 35% 12% 53%
2	H	17	 47% 53%
3	I	2	 100%
3	J	2	 100%
3	K	2	 100%
3	L	2	 50% 50%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 23116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MALTOSE-BINDING PERIPLASMIC PROTEIN, SUPPRESSOR OF FUSED HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	734	5757	3681	967	1090	19	0	0	0
1	B	717	5621	3597	942	1064	18	0	0	0
1	C	725	5687	3640	957	1072	18	0	0	0
1	D	730	5727	3666	962	1081	18	0	0	0

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P0AEX9
A	3	THR	ILE	engineered mutation	UNP P0AEX9
A	360	ALA	GLU	engineered mutation	UNP P0AEX9
A	363	ALA	LYS	engineered mutation	UNP P0AEX9
A	364	ALA	ASP	engineered mutation	UNP P0AEX9
A	368	ASN	ARG	engineered mutation	UNP P0AEX9
A	369	ALA	-	linker	UNP P0AEX9
A	370	ALA	-	linker	UNP P0AEX9
A	371	ALA	-	linker	UNP P0AEX9
A	401	ASP	TRP	engineered mutation	UNP Q9UMX1
A	402	SER	LEU	engineered mutation	UNP Q9UMX1
A	403	PHE	GLY	engineered mutation	UNP Q9UMX1
A	619	PRO	-	linker	UNP Q9UMX1
A	620	SER	-	linker	UNP Q9UMX1
A	621	ARG	-	linker	UNP Q9UMX1
A	622	GLY	-	linker	UNP Q9UMX1
A	625	PRO	-	linker	UNP Q9UMX1
A	718	ALA	PRO	engineered mutation	UNP Q9UMX1
A	?	-	GLU	deletion	UNP Q9UMX1
A	?	-	GLU	deletion	UNP Q9UMX1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	PHE	deletion	UNP Q9UMX1
A	719	ALA	LYS	engineered mutation	UNP Q9UMX1
A	746	VAL	-	expression tag	UNP Q9UMX1
A	747	GLU	-	expression tag	UNP Q9UMX1
A	748	HIS	-	expression tag	UNP Q9UMX1
A	749	HIS	-	expression tag	UNP Q9UMX1
A	750	HIS	-	expression tag	UNP Q9UMX1
A	751	HIS	-	expression tag	UNP Q9UMX1
A	752	HIS	-	expression tag	UNP Q9UMX1
A	753	HIS	-	expression tag	UNP Q9UMX1
B	1	MET	-	expression tag	UNP P0AEX9
B	3	THR	ILE	engineered mutation	UNP P0AEX9
B	360	ALA	GLU	engineered mutation	UNP P0AEX9
B	363	ALA	LYS	engineered mutation	UNP P0AEX9
B	364	ALA	ASP	engineered mutation	UNP P0AEX9
B	368	ASN	ARG	engineered mutation	UNP P0AEX9
B	369	ALA	-	linker	UNP P0AEX9
B	370	ALA	-	linker	UNP P0AEX9
B	371	ALA	-	linker	UNP P0AEX9
B	401	ASP	TRP	engineered mutation	UNP Q9UMX1
B	402	SER	LEU	engineered mutation	UNP Q9UMX1
B	403	PHE	GLY	engineered mutation	UNP Q9UMX1
B	619	PRO	-	linker	UNP Q9UMX1
B	620	SER	-	linker	UNP Q9UMX1
B	621	ARG	-	linker	UNP Q9UMX1
B	622	GLY	-	linker	UNP Q9UMX1
B	625	PRO	-	linker	UNP Q9UMX1
B	718	ALA	PRO	engineered mutation	UNP Q9UMX1
B	?	-	GLU	deletion	UNP Q9UMX1
B	?	-	GLU	deletion	UNP Q9UMX1
B	?	-	PHE	deletion	UNP Q9UMX1
B	719	ALA	LYS	engineered mutation	UNP Q9UMX1
B	746	VAL	-	expression tag	UNP Q9UMX1
B	747	GLU	-	expression tag	UNP Q9UMX1
B	748	HIS	-	expression tag	UNP Q9UMX1
B	749	HIS	-	expression tag	UNP Q9UMX1
B	750	HIS	-	expression tag	UNP Q9UMX1
B	751	HIS	-	expression tag	UNP Q9UMX1
B	752	HIS	-	expression tag	UNP Q9UMX1
B	753	HIS	-	expression tag	UNP Q9UMX1
C	1	MET	-	expression tag	UNP P0AEX9
C	3	THR	ILE	engineered mutation	UNP P0AEX9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	360	ALA	GLU	engineered mutation	UNP P0AEX9
C	363	ALA	LYS	engineered mutation	UNP P0AEX9
C	364	ALA	ASP	engineered mutation	UNP P0AEX9
C	368	ASN	ARG	engineered mutation	UNP P0AEX9
C	369	ALA	-	linker	UNP P0AEX9
C	370	ALA	-	linker	UNP P0AEX9
C	371	ALA	-	linker	UNP P0AEX9
C	401	ASP	TRP	engineered mutation	UNP Q9UMX1
C	402	SER	LEU	engineered mutation	UNP Q9UMX1
C	403	PHE	GLY	engineered mutation	UNP Q9UMX1
C	619	PRO	-	linker	UNP Q9UMX1
C	620	SER	-	linker	UNP Q9UMX1
C	621	ARG	-	linker	UNP Q9UMX1
C	622	GLY	-	linker	UNP Q9UMX1
C	625	PRO	-	linker	UNP Q9UMX1
C	718	ALA	PRO	engineered mutation	UNP Q9UMX1
C	?	-	GLU	deletion	UNP Q9UMX1
C	?	-	GLU	deletion	UNP Q9UMX1
C	?	-	PHE	deletion	UNP Q9UMX1
C	719	ALA	LYS	engineered mutation	UNP Q9UMX1
C	746	VAL	-	expression tag	UNP Q9UMX1
C	747	GLU	-	expression tag	UNP Q9UMX1
C	748	HIS	-	expression tag	UNP Q9UMX1
C	749	HIS	-	expression tag	UNP Q9UMX1
C	750	HIS	-	expression tag	UNP Q9UMX1
C	751	HIS	-	expression tag	UNP Q9UMX1
C	752	HIS	-	expression tag	UNP Q9UMX1
C	753	HIS	-	expression tag	UNP Q9UMX1
D	1	MET	-	expression tag	UNP P0AEX9
D	3	THR	ILE	engineered mutation	UNP P0AEX9
D	360	ALA	GLU	engineered mutation	UNP P0AEX9
D	363	ALA	LYS	engineered mutation	UNP P0AEX9
D	364	ALA	ASP	engineered mutation	UNP P0AEX9
D	368	ASN	ARG	engineered mutation	UNP P0AEX9
D	369	ALA	-	linker	UNP P0AEX9
D	370	ALA	-	linker	UNP P0AEX9
D	371	ALA	-	linker	UNP P0AEX9
D	401	ASP	TRP	engineered mutation	UNP Q9UMX1
D	402	SER	LEU	engineered mutation	UNP Q9UMX1
D	403	PHE	GLY	engineered mutation	UNP Q9UMX1
D	619	PRO	-	linker	UNP Q9UMX1
D	620	SER	-	linker	UNP Q9UMX1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	621	ARG	-	linker	UNP Q9UMX1
D	622	GLY	-	linker	UNP Q9UMX1
D	625	PRO	-	linker	UNP Q9UMX1
D	718	ALA	PRO	engineered mutation	UNP Q9UMX1
D	?	-	GLU	deletion	UNP Q9UMX1
D	?	-	GLU	deletion	UNP Q9UMX1
D	?	-	PHE	deletion	UNP Q9UMX1
D	719	ALA	LYS	engineered mutation	UNP Q9UMX1
D	746	VAL	-	expression tag	UNP Q9UMX1
D	747	GLU	-	expression tag	UNP Q9UMX1
D	748	HIS	-	expression tag	UNP Q9UMX1
D	749	HIS	-	expression tag	UNP Q9UMX1
D	750	HIS	-	expression tag	UNP Q9UMX1
D	751	HIS	-	expression tag	UNP Q9UMX1
D	752	HIS	-	expression tag	UNP Q9UMX1
D	753	HIS	-	expression tag	UNP Q9UMX1

- Molecule 2 is a protein called TRANSCRIPTIONAL ACTIVATOR GLI3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	8	57	35	10	12	0	0	0
2	F	8	57	35	10	12	0	0	0
2	G	8	57	35	10	12	0	0	0
2	H	8	57	35	10	12	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	I	2	23	12	11	0	0	0
3	J	2	23	12	11	0	0	0
3	K	2	23	12	11	0	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	L	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		
4	C	1	Total	Zn	0	0
			1	1		
4	D	1	Total	Zn	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MALTOSE-BINDING PERIPLASMIC PROTEIN, SUPPRESSOR OF FUSED HOMOLOG




- Molecule 1: MALTOSE-BINDING PERIPLASMIC PROTEIN, SUPPRESSOR OF FUSED HOMOLOG



- Molecule 1: MALTOSE-BINDING PERIPLASMIC PROTEIN, SUPPRESSOR OF FUSED HOMOLOG



- Molecule 1: MALTOSE-BINDING PERIPLASMIC PROTEIN, SUPPRESSOR OF FUSED HOMOLOG




- Molecule 2: TRANSCRIPTIONAL ACTIVATOR GLI3




- Molecule 2: TRANSCRIPTIONAL ACTIVATOR GLI3






- Molecule 2: TRANSCRIPTIONAL ACTIVATOR GLI3



- Molecule 2: TRANSCRIPTIONAL ACTIVATOR GLI3



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	116.61Å 136.55Å 116.74Å 90.00° 105.25° 90.00°	Depositor
Resolution (Å)	19.94 – 2.80 19.94 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.94-2.80) 98.5 (19.94-2.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.201 , 0.234 0.209 , 0.242	Depositor DCC
$R_{free}$ test set	1978 reflections (2.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	75.5	Xtrriage
Anisotropy	0.516	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 45.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.044 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	23116	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	96.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/5913	0.83	5/8040 (0.1%)
1	B	0.43	1/5773 (0.0%)	0.84	5/7847 (0.1%)
1	C	0.42	1/5843 (0.0%)	0.86	3/7945 (0.0%)
1	D	0.42	0/5883	0.83	5/8001 (0.1%)
2	E	0.26	0/58	0.72	0/77
2	F	0.26	0/58	0.55	0/77
2	G	0.31	0/58	0.72	0/77
2	H	0.25	0/58	0.64	0/77
All	All	0.42	2/23644 (0.0%)	0.84	18/32141 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	41	PRO	N-CD	5.43	1.55	1.47
1	C	391	PRO	N-CD	5.05	1.54	1.47

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	666	THR	N-CA-C	8.10	119.89	111.14
1	D	337	MET	N-CA-C	6.26	118.19	111.36
1	D	594	ARG	N-CA-C	-6.10	104.74	111.82
1	B	56	ASP	N-CA-C	5.63	119.69	112.26
1	A	109	ILE	CB-CA-C	-5.62	106.11	111.44

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5757	0	5587	77	0
1	B	5621	0	5442	84	0
1	C	5687	0	5522	91	0
1	D	5727	0	5562	62	0
2	E	57	0	49	0	0
2	F	57	0	49	0	0
2	G	57	0	49	1	0
2	H	57	0	49	0	0
3	I	23	0	21	0	0
3	J	23	0	21	0	0
3	K	23	0	21	0	0
3	L	23	0	21	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
All	All	23116	0	22393	311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:413:MET:HE3	1:B:430:TYR:CE2	1.47	1.45
1:C:72:ALA:CB	1:C:105:ILE:HD13	1.45	1.44
1:A:585:PHE:CD1	1:A:591:LEU:HD11	1.72	1.25
1:C:72:ALA:HB1	1:C:105:ILE:HD13	1.19	1.19
1:D:51:VAL:HB	1:D:56:ASP:OD1	1.41	1.18

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/753 (96%)	702 (97%)	22 (3%)	2 (0%)	36	66
1	B	705/753 (94%)	691 (98%)	14 (2%)	0	100	100
1	C	717/753 (95%)	703 (98%)	14 (2%)	0	100	100
1	D	722/753 (96%)	707 (98%)	15 (2%)	0	100	100
2	E	6/17 (35%)	6 (100%)	0	0	100	100
2	F	6/17 (35%)	6 (100%)	0	0	100	100
2	G	6/17 (35%)	6 (100%)	0	0	100	100
2	H	6/17 (35%)	6 (100%)	0	0	100	100
All	All	2894/3080 (94%)	2827 (98%)	65 (2%)	2 (0%)	48	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	679	GLU
1	A	678	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	608/626 (97%)	600 (99%)	8 (1%)	61	86
1	B	593/626 (95%)	589 (99%)	4 (1%)	76	91
1	C	600/626 (96%)	597 (100%)	3 (0%)	81	93

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	605/626 (97%)	600 (99%)	5 (1%)	73	90
2	E	6/12 (50%)	6 (100%)	0	100	100
2	F	6/12 (50%)	6 (100%)	0	100	100
2	G	6/12 (50%)	5 (83%)	1 (17%)	2	8
2	H	6/12 (50%)	6 (100%)	0	100	100
All	All	2430/2552 (95%)	2409 (99%)	21 (1%)	70	89

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	695	TRP
1	D	665	ILE
2	G	333	SER
1	D	715	LEU
1	D	493	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	635	HIS
1	D	553	GLN
1	C	273	ASN
1	D	490	GLN
1	C	206	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GLC	I	1	3	12,12,12	0.54	0	17,17,17	0.52	0
3	GLC	I	2	3	11,11,12	0.64	0	15,15,17	0.66	0
3	GLC	J	1	3	12,12,12	0.55	0	17,17,17	0.62	0
3	GLC	J	2	3	11,11,12	0.63	0	15,15,17	0.58	0
3	GLC	K	1	3	12,12,12	0.54	0	17,17,17	0.67	0
3	GLC	K	2	3	11,11,12	0.61	0	15,15,17	0.74	0
3	GLC	L	1	3	12,12,12	0.56	0	17,17,17	0.91	1 (5%)
3	GLC	L	2	3	11,11,12	0.60	0	15,15,17	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	I	1	3	-	0/2/22/22	0/1/1/1
3	GLC	I	2	3	-	1/2/19/22	0/1/1/1
3	GLC	J	1	3	-	2/2/22/22	0/1/1/1
3	GLC	J	2	3	-	0/2/19/22	0/1/1/1
3	GLC	K	1	3	-	2/2/22/22	0/1/1/1
3	GLC	K	2	3	-	0/2/19/22	0/1/1/1
3	GLC	L	1	3	-	2/2/22/22	0/1/1/1
3	GLC	L	2	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	1	GLC	C1-C2-C3	2.20	114.84	110.36

There are no chirality outliers.


5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	L	1	GLC	C4-C5-C6-O6
3	L	1	GLC	O5-C5-C6-O6
3	J	1	GLC	C4-C5-C6-O6
3	J	1	GLC	O5-C5-C6-O6
3	K	1	GLC	C4-C5-C6-O6


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	734/753 (97%)	-0.35	5 (0%) 84 77	49, 79, 122, 165	0
1	B	717/753 (95%)	-0.20	3 (0%) 88 84	51, 92, 163, 199	0
1	C	725/753 (96%)	-0.06	2 (0%) 90 86	54, 109, 174, 214	0
1	D	730/753 (96%)	-0.28	2 (0%) 90 86	46, 83, 142, 217	0
2	E	8/17 (47%)	0.12	0 100 100	63, 73, 93, 94	0
2	F	8/17 (47%)	0.21	0 100 100	89, 96, 118, 120	0
2	G	8/17 (47%)	0.14	0 100 100	76, 88, 99, 108	0
2	H	8/17 (47%)	-0.28	0 100 100	77, 82, 96, 101	0
All	All	2938/3080 (95%)	-0.22	12 (0%) 88 84	46, 88, 158, 217	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	607	SER	2.5
1	B	614	ALA	2.5
1	A	735	ILE	2.4
1	D	735	ILE	2.4
1	A	3	THR	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GLC	K	1	12/12	0.89	0.08	67,77,85,86	0
3	GLC	L	1	12/12	0.89	0.08	52,59,69,69	0
3	GLC	K	2	11/12	0.92	0.07	61,73,84,94	0
3	GLC	J	1	12/12	0.93	0.09	51,63,80,93	0
3	GLC	I	2	11/12	0.93	0.07	48,56,65,69	0
3	GLC	L	2	11/12	0.94	0.07	47,56,71,73	0
3	GLC	I	1	12/12	0.95	0.06	48,60,73,84	0
3	GLC	J	2	11/12	0.96	0.07	50,57,67,70	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around Chain J:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain K:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ZN	A	910	1/1	0.93	0.06	65,65,65,65	0
4	ZN	B	910	1/1	0.99	0.03	75,75,75,75	0
4	ZN	C	910	1/1	1.00	0.02	57,57,57,57	0
4	ZN	D	910	1/1	1.00	0.01	74,74,74,74	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.