



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:39 PM UTC

PDB ID : 9BL4 / pdb_00009bl4
Title : KIR3DL1*086 in complex with HLA-B*57:03 presenting the AW10 peptide
Authors : Faoro, C.; Rossjohn, J.
Deposited on : 2024-04-29
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	4% 91% 8%
2	B	100	5% 92% 7%
3	C	10	30% 90% 10%
4	G	305	14% 81% 10% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA-B alpha chain (B*5703GB).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	1	5	0
			2226	1397	399	421	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	1	0
			800	511	139	148	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Catenin alpha-1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	0	0	0
			75	49	12	14			

- Molecule 4 is a protein called Killer cell immunoglobulin-like receptor 3DL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	278	Total	C	N	O	S	0	1	0
			2118	1346	378	380	14			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	88	ALA	PRO	conflict	UNP I6LEK9
G	166	PHE	LEU	conflict	UNP I6LEK9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	300	HIS	-	expression tag	UNP I6LEK9
G	301	HIS	-	expression tag	UNP I6LEK9
G	302	HIS	-	expression tag	UNP I6LEK9
G	303	HIS	-	expression tag	UNP I6LEK9
G	304	HIS	-	expression tag	UNP I6LEK9
G	305	HIS	-	expression tag	UNP I6LEK9

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	200	Total	O	0	0
			200	200		
6	B	73	Total	O	0	0
			73	73		
6	C	9	Total	O	0	0
			9	9		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	G	172	Total	O	0	0
			172	172		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA-B alpha chain (B*5703GB)




- Molecule 2: Beta-2-microglobulin



- Molecule 3: Catenin alpha-1 peptide



- Molecule 4: Killer cell immunoglobulin-like receptor 3DL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.94Å 60.54Å 64.90Å 95.37° 96.50° 106.02°	Depositor
Resolution (Å)	44.63 – 1.75 44.63 – 1.75	Depositor EDS
% Data completeness (in resolution range)	93.1 (44.63-1.75) 93.1 (44.63-1.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.75Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.191 , 0.227 (Not available) , 0.239	Depositor DCC
R_{free} test set	3171 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.135	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5715	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/2301	0.51	1/3133 (0.0%)
2	B	0.28	0/826	0.48	0/1124
3	C	0.89	0/77	0.99	0/105
4	G	0.33	0/2185	0.58	0/2972
All	All	0.33	0/5389	0.54	1/7334 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	TYR	CA-CB-CG	-5.23	104.49	113.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2226	0	2084	15	0
2	B	800	0	738	9	0
3	C	75	0	76	1	0
4	G	2118	0	1971	17	0
5	G	42	0	39	2	0
6	A	200	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	73	0	0	0	0
6	C	9	0	0	0	0
6	G	172	0	0	1	0
All	All	5715	0	4908	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:G:93:VAL:HG21	5:G:401:NAG:H82	1.73	0.69
1:A:192:HIS:CE1	2:B:98:ASP:HB3	2.34	0.63
4:G:8:PRO:HB3	4:G:30:TYR:HB2	1.83	0.59
4:G:221:LEU:HD11	4:G:292:VAL:HG21	1.87	0.57
1:A:23:ILE:HD13	2:B:54:LEU:HB3	1.87	0.56
4:G:273:PHE:CE2	4:G:286:PRO:HB3	2.45	0.52
4:G:250:LYS:HB2	4:G:252:ASN:OD1	2.10	0.51
2:B:6:LYS:NZ	2:B:29:GLY:HA3	2.27	0.49
4:G:112:LEU:HD23	4:G:195:VAL:HB	1.95	0.49
1:A:204:TRP:HZ2	2:B:98:ASP:O	1.95	0.49
4:G:133:PHE:HA	4:G:144:SER:O	2.14	0.48
1:A:12:MET:HE2	1:A:12:MET:HB2	1.75	0.47
4:G:235:SER:HB3	4:G:242:GLU:HG2	1.96	0.47
1:A:49:ALA:O	1:A:52:ILE:HG22	2.16	0.46
4:G:7:LYS:HG2	4:G:89:SER:HA	1.98	0.45
4:G:47:ILE:HD12	4:G:48:PRO:HD2	1.99	0.44
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.52	0.44
4:G:6:ASP:N	6:G:516:HOH:O	2.50	0.44
2:B:98:ASP:OD1	2:B:98:ASP:N	2.51	0.44
4:G:236:ARG:HD3	4:G:270:TYR:CZ	2.53	0.44
1:A:238:ASP:O	1:A:239:ARG:HB2	2.17	0.44
4:G:30:TYR:CE2	4:G:31:ARG:HG3	2.52	0.43
4:G:41:LYS:HA	4:G:71:ASN:O	2.18	0.43
4:G:13:TRP:CD2	4:G:14:PRO:HA	2.53	0.43
1:A:215:LEU:HD12	1:A:261:VAL:HG22	2.01	0.43
4:G:254:THR:HG21	5:G:403:NAG:HN2	1.83	0.43
1:A:214:THR:OG1	1:A:262:GLN:HB2	2.20	0.42
1:A:35:ARG:HD3	2:B:53:ASP:OD2	2.20	0.42
1:A:98:MET:HE2	6:A:423:HOH:O	2.19	0.42
1:A:83:ARG:HG3	1:A:84:TYR:N	2.35	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:GLN:HG2	1:A:269:PRO:HB3	2.02	0.41
1:A:159:TYR:HB2	3:C:3:LEU:HD13	2.03	0.40
2:B:39:LEU:HD13	2:B:68:THR:HG22	2.04	0.40
4:G:30:TYR:CZ	4:G:86:SER:HB3	2.57	0.40
1:A:23:ILE:CD1	2:B:54:LEU:HB3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/276 (101%)	272 (98%)	7 (2%)	0	100	100
2	B	99/100 (99%)	99 (100%)	0	0	100	100
3	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
4	G	271/305 (89%)	266 (98%)	5 (2%)	0	100	100
All	All	657/691 (95%)	644 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/230 (98%)	220 (97%)	6 (3%)	39	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	85/95 (90%)	84 (99%)	1 (1%)	63	49
3	C	8/8 (100%)	8 (100%)	0	100	100
4	G	222/261 (85%)	219 (99%)	3 (1%)	59	44
All	All	541/594 (91%)	531 (98%)	10 (2%)	50	33

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	MET
1	A	14	ARG
1	A	54	GLN
1	A	83	ARG
1	A	111	ARG
1	A	170	ARG
2	B	98	ASP
4	G	102	LYS
4	G	142	ASP
4	G	154	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN
1	A	114	ASN
2	B	89	GLN
4	G	20	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	G	403	4	14,14,15	0.23	0	17,19,21	0.41	0
5	NAG	G	401	4	14,14,15	0.38	0	17,19,21	0.69	1 (5%)
5	NAG	G	402	4	14,14,15	0.20	0	17,19,21	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	403	4	-	2/6/23/26	0/1/1/1
5	NAG	G	401	4	-	0/6/23/26	0/1/1/1
5	NAG	G	402	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	401	NAG	C1-O5-C5	2.26	115.22	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	402	NAG	O5-C5-C6-O6
5	G	403	NAG	O5-C5-C6-O6
5	G	403	NAG	C4-C5-C6-O6
5	G	402	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	403	NAG	1	0
5	G	401	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	0.38	11 (3%) 42 48	9, 26, 51, 74	6 (2%)
2	B	100/100 (100%)	0.48	5 (5%) 34 39	15, 29, 62, 78	1 (1%)
3	C	10/10 (100%)	1.37	3 (30%) 1 1	23, 32, 40, 41	0
4	G	278/305 (91%)	0.82	42 (15%) 5 6	11, 30, 63, 99	1 (0%)
All	All	664/691 (96%)	0.60	61 (9%) 14 17	9, 28, 58, 99	8 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	263	PRO	5.7
4	G	219	VAL	5.2
1	A	194	ILE	5.1
4	G	49	ILE	5.1
4	G	35	ASN	4.3
2	B	73	THR	4.2
4	G	265	THR	3.9
4	G	50	PHE	3.9
4	G	218	SER	3.8
4	G	13	TRP	3.7
4	G	80	HIS	3.6
4	G	85	TRP	3.5
4	G	47	ILE	3.5
1	A	225	THR	3.5
4	G	294	VAL	3.4
4	G	252	ASN	3.4
4	G	239	GLY	3.4
2	B	99	MET	3.2
4	G	251	VAL	3.2
1	A	1	GLY	3.2
3	C	6	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	G	48	PRO	3.0
4	G	240	ALA	3.0
1	A	221	GLY	3.0
4	G	46	HIS	2.9
1	A	42[A]	SER	2.9
1	A	41	ALA	2.8
4	G	6	ASP	2.8
4	G	126	ASP	2.8
4	G	79	PRO	2.8
3	C	5	LEU	2.8
4	G	44	ARG	2.8
2	B	0	MET	2.8
4	G	208	GLN	2.7
4	G	57	GLU	2.7
2	B	1	ILE	2.7
4	G	238	GLY	2.6
1	A	223	ASP	2.5
4	G	32	HIS	2.5
1	A	192	HIS	2.4
4	G	220	THR	2.4
4	G	41	LYS	2.4
4	G	82	PRO	2.3
4	G	45	ILE	2.3
3	C	7	ALA	2.3
1	A	58	GLU	2.2
2	B	97	ARG	2.2
4	G	56	GLN	2.2
4	G	34	PHE	2.2
4	G	151	HIS	2.2
4	G	216	GLY	2.2
4	G	101	ARG	2.2
4	G	278	HIS	2.2
4	G	231	MET	2.1
1	A	274	TRP	2.1
4	G	242	GLU	2.1
4	G	264	ALA	2.1
4	G	217	GLU	2.1
4	G	55	PHE	2.1
4	G	52	GLY	2.0
1	A	227	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	G	403	14/15	0.58	0.21	84,92,99,105	0
5	NAG	G	402	14/15	0.80	0.15	55,63,72,74	0
5	NAG	G	401	14/15	0.83	0.12	44,54,57,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.