



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:37 AM UTC

PDB ID : 4BMK / pdb_00004bmk
Title : Serine Palmitoyltransferase K265A from *S. paucimobilis* with bound PLP-Myriocin Aldimine
Authors : Wadsworth, J.M.; Clarke, D.J.; McMahon, S.A.; Beattie, A.E.; Lowther, J.; Dunn, T.M.; Naismith, J.H.; Campopiano, D.J.
Deposited on : 2013-05-09
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	442	5% 85% 6% 10%
1	B	442	10% 81% 6% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1424	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6602 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE PALMITOYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	399	Total	C	N	O	S	25	8	1
			3040	1927	524	565	24			
1	B	386	Total	C	N	O	S	21	3	0
			2927	1849	505	550	23			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	GLY	-	expression tag	UNP Q93UV0
A	-19	SER	-	expression tag	UNP Q93UV0
A	-18	SER	-	expression tag	UNP Q93UV0
A	-17	HIS	-	expression tag	UNP Q93UV0
A	-16	HIS	-	expression tag	UNP Q93UV0
A	-15	HIS	-	expression tag	UNP Q93UV0
A	-14	HIS	-	expression tag	UNP Q93UV0
A	-13	HIS	-	expression tag	UNP Q93UV0
A	-12	HIS	-	expression tag	UNP Q93UV0
A	-11	SER	-	expression tag	UNP Q93UV0
A	-10	SER	-	expression tag	UNP Q93UV0
A	-9	GLY	-	expression tag	UNP Q93UV0
A	-8	LEU	-	expression tag	UNP Q93UV0
A	-7	VAL	-	expression tag	UNP Q93UV0
A	-6	PRO	-	expression tag	UNP Q93UV0
A	-5	ARG	-	expression tag	UNP Q93UV0
A	-4	GLY	-	expression tag	UNP Q93UV0
A	-3	SER	-	expression tag	UNP Q93UV0
A	-2	HIS	-	expression tag	UNP Q93UV0
A	-1	MET	-	expression tag	UNP Q93UV0
A	0	ALA	-	expression tag	UNP Q93UV0
A	1	SER	-	expression tag	UNP Q93UV0
A	421	LEU	-	expression tag	UNP Q93UV0
A	265	ALA	LYS	engineered mutation	UNP Q93UV0
B	-20	GLY	-	expression tag	UNP Q93UV0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	SER	-	expression tag	UNP Q93UV0
B	-18	SER	-	expression tag	UNP Q93UV0
B	-17	HIS	-	expression tag	UNP Q93UV0
B	-16	HIS	-	expression tag	UNP Q93UV0
B	-15	HIS	-	expression tag	UNP Q93UV0
B	-14	HIS	-	expression tag	UNP Q93UV0
B	-13	HIS	-	expression tag	UNP Q93UV0
B	-12	HIS	-	expression tag	UNP Q93UV0
B	-11	SER	-	expression tag	UNP Q93UV0
B	-10	SER	-	expression tag	UNP Q93UV0
B	-9	GLY	-	expression tag	UNP Q93UV0
B	-8	LEU	-	expression tag	UNP Q93UV0
B	-7	VAL	-	expression tag	UNP Q93UV0
B	-6	PRO	-	expression tag	UNP Q93UV0
B	-5	ARG	-	expression tag	UNP Q93UV0
B	-4	GLY	-	expression tag	UNP Q93UV0
B	-3	SER	-	expression tag	UNP Q93UV0
B	-2	HIS	-	expression tag	UNP Q93UV0
B	-1	MET	-	expression tag	UNP Q93UV0
B	0	ALA	-	expression tag	UNP Q93UV0
B	1	SER	-	expression tag	UNP Q93UV0
B	421	LEU	-	expression tag	UNP Q93UV0
B	265	ALA	LYS	engineered mutation	UNP Q93UV0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is Decarboxylated Myriocin (CCD ID: MYB) (formula: $C_{20}H_{39}NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			13	9	1	3		
3	B	1	Total	C	N	O	0	0
			13	9	1	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	313	Total	O	0	0
			313	313		
5	B	254	Total	O	0	0
			254	254		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.91Å 98.71Å 133.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.14 – 1.62 56.14 – 1.62	Depositor EDS
% Data completeness (in resolution range)	98.7 (56.14-1.62) 98.7 (56.14-1.62)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.189 , 0.211 0.200 , 0.221	Depositor DCC
R_{free} test set	5148 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6602	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, GOL, MYB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	3/3120 (0.1%)	0.85	3/4217 (0.1%)
1	B	0.69	1/2987 (0.0%)	0.82	2/4031 (0.0%)
All	All	0.79	4/6107 (0.1%)	0.83	5/8248 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	207	MET	CB-CG	-28.74	0.66	1.52
1	A	175	ARG	CG-CD	11.37	1.86	1.52
1	B	186	LYS	CB-CG	-5.47	1.36	1.52
1	A	186	LYS	CB-CG	-5.46	1.36	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	MET	CA-CB-CG	11.76	137.62	114.10
1	A	175	ARG	CB-CG-CD	-8.78	91.12	111.30
1	B	207	MET	CA-CB-CG	7.62	129.34	114.10
1	A	186	LYS	CA-CB-CG	6.26	126.63	114.10
1	B	186	LYS	CA-CB-CG	5.82	125.73	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3040	0	3071	16	0
1	B	2927	0	2931	27	0
2	A	15	0	6	1	0
2	B	15	0	6	1	0
3	A	13	0	14	1	0
3	B	13	0	14	0	0
4	A	12	0	16	4	0
5	A	313	0	0	0	0
5	B	254	0	0	9	0
All	All	6602	0	6058	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:353:MET:HG2	5:B:2238:HOH:O	1.66	0.96
1:B:310:ARG:HD2	5:B:2214:HOH:O	1.77	0.84
1:B:388:LEU:HD22	5:B:2238:HOH:O	1.84	0.77
1:B:363:MET:HE1	1:B:412:ALA:C	2.12	0.74
1:B:253:GLU:OE2	5:B:2192:HOH:O	2.12	0.66
1:B:208:LEU:HB2	5:B:2165:HOH:O	1.96	0.65
1:B:73:TYR:OH	1:B:390:ARG:NH2	2.29	0.64
1:B:363:MET:CE	1:B:412:ALA:HB1	2.27	0.64
1:A:354:LEU:HD13	1:A:360:ALA:HA	1.80	0.62
1:B:354:LEU:HD13	1:B:360:ALA:HA	1.81	0.60
1:B:363:MET:CE	1:B:412:ALA:CB	2.86	0.53
1:B:363:MET:HE1	1:B:412:ALA:CB	2.38	0.53
1:B:393:ILE:HD12	1:B:393:ILE:C	2.34	0.53
1:A:260[B]:VAL:HG22	1:A:274:VAL:HG22	1.91	0.52
1:B:207:MET:HB3	5:B:2164:HOH:O	2.08	0.52
1:A:393:ILE:C	1:A:393:ILE:HD12	2.35	0.51
1:B:339:ARG:NH1	1:B:353:MET:SD	2.84	0.51
1:A:339:ARG:NH1	1:A:353[A]:MET:SD	2.85	0.50
1:A:319:ARG:HE	4:A:1424:GOL:C3	2.24	0.50
1:B:363:MET:HE1	1:B:412:ALA:HB1	1.94	0.49
1:B:322:LEU:C	1:B:322:LEU:HD23	2.38	0.48
1:B:390:ARG:CD	5:B:2244:HOH:O	2.61	0.48
1:A:404:THR:CG2	1:A:408:MET:HE3	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:VAL:HG12	1:A:44:THR:O	2.15	0.47
1:A:269[B]:THR:HG23	1:A:270:VAL:H	1.80	0.47
1:A:322:LEU:C	1:A:322:LEU:HD23	2.39	0.47
1:B:269:THR:HG23	1:B:270:VAL:H	1.80	0.46
1:B:388:LEU:CD2	5:B:2238:HOH:O	2.54	0.45
1:B:363:MET:HA	1:B:363:MET:HE3	1.98	0.45
1:B:269:THR:HG23	1:B:270:VAL:N	2.32	0.44
1:A:363:MET:HB3	1:A:389:LEU:HD11	2.00	0.44
2:A:1421:PLP:O3	3:A:1422:MYB:N2	2.50	0.43
1:B:204:VAL:HG22	1:B:211:ILE:CD1	2.49	0.43
1:B:353:MET:HE3	5:B:2238:HOH:O	2.17	0.43
1:B:231:ASP:OD2	2:B:1421:PLP:N1	2.52	0.42
1:A:319:ARG:HE	4:A:1424:GOL:H32	1.85	0.42
1:A:202:GLU:HB2	1:A:205:TYR:CE1	2.56	0.41
1:A:319:ARG:HE	4:A:1424:GOL:H31	1.85	0.41
1:A:269[B]:THR:HG23	1:A:270:VAL:N	2.35	0.41
1:A:320:GLU:OE2	4:A:1424:GOL:O3	2.33	0.41
1:A:204:VAL:HG22	1:A:211:ILE:HG22	2.03	0.41
1:B:204:VAL:HG22	1:B:211:ILE:HD12	2.03	0.40
1:B:269:THR:HG21	1:B:305:ALA:HB2	2.02	0.40
1:B:205:TYR:HD2	1:B:210:ASP:OD1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/442 (92%)	394 (97%)	11 (3%)	0	100	100
1	B	385/442 (87%)	373 (97%)	12 (3%)	0	100	100
All	All	790/884 (89%)	767 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/345 (93%)	321 (100%)	0	100	100
1	B	307/345 (89%)	305 (99%)	2 (1%)	76	62
All	All	628/690 (91%)	626 (100%)	2 (0%)	86	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	38	LEU
1	B	194	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	357	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	B	1421	3	15,15,16	1.04	1 (6%)	21,22,23	2.49	9 (42%)
4	GOL	A	1423	-	5,5,5	0.33	0	5,5,5	0.64	0
4	GOL	A	1424	-	5,5,5	0.20	0	5,5,5	0.32	0
3	MYB	A	1422	2	12,12,24	1.26	1 (8%)	9,14,27	2.00	3 (33%)
2	PLP	A	1421	3	15,15,16	0.82	0	21,22,23	2.48	8 (38%)
3	MYB	B	1422	2	12,12,24	1.30	1 (8%)	9,14,27	1.65	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	1421	3	-	3/6/6/8	0/1/1/1
4	GOL	A	1423	-	-	0/4/4/4	-
4	GOL	A	1424	-	-	0/4/4/4	-
3	MYB	A	1422	2	-	6/15/15/28	-
2	PLP	A	1421	3	-	5/6/6/8	0/1/1/1
3	MYB	B	1422	2	-	9/15/15/28	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1422	MYB	C11-C12	3.96	1.54	1.31
3	A	1422	MYB	C11-C12	3.85	1.53	1.31
2	B	1421	PLP	C5-C4	2.62	1.43	1.40

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1421	PLP	O4P-C5A-C5	5.80	120.23	109.36
2	B	1421	PLP	O2P-P-O4P	-5.75	91.68	106.67
2	B	1421	PLP	C5A-C5-C6	-4.72	111.67	119.36
3	A	1422	MYB	C10-C11-C12	-4.62	109.23	126.21
2	B	1421	PLP	O4P-C5A-C5	4.40	117.60	109.36
2	A	1421	PLP	O2P-P-O4P	-4.01	96.21	106.67
2	A	1421	PLP	O4P-P-O1P	-4.00	95.62	106.44
2	A	1421	PLP	O3P-P-O4P	3.72	116.38	106.67
2	B	1421	PLP	C4A-C4-C5	3.66	124.72	120.94
2	A	1421	PLP	C4A-C4-C5	3.66	124.72	120.94
3	B	1422	MYB	C10-C11-C12	-3.53	113.25	126.21
2	A	1421	PLP	C5A-C5-C6	-2.98	114.50	119.36
2	B	1421	PLP	C5A-C5-C4	2.81	128.18	122.64
2	A	1421	PLP	C6-C5-C4	2.57	120.20	118.10
2	A	1421	PLP	O3P-P-O2P	2.54	117.31	107.80
2	B	1421	PLP	C6-C5-C4	2.49	120.14	118.10
2	B	1421	PLP	O3P-P-O4P	2.42	112.98	106.67
2	B	1421	PLP	C3-C4-C5	-2.18	115.97	118.59
3	A	1422	MYB	CBN-C10-C11	2.17	116.28	112.88
3	A	1422	MYB	C13-C12-C11	-2.14	113.55	126.42
2	B	1421	PLP	O3P-P-O2P	2.02	115.39	107.80

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1421	PLP	C4-C5-C5A-O4P
2	A	1421	PLP	C6-C5-C5A-O4P
2	A	1421	PLP	C5A-O4P-P-O1P
2	A	1421	PLP	C5A-O4P-P-O2P
2	B	1421	PLP	C4-C5-C5A-O4P
2	B	1421	PLP	C6-C5-C5A-O4P
3	A	1422	MYB	N2-C6-C8-OAJ
3	A	1422	MYB	N2-C6-C8-CBN
3	A	1422	MYB	C7-C6-C8-OAJ
3	A	1422	MYB	C7-C6-C8-CBN
3	B	1422	MYB	N2-C6-C7-O4
3	B	1422	MYB	N2-C6-C8-OAJ
3	B	1422	MYB	N2-C6-C8-CBN
3	B	1422	MYB	C7-C6-C8-OAJ
3	B	1422	MYB	C7-C6-C8-CBN
3	B	1422	MYB	C10-C11-C12-C13
3	B	1422	MYB	C11-C12-C13-C14

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	1422	MYB	C8-C6-C7-O4
2	A	1421	PLP	C5A-O4P-P-O3P
2	B	1421	PLP	C5A-O4P-P-O1P
3	B	1422	MYB	CBN-C10-C11-C12
3	A	1422	MYB	N2-C6-C7-O4
3	A	1422	MYB	C11-C12-C13-C14

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1421	PLP	1	0
4	A	1424	GOL	4	0
3	A	1422	MYB	1	0
2	A	1421	PLP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/442 (90%)	0.18	24 (6%) 27 29	6, 16, 30, 52	14 (3%)
1	B	386/442 (87%)	0.41	46 (11%) 9 9	7, 17, 48, 70	8 (2%)
All	All	785/884 (88%)	0.29	70 (8%) 15 16	6, 16, 40, 70	22 (2%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	38	LEU	6.9
1	B	73	TYR	5.5
1	A	381	ALA	5.0
1	A	383	PRO	4.9
1	B	417	GLY	4.9
1	B	42	GLY	4.8
1	A	387	PHE	4.8
1	B	354	LEU	4.7
1	B	356	ASP	4.6
1	B	43	VAL	4.6
1	A	384	ALA	4.6
1	A	379	PRO	4.6
1	B	39	LEU	4.5
1	A	382	THR	4.5
1	A	380	PRO	4.4
1	A	385	GLY	4.3
1	B	41	SER	4.3
1	B	205	TYR	4.1
1	B	415	ALA	4.0
1	A	378	ARG	4.0
1	A	73	TYR	3.8
1	B	355	GLU	3.8
1	B	388	LEU	3.7
1	B	208	LEU	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	386	THR	3.6
1	B	357	GLN	3.5
1	B	391	CYS	3.4
1	B	360	ALA	3.4
1	B	353	MET	3.4
1	B	416	VAL	3.3
1	B	358	GLU	3.2
1	B	40	ASP	3.1
1	B	377	ALA	2.9
1	B	35	ARG	2.9
1	B	359	GLN	2.9
1	B	389	LEU	2.9
1	B	335	ALA	2.8
1	B	336	MET	2.8
1	B	352	VAL	2.7
1	B	37	LYS	2.6
1	B	411	ALA	2.6
1	A	390	ARG	2.6
1	A	47	PHE	2.5
1	A	316	HIS	2.5
1	A	357	GLN	2.4
1	A	355	GLU	2.4
1	B	390	ARG	2.4
1	B	207	MET	2.4
1	A	377	ALA	2.4
1	A	414	ARG	2.4
1	A	358	GLU	2.3
1	B	361	ALA	2.3
1	B	47	PHE	2.3
1	B	338	PHE	2.3
1	A	207	MET	2.3
1	B	412	ALA	2.2
1	B	362	MET	2.2
1	A	418	VAL	2.2
1	B	339	ARG	2.2
1	B	179	ASN	2.2
1	B	22	ARG	2.2
1	B	337	GLY	2.2
1	B	413	GLY	2.2
1	B	206	SER	2.1
1	A	34	GLU	2.1
1	B	36	GLN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	356	ASP	2.0
1	B	342	THR	2.0
1	B	182	GLU	2.0
1	A	416	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MYB	A	1422	13/25	0.66	0.22	40,42,45,45	0
4	GOL	A	1424	6/6	0.71	0.18	38,40,43,45	0
3	MYB	B	1422	13/25	0.77	0.16	32,35,37,37	0
4	GOL	A	1423	6/6	0.94	0.11	21,23,24,24	0
2	PLP	A	1421	15/16	0.94	0.11	16,30,33,36	0
2	PLP	B	1421	15/16	0.95	0.10	16,27,30,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MYB A 1422:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around MYB B 1422:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.