



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:31 PM EDT

PDB ID : 3BND / pdb_00003bnd
Title : Lipoygenase-1 (Soybean), I553V Mutant
Authors : Tomchick, D.R.
Deposited on : 2007-12-14
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	839	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Seed lipoxygenase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	817	Total	C	N	O	S	0	14	0
			6617	4236	1118	1243	20			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	GLU	SER	engineered mutation	UNP P08170
A	553	VAL	ILE	engineered mutation	UNP P08170

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	641	Total	O	0	0
			641	641		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Seed lipoxygenase-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	94.01Å 93.00Å 49.35Å 90.00° 90.61° 90.00°	Depositor
Resolution (Å)	27.37 – 1.60 27.37 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.37-1.60) 99.5 (27.37-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.168 , 0.200 0.171 , 0.204	Depositor DCC
R_{free} test set	3349 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.004 for k,h,-l 0.006 for -k,-h,-l 0.023 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7259	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	3/6782 (0.0%)	0.91	6/9216 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	629	VAL	CA-CB	6.81	1.57	1.54
1	A	90	SER	CB-OG	5.46	1.53	1.42
1	A	341	MET	SD-CE	-5.37	1.66	1.79

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	308	ILE	CA-C-N	-9.54	109.50	121.04
1	A	308	ILE	C-N-CA	-9.54	109.50	121.04
1	A	211	THR	N-CA-C	-5.45	105.42	111.36
1	A	279	THR	CA-C-N	-5.41	114.21	119.78
1	A	279	THR	C-N-CA	-5.41	114.21	119.78
1	A	663	LYS	N-CA-C	5.19	116.37	109.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6617	0	6552	33	0
2	A	1	0	0	0	0
3	A	641	0	0	5	0
All	All	7259	0	6552	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:LEU:HD23	1:A:750[B]:VAL:HG22	1.72	0.71
1:A:270:PHE:HD2	1:A:746[A]:ILE:CD1	2.07	0.66
1:A:88:ASP:OD1	1:A:88:ASP:N	2.24	0.66
1:A:552:ILE:HG21	1:A:750[B]:VAL:HG21	1.80	0.64
1:A:301:ARG:HG3	1:A:323:HIS:NE2	2.14	0.62
1:A:308:ILE:HD11	1:A:321:GLY:HA2	1.86	0.57
1:A:262:LEU:HD21	1:A:750[B]:VAL:HG23	1.86	0.57
1:A:270:PHE:HD2	1:A:746[A]:ILE:HD11	1.69	0.56
1:A:7:LYS:HD3	1:A:84:HIS:HB3	1.88	0.55
1:A:262:LEU:CD2	1:A:750[B]:VAL:HG22	2.36	0.54
1:A:270:PHE:HD2	1:A:746[A]:ILE:HD13	1.73	0.53
1:A:196:ASP:OD2	1:A:226[B]:VAL:HG22	2.10	0.52
1:A:354[B]:VAL:CG2	1:A:700:TYR:HB2	2.41	0.51
1:A:7:LYS:CD	1:A:84:HIS:HB3	2.41	0.51
1:A:262:LEU:CD2	1:A:750[B]:VAL:CG2	2.90	0.49
1:A:15[A]:MET:HE2	1:A:76:ALA:O	2.14	0.48
1:A:318:ARG:NH2	3:A:1345:HOH:O	2.47	0.48
1:A:497:MET:SD	1:A:567:SER:HA	2.54	0.47
1:A:262:LEU:HD21	1:A:750[B]:VAL:CG2	2.44	0.47
1:A:649:TRP:HB2	1:A:669:LEU:HD22	1.97	0.47
1:A:76:ALA:HB1	1:A:249:LEU:HD11	1.96	0.46
1:A:41:SER:HB3	1:A:63:PHE:CE2	2.51	0.46
1:A:270:PHE:CD2	1:A:746[A]:ILE:HD13	2.50	0.46
1:A:606:GLU:HG2	3:A:1569:HOH:O	2.18	0.44
1:A:15[A]:MET:HE2	1:A:78:GLU:HG3	2.00	0.43
1:A:750[A]:VAL:O	1:A:754:LEU:HG	2.18	0.43
1:A:340:TRP:CH2	1:A:341:MET:HG2	2.55	0.41
1:A:495:GLN:NE2	1:A:697:GLN:OE1	2.47	0.41
1:A:509:PRO:HB3	1:A:656:GLY:HA3	2.03	0.41
1:A:308:ILE:HG23	3:A:1564:HOH:O	2.20	0.41
1:A:91:MET:HA	3:A:1235:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:530:PRO:HG2	1:A:767:ARG:CZ	2.52	0.40
1:A:713:ARG:CZ	3:A:1437:HOH:O	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	825/839 (98%)	793 (96%)	31 (4%)	1 (0%)	48 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	726/730 (100%)	720 (99%)	6 (1%)	73 59

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	70	SER
1	A	88	ASP
1	A	115	GLU
1	A	592	ILE
1	A	608	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	502	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	817/839 (97%)	0.56	61 (7%)	20 20	9, 22, 36, 51	14 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	89	GLY	6.1
1	A	121	GLY	5.1
1	A	77	GLY	4.6
1	A	116	ALA	4.5
1	A	72	PRO	4.2
1	A	71	LEU	4.2
1	A	457	GLY	4.2
1	A	76	ALA	4.0
1	A	289	VAL	4.0
1	A	90	SER	3.8
1	A	246	LEU	3.8
1	A	285	SER	3.8
1	A	293	TYR	3.6
1	A	75	GLY	3.5
1	A	286	PHE	3.5
1	A	88	ASP	3.3
1	A	295	GLY	3.1
1	A	36	LEU	3.1
1	A	69	THR	3.1
1	A	73	THR	3.0
1	A	292	LEU	3.0
1	A	308	ILE	3.0
1	A	302	ASP	3.0
1	A	284	HIS	2.9
1	A	74	LEU	2.9
1	A	54	GLY	2.9
1	A	251	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	301	ARG	2.8
1	A	34	ALA	2.7
1	A	70	SER	2.7
1	A	454	HIS	2.6
1	A	296	GLY	2.6
1	A	458	ASP	2.6
1	A	662	ASP	2.6
1	A	287	GLN	2.5
1	A	13	VAL	2.5
1	A	290	HIS	2.5
1	A	160	GLU	2.5
1	A	272	SER	2.5
1	A	305	SER	2.5
1	A	213	PRO	2.5
1	A	226[A]	VAL	2.5
1	A	31	ASN	2.4
1	A	456	ALA	2.4
1	A	10	GLY	2.3
1	A	643	SER	2.3
1	A	93	ILE	2.3
1	A	63	PHE	2.3
1	A	130	TRP	2.3
1	A	262	LEU	2.2
1	A	91	MET	2.2
1	A	595	PRO	2.2
1	A	18	ASN	2.2
1	A	791	GLU	2.1
1	A	87	TRP	2.1
1	A	291	ASP	2.1
1	A	303	VAL	2.1
1	A	386	ASP	2.1
1	A	8	ILE	2.1
1	A	212	PHE	2.1
1	A	11	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FE	A	840	1/1	0.99	0.11	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.