



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:14 AM UTC

PDB ID : 6BN9 / pdb_00006bn9
Title : Crystal structure of DDB1-CRBN-BRD4(BD1) complex bound to dBET70 PROTAC
Authors : Nowak, R.P.; DeAngelo, S.L.; Buckley, D.; Bradner, J.E.; Fischer, E.S.
Deposited on : 2017-11-16
Resolution : 4.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 4.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (4.84-3.90)
Clashscore	190562	1132 (4.84-3.90)
Ramachandran outliers	187476	1024 (4.84-3.90)
Sidechain outliers	187428	1008 (4.84-3.90)
RSRZ outliers	180081	1081 (4.84-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	864	
2	B	463	
3	C	127	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA damage-binding protein 1, DNA damage-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	808	Total	C	N	O	S	0	0	0
			6249	3962	1043	1209	35			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-27	MET	-	initiating methionine	UNP Q16531
A	-26	GLY	-	expression tag	UNP Q16531
A	-25	SER	-	expression tag	UNP Q16531
A	-24	SER	-	expression tag	UNP Q16531
A	-23	HIS	-	expression tag	UNP Q16531
A	-22	HIS	-	expression tag	UNP Q16531
A	-21	HIS	-	expression tag	UNP Q16531
A	-20	HIS	-	expression tag	UNP Q16531
A	-19	HIS	-	expression tag	UNP Q16531
A	-18	HIS	-	expression tag	UNP Q16531
A	-17	SER	-	expression tag	UNP Q16531
A	-16	ALA	-	expression tag	UNP Q16531
A	-15	ALA	-	expression tag	UNP Q16531
A	-14	HIS	-	expression tag	UNP Q16531
A	-13	ILE	-	expression tag	UNP Q16531
A	-12	VAL	-	expression tag	UNP Q16531
A	-11	MET	-	expression tag	UNP Q16531
A	-10	VAL	-	expression tag	UNP Q16531
A	-9	ASP	-	expression tag	UNP Q16531
A	-8	ALA	-	expression tag	UNP Q16531
A	-7	TYR	-	expression tag	UNP Q16531
A	-6	LYS	-	expression tag	UNP Q16531
A	-5	PRO	-	expression tag	UNP Q16531
A	-4	THR	-	expression tag	UNP Q16531
A	-3	LYS	-	expression tag	UNP Q16531
A	-2	GLY	-	expression tag	UNP Q16531

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q16531
A	0	ARG	-	expression tag	UNP Q16531
A	700	GLY	-	linker	UNP Q16531
A	701	ASN	-	linker	UNP Q16531
A	702	GLY	-	linker	UNP Q16531
A	703	ASN	-	linker	UNP Q16531
A	704	SER	-	linker	UNP Q16531
A	705	GLY	-	linker	UNP Q16531

- Molecule 2 is a protein called Protein cereblon.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	375	Total	C	N	O	S	0	0	0
			3005	1919	512	550	24			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MET	-	initiating methionine	UNP Q96SW2
B	-19	GLY	-	expression tag	UNP Q96SW2
B	-18	SER	-	expression tag	UNP Q96SW2
B	-17	SER	-	expression tag	UNP Q96SW2
B	-16	HIS	-	expression tag	UNP Q96SW2
B	-15	HIS	-	expression tag	UNP Q96SW2
B	-14	HIS	-	expression tag	UNP Q96SW2
B	-13	HIS	-	expression tag	UNP Q96SW2
B	-12	HIS	-	expression tag	UNP Q96SW2
B	-11	HIS	-	expression tag	UNP Q96SW2
B	-10	SER	-	expression tag	UNP Q96SW2
B	-9	ALA	-	expression tag	UNP Q96SW2
B	-8	VAL	-	expression tag	UNP Q96SW2
B	-7	ASP	-	expression tag	UNP Q96SW2
B	-6	GLU	-	expression tag	UNP Q96SW2
B	-5	ASN	-	expression tag	UNP Q96SW2
B	-4	LEU	-	expression tag	UNP Q96SW2
B	-3	TYR	-	expression tag	UNP Q96SW2
B	-2	PHE	-	expression tag	UNP Q96SW2
B	-1	GLN	-	expression tag	UNP Q96SW2
B	0	GLY	-	expression tag	UNP Q96SW2
B	1	GLY	-	expression tag	UNP Q96SW2

- Molecule 3 is a protein called Bromodomain-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	127	Total	C	N	O	S	0	0	0
			1059	685	173	194	7			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	43	MET	THR	engineered mutation	UNP O60885


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA damage-binding protein 1, DNA damage-binding protein 1



- Molecule 2: Protein cereblon



- Molecule 3: Bromodomain-containing protein 4



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	117.60Å 117.60Å 597.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	100.40 – 4.38 100.40 – 4.38	Depositor EDS
% Data completeness (in resolution range)	99.5 (100.40-4.38) 99.5 (100.40-4.38)	Depositor EDS
R_{merge}	0.35	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.41 (at 4.47Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.275 , 0.301 0.275 , 0.294	Depositor DCC
R_{free} test set	851 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	206.8	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 679.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	10314	wwPDB-VP
Average B, all atoms (Å ²)	278.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.10	0/6362	0.29	0/8624
2	B	0.11	0/3077	0.30	0/4177
3	C	0.10	0/1089	0.25	0/1481
All	All	0.10	0/10528	0.29	0/14282

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6249	0	6105	46	0
2	B	3005	0	2985	27	0
3	C	1059	0	1047	6	0
4	B	1	0	0	0	0
All	All	10314	0	10137	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ALA:HB1	1:A:327:ARG:HG3	1.54	0.90
1:A:327:ARG:NH2	2:B:200:GLU:OE1	2.29	0.66
1:A:980:ASP:HB3	1:A:983:ALA:HB2	1.79	0.64
1:A:1005:ASN:ND2	2:B:237:LEU:O	2.34	0.60
1:A:1109:VAL:HG12	1:A:1129:LEU:HD22	1.85	0.59
1:A:143:ILE:HG12	1:A:154:ALA:HB2	1.85	0.58
3:C:103:THR:O	3:C:135:ASN:ND2	2.37	0.57
1:A:327:ARG:HD2	2:B:199:LEU:CD2	2.35	0.56
2:B:168:LEU:HD11	2:B:183:GLN:HB2	1.88	0.55
1:A:983:ALA:HB3	1:A:989:ARG:HG2	1.88	0.54
2:B:197:VAL:HG12	2:B:237:LEU:HD12	1.88	0.54
2:B:152:ILE:HB	3:C:152:ALA:HB2	1.90	0.54
2:B:355:TYR:HD1	2:B:356:VAL:H	1.55	0.53
1:A:5:TYR:HB2	1:A:1043:LEU:HD11	1.90	0.53
1:A:195:VAL:HG22	1:A:202:PHE:HE1	1.73	0.52
1:A:812:TYR:CZ	2:B:241:PRO:HB3	2.45	0.52
1:A:275:ASP:OD1	1:A:279:ARG:N	2.42	0.52
1:A:952:ASN:ND2	1:A:969:GLU:OE1	2.37	0.51
1:A:310:ILE:HG21	1:A:328:LEU:HD12	1.93	0.50
1:A:118:THR:OG1	1:A:134:ARG:NH2	2.44	0.50
1:A:785:GLU:HB2	2:B:229:LYS:HZ1	1.77	0.49
1:A:835:MET:HB2	1:A:845:GLN:HB3	1.94	0.49
1:A:1030:PHE:CZ	1:A:1038:GLY:HA3	2.48	0.49
1:A:59:GLY:HA2	1:A:1073:TRP:CZ3	2.47	0.49
1:A:812:TYR:OH	2:B:241:PRO:HB3	2.14	0.48
1:A:928:ARG:HB3	1:A:950:ASN:O	2.14	0.48
2:B:257:ILE:HG23	2:B:315:MET:HE1	1.97	0.47
1:A:358:PRO:HD2	1:A:380:GLY:HA2	1.97	0.46
1:A:909:ILE:HD12	1:A:925:ASP:HB2	1.96	0.46
2:B:346:MET:HA	2:B:359:THR:O	2.16	0.46
1:A:1106:GLN:HA	1:A:1109:VAL:HG22	1.97	0.46
2:B:318:CYS:HA	2:B:426:ILE:HG22	1.97	0.46
1:A:188:ARG:NH1	1:A:216:ALA:O	2.49	0.46
1:A:954:MET:HE1	1:A:957:VAL:HG22	1.98	0.46
2:B:351:ASN:OD1	2:B:355:TYR:N	2.48	0.46
3:C:81:TRP:CG	3:C:82:PRO:HD3	2.51	0.46
1:A:1005:ASN:OD1	2:B:238:THR:HA	2.17	0.45
2:B:199:LEU:HD12	2:B:202:LEU:CD1	2.46	0.45
2:B:189:VAL:HG22	2:B:278:ILE:HD11	1.98	0.45
1:A:1005:ASN:HD21	2:B:238:THR:HA	1.82	0.45
1:A:889:ARG:HD2	1:A:891:TYR:CZ	2.51	0.45
1:A:1112:LEU:HD23	1:A:1124:ALA:HB3	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:80:ALA:O	3:C:84:GLN:HG3	2.16	0.44
1:A:382:PHE:H	1:A:720:SER:HG	1.64	0.44
1:A:331:SER:HB2	1:A:353:PHE:HB2	1.99	0.44
1:A:218:MET:HE1	1:A:258:ILE:HG22	2.00	0.43
3:C:71:LEU:HD13	3:C:114:LEU:HD12	1.99	0.43
1:A:985:THR:O	1:A:989:ARG:HG3	2.19	0.43
1:A:1005:ASN:ND2	2:B:238:THR:HA	2.34	0.43
2:B:352:PRO:HG3	2:B:378:HIS:CG	2.53	0.43
1:A:933:LEU:HD23	1:A:944:GLU:HA	2.01	0.42
1:A:785:GLU:HB2	2:B:229:LYS:NZ	2.34	0.42
3:C:77:HIS:CE1	3:C:79:PHE:HB2	2.54	0.42
1:A:886:SER:HA	1:A:910:MET:HA	2.02	0.42
1:A:795:ASP:HB2	1:A:802:LEU:HD11	2.01	0.41
2:B:288:LEU:HD23	2:B:288:LEU:HA	1.85	0.41
1:A:973:ASN:OD1	1:A:999:HIS:ND1	2.52	0.41
1:A:1133:VAL:O	1:A:1137:THR:HG23	2.20	0.41
2:B:257:ILE:HG21	2:B:284:VAL:HG21	2.03	0.41
1:A:1071:SER:O	1:A:1075:SER:OG	2.32	0.41
2:B:326:CYS:C	2:B:328:GLU:H	2.29	0.41
1:A:850:VAL:HB	1:A:862:ALA:HB3	2.01	0.41
2:B:232:PHE:O	2:B:242:ARG:HG2	2.20	0.41
1:A:273:LEU:O	1:A:280:LEU:HD12	2.21	0.41
1:A:130:MET:HE3	1:A:176:PRO:HG2	2.03	0.40
1:A:954:MET:HA	1:A:969:GLU:HA	2.03	0.40
2:B:141:ILE:HG23	2:B:157:VAL:HG13	2.03	0.40
2:B:323:CYS:HB3	2:B:326:CYS:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	798/864 (92%)	742 (93%)	47 (6%)	9 (1%)	11	45
2	B	371/463 (80%)	348 (94%)	22 (6%)	1 (0%)	36	71
3	C	125/127 (98%)	122 (98%)	3 (2%)	0	100	100
All	All	1294/1454 (89%)	1212 (94%)	72 (6%)	10 (1%)	16	52

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	367	LEU
1	A	368	GLU
1	A	1076	PHE
1	A	36	ASN
1	A	317	LEU
1	A	1112	LEU
1	A	746	SER
2	B	50	ASP
1	A	48	GLY
1	A	310	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	680/749 (91%)	680 (100%)	0	100	100
2	B	335/416 (80%)	335 (100%)	0	100	100
3	C	119/120 (99%)	119 (100%)	0	100	100
All	All	1134/1285 (88%)	1134 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	107	ASN
1	A	203	ASN
1	A	319	ASN
1	A	991	HIS
1	A	1025	GLN
1	A	1049	ASN
2	B	95	GLN
2	B	148	GLN
2	B	225	GLN
2	B	297	GLN
3	C	52	ASN
3	C	64	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	808/864 (93%)	0.34	51 (6%) 26 27	212, 287, 355, 409	0
2	B	375/463 (80%)	0.51	31 (8%) 17 21	197, 250, 310, 336	0
3	C	127/127 (100%)	0.44	11 (8%) 16 19	232, 284, 344, 359	0
All	All	1310/1454 (90%)	0.40	93 (7%) 22 24	197, 275, 350, 409	0

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	TYR	6.7
1	A	273	LEU	4.6
2	B	104	PRO	4.4
2	B	49	PHE	4.2
1	A	957	VAL	4.1
2	B	340	LEU	4.0
1	A	281	PHE	3.9
1	A	958	GLU	3.9
1	A	272	LEU	3.9
1	A	966	LEU	3.8
1	A	283	LEU	3.7
3	C	92	LEU	3.6
1	A	787	GLU	3.6
1	A	282	MET	3.5
1	A	328	LEU	3.5
2	B	157	VAL	3.5
3	C	159	GLN	3.4
1	A	922	LEU	3.4
2	B	89	MET	3.3
2	B	117	ASP	3.3
3	C	74	LEU	3.3
1	A	305	LEU	3.3
3	C	81	TRP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	719	GLU	3.1
2	B	233	HIS	3.1
1	A	717	LEU	3.1
1	A	245	TYR	3.0
1	A	336	LEU	3.0
2	B	388	VAL	3.0
3	C	91	LYS	3.0
1	A	132	GLY	3.0
1	A	965	PHE	3.0
1	A	1007	PHE	3.0
3	C	70	VAL	3.0
2	B	348	ALA	3.0
2	B	236	ASN	2.9
2	B	359	THR	2.9
1	A	926	LEU	2.9
1	A	243	ASP	2.9
2	B	292	ASP	2.9
2	B	352	PRO	2.9
3	C	73	THR	2.9
2	B	105	GLN	2.8
1	A	382	PHE	2.8
1	A	960	LEU	2.7
3	C	157	PHE	2.7
2	B	152	ILE	2.7
2	B	144	TYR	2.7
2	B	98	PRO	2.7
1	A	959	ILE	2.7
2	B	242	ARG	2.6
1	A	280	LEU	2.6
2	B	128	VAL	2.6
1	A	1011	SER	2.5
2	B	44	PRO	2.5
1	A	210	GLU	2.5
2	B	390	GLN	2.5
1	A	924	GLY	2.5
2	B	381	PHE	2.5
1	A	48	GLY	2.4
2	B	199	LEU	2.4
1	A	219	VAL	2.4
1	A	913	TYR	2.4
2	B	349	TYR	2.4
1	A	99	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	733	PHE	2.3
1	A	800	GLU	2.3
1	A	140	PHE	2.3
3	C	93	ASN	2.3
1	A	900	ARG	2.2
1	A	1111	ASN	2.2
1	A	1028	VAL	2.2
2	B	69	GLY	2.2
3	C	42	SER	2.2
1	A	148	ASP	2.1
3	C	129	PHE	2.1
1	A	270	ARG	2.1
2	B	380	TRP	2.1
1	A	930	VAL	2.1
2	B	97	LEU	2.1
2	B	343	CYS	2.1
1	A	932	LEU	2.1
1	A	909	ILE	2.1
1	A	860	THR	2.1
1	A	390	ILE	2.1
1	A	176	PRO	2.1
2	B	416	GLY	2.0
1	A	285	LEU	2.0
1	A	225	PRO	2.0
2	B	400	TRP	2.0
1	A	364	VAL	2.0
1	A	1040	VAL	2.0
2	B	85	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	B	501	1/1	1.00	0.09	197,197,197,197	0

6.5 Other polymers [i](#)

There are no such residues in this entry.