



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:48 PM UTC

PDB ID : 8BNY / pdb_00008bny
Title : Structure of the tetramerization domain of pLS20 conjugation repressor Rco
Authors : Bernardo, N.; Crespo, I.; Meijer, W.J.J.; Boer, D.R.
Deposited on : 2022-11-14
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4041 (1.44-1.40)
Clashscore	190562	4154 (1.44-1.40)
Ramachandran outliers	187476	4083 (1.44-1.40)
Sidechain outliers	187428	4082 (1.44-1.40)
RSRZ outliers	180081	4039 (1.44-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	4% 18% 78%
1	B	161	% 21% 79%
1	C	161	% 20% 78%
1	D	161	3% 19% 78%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Immunity repressor protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	35	Total	C	N	O	S	0	0	0
			300	194	50	55	1			
1	B	34	Total	C	N	O	S	0	1	0
			297	190	51	55	1			
1	C	35	Total	C	N	O	S	0	0	0
			297	191	49	56	1			
1	D	36	Total	C	N	O	S	0	0	0
			306	197	51	57	1			


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	11	Total	O	0	0
			11	11		
3	B	18	Total	O	0	0
			18	18		
3	C	17	Total	O	0	0
			17	17		
3	D	16	Total	O	0	0
			16	16		

- Molecule 1: Immunity repressor protein





MET	GLY	ASN	ARG	GLU	GLN	PHE	ASP	LEU	SER	LYS	TYR	LEU	LEU	LEU	GLU	LYS	ARG	LYS	GLN	LYS	GLY	LEU	SER	SER	SER	ALA	TYR	ILE	ILE	SER	MET	LEU	GLU	LYS	GLY	GLU	ARG	LYS	LYS	ARG	PRO	THR	GLN	ALA	VAL	ILE	ILE	LYS	LYS	LEU	ALA	THR	ALA
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

LEU	SER	ILE	PRO	ASN	ILE	GLU	MET	GLN	LEU	ILE	GLN	ILE	MET	GLU	TYR	ALA	ALA	MET	GLU	GLU	ASP	HIS	VAL	LYS	GLU	LYS	ASP	LYS	ASP	THR	SER	LYS	GLN	ILE	LYS	LEU	PHE	ASP	LEU	THR	GLY	LEU	THR	GLU	ALA	SER	ILE	ILE	ASN	GLN	ILE	ASP	GLN	GLN	LYS	LEU	LYS	SER	SER
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

GLU	ASN	ASN	S124	V125	I126	F129	T132	G133	L134	S135	R152	Y155	K159	LYS	ASP
-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	35.33Å 36.18Å 109.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.36 – 1.43 34.36 – 1.43	Depositor EDS
% Data completeness (in resolution range)	69.2 (34.36-1.43) 69.2 (34.36-1.43)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.67 (at 1.43Å)	Xtriage
Refinement program	PHENIX 1.12-2829	Depositor
R, R_{free}	0.233 , 0.249 0.234 , 0.251	Depositor DCC
R_{free} test set	905 reflections (3.24%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.038 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1263	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/302	0.46	0/400
1	B	0.26	0/302	0.41	0/400
1	C	0.31	0/299	0.48	0/397
1	D	0.27	0/308	0.45	0/408
All	All	0.26	0/1211	0.45	0/1605

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	300	0	319	4	0
1	B	297	0	313	0	0
1	C	297	0	311	2	0
1	D	306	0	324	2	0
2	A	1	0	0	0	0
3	A	11	0	0	0	0
3	B	18	0	0	0	0
3	C	17	0	0	0	0
3	D	16	0	0	0	0
All	All	1263	0	1267	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:ARG:O	1:A:156:MET:HG2	2.12	0.49
1:A:144:GLU:OE1	1:C:141:ARG:NH1	2.50	0.45
1:A:125:VAL:HB	1:D:132:THR:HA	2.01	0.42
1:A:136:GLU:O	1:A:140:GLU:HG2	2.19	0.42
1:C:144:GLU:OE2	1:D:152:ARG:NH2	2.33	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	33/161 (20%)	32 (97%)	1 (3%)	0	100	100
1	B	33/161 (20%)	32 (97%)	1 (3%)	0	100	100
1	C	33/161 (20%)	32 (97%)	1 (3%)	0	100	100
1	D	34/161 (21%)	33 (97%)	1 (3%)	0	100	100
All	All	133/644 (21%)	129 (97%)	4 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	34/148 (23%)	34 (100%)	0	100	100
1	B	34/148 (23%)	34 (100%)	0	100	100
1	C	34/148 (23%)	34 (100%)	0	100	100
1	D	35/148 (24%)	32 (91%)	3 (9%)	10	0
All	All	137/592 (23%)	134 (98%)	3 (2%)	45	13

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	134	LEU
1	D	135	SER
1	D	155	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	35/161 (21%)	1.31	7 (20%) 3 3	25, 40, 81, 99	0
1	B	34/161 (21%)	0.52	1 (2%) 53 57	20, 28, 53, 70	1 (2%)
1	C	35/161 (21%)	0.37	1 (2%) 53 57	15, 27, 46, 67	0
1	D	36/161 (22%)	0.94	5 (13%) 6 8	21, 38, 61, 65	0
All	All	140/644 (21%)	0.79	14 (10%) 12 13	15, 35, 64, 99	1 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	LEU	4.7
1	C	158	LEU	4.3
1	D	126	ILE	3.6
1	A	126	ILE	3.1
1	A	157	LYS	3.0
1	A	125	VAL	2.9
1	A	155	TYR	2.7
1	D	134	LEU	2.6
1	D	125	VAL	2.4
1	A	159	LYS	2.2
1	D	129	PHE	2.1
1	B	157	LYS	2.1
1	A	128	PHE	2.1
1	D	133	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CL	A	201	1/1	0.78	0.19	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.