



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:48 PM UTC

PDB ID : 2BOP / pdb_00002bop
Title : CRYSTAL STRUCTURE AT 1.7 ANGSTROMS OF THE BOVINE PAPILLOMAVIRUS-1 E2 DNA-BINDING DOMAIN BOUND TO ITS DNA TARGET
Authors : Hegde, R.S.; Grossman, S.R.; Laimins, L.A.; Sigler, P.B.
Deposited on : 1994-01-13
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	17	
2	A	85	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*CP*GP*AP*CP*CP*GP*AP*CP*GP*TP*CP*GP*GP*TP*CP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	17	Total	C	N	O	P	0	0	0
			344	163	65	100	16			

- Molecule 2 is a protein called PROTEIN (E2).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	85	Total	C	N	O	S	0	0	0
			677	426	125	122	4			

- Molecule 3 is YTTERBIUM (III) ION (CCD ID: YB) (formula: Yb).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Yb	0	0
			1	1		
3	A	1	Total	Yb	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	52	Total	O	0	0
			52	52		
4	A	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*CP*CP*GP*AP*CP*CP*GP*AP*CP*GP*TP*CP*GP*GP*TP*CP*G)-3')

Chain B: 



- Molecule 2: PROTEIN (E2)

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	R 3 2	Depositor
Cell constants a, b, c, α , β , γ	64.60Å 64.60Å 64.60Å 61.80° 61.80° 61.80°	Depositor
Resolution (Å)	6.00 – 1.70	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-1.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	NUCLSQ, PROFFT	Depositor
R, R_{free}	0.201 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1144	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: YB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	2.31	15/385 (3.9%)	2.32	23/592 (3.9%)
2	A	1.77	8/694 (1.2%)	2.07	18/936 (1.9%)
All	All	1.98	23/1079 (2.1%)	2.17	41/1528 (2.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4
2	A	1	6
All	All	1	10

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	9	DC	P-O5'	11.29	1.83	1.60
1	B	11	DT	P-OP1	-11.07	1.26	1.48
1	B	17	DG	P-O5'	-9.94	1.40	1.60
1	B	9	DC	O3'-P	9.39	1.75	1.61
1	B	8	DA	P-OP2	-9.07	1.30	1.48
1	B	17	DG	P-OP1	8.71	1.65	1.48
1	B	10	DG	O3'-P	-8.14	1.49	1.61
1	B	7	DG	P-OP1	7.66	1.63	1.48
1	B	2	DC	P-OP2	-7.54	1.33	1.48
1	B	6	DC	P-OP2	-7.49	1.33	1.48
1	B	16	DC	O3'-P	7.21	1.72	1.61
1	B	16	DC	P-O5'	-6.62	1.47	1.60
1	B	9	DC	P-OP2	-6.27	1.35	1.48
2	A	365	ASP	C-O	6.09	1.31	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	340	CYS	N-CA	6.08	1.53	1.46
2	A	342	ARG	N-CA	5.75	1.53	1.46
1	B	16	DC	P-OP2	-5.70	1.37	1.48
2	A	381	GLY	N-CA	5.45	1.52	1.45
2	A	348	ASN	C-O	5.28	1.30	1.23
2	A	399	MET	N-CA	5.24	1.51	1.45
1	B	2	DC	O3'-P	5.17	1.69	1.61
2	A	394	PRO	C-N	-5.14	1.27	1.33
2	A	339	LYS	C-O	5.09	1.29	1.24

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	400	ASN	CA-CB-CG	-10.57	102.03	112.60
1	B	17	DG	O5'-P-OP2	9.84	137.52	108.00
2	A	387	GLN	CA-C-O	-8.26	111.80	120.55
1	B	7	DG	P-O5'-C5'	-8.02	107.97	120.00
2	A	392	HIS	CA-CB-CG	-7.98	105.82	113.80
2	A	326	SER	N-CA-CB	7.40	123.09	110.50
2	A	396	PRO	CA-C-O	7.14	126.04	120.90
2	A	352	ARG	CD-NE-CZ	-7.04	114.55	124.40
1	B	17	DG	O5'-C5'-C4'	-7.01	100.28	110.80
1	B	3	DG	P-O5'-C5'	6.95	130.42	120.00
1	B	5	DC	C5'-C4'-C3'	6.66	124.89	114.90
1	B	3	DG	O3'-P-O5'	-6.57	94.15	104.00
1	B	8	DA	P-O5'-C5'	-6.56	110.16	120.00
1	B	7	DG	OP1-P-OP2	-6.54	100.37	120.00
2	A	337	GLN	O-C-N	6.44	128.70	122.07
1	B	16	DC	O3'-P-O5'	-6.39	94.42	104.00
1	B	9	DC	O5'-C5'-C4'	-6.31	101.34	110.80
1	B	14	DG	OP1-P-O3'	-6.17	89.49	108.00
2	A	403	GLY	O-C-N	-6.06	117.79	123.48
2	A	355	ASN	CA-CB-CG	-5.97	106.63	112.60
1	B	9	DC	O3'-P-O5'	-5.90	95.15	104.00
1	B	15	DT	N3-C4-O4	-5.83	113.86	122.60
2	A	379	THR	CA-CB-OG1	-5.70	101.06	109.60
2	A	402	SER	CA-C-O	-5.69	114.45	121.11
1	B	16	DC	N1-C1'-C2'	-5.67	105.00	113.50
1	B	16	DC	O5'-P-OP2	5.50	124.51	108.00
2	A	370	ARG	NE-CZ-NH1	5.50	127.00	121.50
2	A	384	SER	N-CA-CB	5.44	117.90	110.01
2	A	373	GLN	CB-CG-CD	-5.44	103.35	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	DA	C5'-C4'-C3'	5.40	123.00	114.90
1	B	17	DG	OP1-P-OP2	-5.40	103.81	120.00
1	B	7	DG	O5'-C5'-C4'	-5.37	102.75	110.80
1	B	8	DA	O5'-C5'-C4'	-5.36	102.76	110.80
2	A	326	SER	CB-CA-C	5.35	120.27	110.10
2	A	370	ARG	CA-C-O	-5.28	115.17	121.56
2	A	393	VAL	N-CA-CB	-5.27	104.44	111.08
1	B	9	DC	O5'-P-OP1	-5.27	93.19	109.00
1	B	6	DC	P-O5'-C5'	5.24	127.86	120.00
1	B	2	DC	C2'-C3'-O3'	5.21	119.31	111.50
2	A	344	ARG	CD-NE-CZ	5.16	131.62	124.40
1	B	10	DG	OP1-P-OP2	5.10	135.31	120.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	326	SER	CA

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	348	ASN	Mainchain
2	A	371	GLN	Sidechain
2	A	381	GLY	Mainchain
2	A	391	LYS	Mainchain
2	A	403	GLY	Mainchain
2	A	406	ALA	Mainchain
1	B	12	DC	Sidechain
1	B	6	DC	Sidechain
1	B	8	DA	Sidechain
1	B	9	DC	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	344	0	191	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	677	0	657	8	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	69	0	0	2	2
4	B	52	0	0	1	0
All	All	1144	0	848	10	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:DC:OP2	4:B:621:HOH:O	1.84	0.96
2:A:400:ASN:OD1	4:A:545:HOH:O	1.86	0.92
2:A:351:HIS:ND1	4:A:568:HOH:O	2.00	0.86
2:A:364:ALA:HB2	2:A:371:GLN:HG2	1.81	0.62
2:A:338:VAL:CG1	2:A:376:ILE:HG23	2.37	0.54
2:A:396:PRO:HG2	2:A:399:MET:SD	2.53	0.49
2:A:400:ASN:OD1	2:A:401:ILE:N	2.47	0.47
1:B:10:DG:H2''	1:B:11:DT:H72	1.98	0.45
2:A:338:VAL:HG11	2:A:376:ILE:HG23	2.01	0.41
2:A:347:LYS:HE3	2:A:347:LYS:HB3	1.63	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:569:HOH:O	4:A:569:HOH:O[4_776]	1.65	0.55
4:A:540:HOH:O	4:A:587:HOH:O[4_776]	2.07	0.13

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	83/85 (98%)	80 (96%)	3 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	73/73 (100%)	69 (94%)	4 (6%)	19	6

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	330	LEU
2	A	331	ILE
2	A	345	VAL
2	A	371	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	336	ASN
2	A	348	ASN
2	A	392	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.