



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 01:32 PM UTC

PDB ID : 6BOX / pdb_00006box
Title : Structure of the *S. pombe* Clr4 catalytic domain bound to SAH
Authors : Currie, M.A.; Moazed, D.
Deposited on : 2017-11-21
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6062 (2.44-2.40)
Clashscore	190562	6562 (2.44-2.40)
Ramachandran outliers	187476	6481 (2.44-2.40)
Sidechain outliers	187428	6482 (2.44-2.40)
RSRZ outliers	180081	6066 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	12% 84% 5% 11%
1	B	303	11% 82% 6% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8428 atoms, of which 4035 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-lysine N-methyltransferase, H3 lysine-9 specific.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	270	Total	C	H	N	O	S	0	0	0
			4160	1357	2006	368	412	17			
1	B	267	Total	C	H	N	O	S	0	0	0
			4112	1331	1991	364	409	17			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	188	GLY	-	expression tag	UNP O60016
A	189	PRO	-	expression tag	UNP O60016
A	190	LEU	-	expression tag	UNP O60016
A	191	GLY	-	expression tag	UNP O60016
B	188	GLY	-	expression tag	UNP O60016
B	189	PRO	-	expression tag	UNP O60016
B	190	LEU	-	expression tag	UNP O60016
B	191	GLY	-	expression tag	UNP O60016

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Zn	0	0
			4	4		
2	B	4	Total	Zn	0	0
			4	4		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	S	0	0
			45	14	19	6	5	1		
3	B	1	Total	C	H	N	O	S	0	0
			45	14	19	6	5	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	28	Total	O	0	0
			28	28		

i

- Molecule 1: Histone-lysine N-methyltransferase, H3 lysine-9 specific



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	125.62Å 71.25Å 87.24Å 90.00° 120.37° 90.00°	Depositor
Resolution (Å)	54.19 – 2.41 54.19 – 2.41	Depositor EDS
% Data completeness (in resolution range)	96.6 (54.19-2.41) 96.7 (54.19-2.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.42Å)	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.231 , 0.276 0.233 , 0.278	Depositor DCC
R_{free} test set	1269 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtriage
Anisotropy	0.438	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 29.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.000 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8428	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2204	0.31	0/2979
1	B	0.11	0/2169	0.30	0/2934
All	All	0.14	0/4373	0.30	0/5913

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2154	2006	2028	9	0
1	B	2121	1991	1996	12	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	26	19	19	1	0
3	B	26	19	19	0	0
4	A	30	0	0	0	0
4	B	28	0	0	0	0
All	All	4393	4035	4062	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:381:TYR:OH	1:B:406:ARG:NH1	2.24	0.70
1:B:320:ARG:HD3	1:B:323:GLN:OE1	1.93	0.68
1:B:315:MET:SD	1:B:320:ARG:NH1	2.70	0.64
1:B:313:CYS:O	1:B:320:ARG:NH2	2.34	0.60
1:A:306:GLU:OE1	1:A:423:ARG:NH2	2.38	0.56
1:B:393:GLU:N	1:B:393:GLU:OE1	2.39	0.56
1:B:423:ARG:NH1	1:B:432:ASP:OD2	2.40	0.55
1:A:485:ARG:HE	1:A:487:TRP:HZ2	1.53	0.53
1:B:290:TYR:OH	1:B:306:GLU:OE2	2.24	0.50
1:A:275:ARG:NE	1:A:275:ARG:HA	2.28	0.49
1:A:423:ARG:NH1	1:A:432:ASP:OD2	2.47	0.48
3:A:505:SAH:H5'2	3:A:505:SAH:H8	1.96	0.48
1:A:234:SER:OG	1:A:236:ASP:OD1	2.34	0.45
1:B:284:GLU:HB2	1:B:285:PRO:HD3	1.98	0.45
1:B:283:ASP:O	1:B:286:THR:HG22	2.17	0.45
1:B:320:ARG:CD	1:B:323:GLN:OE1	2.64	0.43
1:A:274:SER:O	1:A:275:ARG:HB2	2.19	0.43
1:A:457:PHE:CE1	1:B:487:TRP:HZ2	2.37	0.42
1:A:451:TYR:HB2	1:A:457:PHE:CE1	2.56	0.41
1:A:381:TYR:OH	1:A:406:ARG:NH1	2.54	0.41
1:B:388:PHE:CZ	1:B:422:VAL:HG21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/303 (86%)	251 (96%)	10 (4%)	1 (0%)	30	42
1	B	261/303 (86%)	251 (96%)	10 (4%)	0	100	100
All	All	523/606 (86%)	502 (96%)	20 (4%)	1 (0%)	43	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	263	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/264 (89%)	236 (100%)	0	100	100
1	B	232/264 (88%)	232 (100%)	0	100	100
All	All	468/528 (89%)	468 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	238	GLN
1	A	425	HIS
1	B	238	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 8 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SAH	B	505	-	27,28,28	3.02	9 (33%)	36,40,40	2.38	13 (36%)
3	SAH	A	505	-	27,28,28	3.00	9 (33%)	36,40,40	2.36	12 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	B	505	-	-	3/15/31/31	0/3/3/3
3	SAH	A	505	-	-	6/15/31/31	0/3/3/3

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	505	SAH	O4'-C1'	8.44	1.61	1.42
3	A	505	SAH	O4'-C1'	8.29	1.61	1.42
3	A	505	SAH	O4'-C4'	-6.84	1.29	1.45
3	B	505	SAH	O4'-C4'	-6.75	1.30	1.45
3	A	505	SAH	C2'-C1'	-6.61	1.32	1.53
3	B	505	SAH	C2'-C1'	-6.51	1.33	1.53
3	B	505	SAH	C6-N6	5.89	1.49	1.34
3	A	505	SAH	C6-N6	5.83	1.49	1.34
3	B	505	SAH	O2'-C2'	2.92	1.50	1.43
3	A	505	SAH	O2'-C2'	2.90	1.50	1.43
3	B	505	SAH	C5-C4	-2.60	1.34	1.39
3	A	505	SAH	C5-C4	-2.53	1.34	1.39
3	A	505	SAH	C5-N7	-2.35	1.34	1.39
3	B	505	SAH	C5-N7	-2.34	1.34	1.39
3	A	505	SAH	O3'-C3'	-2.34	1.37	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	505	SAH	O3'-C3'	-2.32	1.37	1.43
3	A	505	SAH	C8-N9	-2.10	1.34	1.37
3	B	505	SAH	C8-N9	-2.08	1.34	1.37

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	505	SAH	N6-C6-N1	-5.96	105.10	118.38
3	A	505	SAH	N6-C6-N1	-5.82	105.40	118.38
3	B	505	SAH	N3-C2-N1	-5.56	120.17	128.58
3	A	505	SAH	N3-C2-N1	-5.42	120.38	128.58
3	B	505	SAH	C5-C6-N6	4.90	135.42	123.29
3	A	505	SAH	C5-C4-N3	-4.86	120.02	126.72
3	A	505	SAH	C5-C6-N6	4.69	134.91	123.29
3	B	505	SAH	C5-C4-N3	-4.68	120.27	126.72
3	B	505	SAH	N9-C8-N7	-4.29	107.84	113.94
3	A	505	SAH	N9-C8-N7	-4.12	108.10	113.94
3	A	505	SAH	C5'-SD-CG	3.31	112.09	102.26
3	A	505	SAH	C2-N3-C4	3.21	119.67	111.83
3	B	505	SAH	C5-N7-C8	3.18	108.45	103.45
3	B	505	SAH	C2-N3-C4	3.17	119.58	111.83
3	A	505	SAH	C5-N7-C8	3.00	108.17	103.45
3	A	505	SAH	N3-C4-N9	2.93	132.15	127.17
3	B	505	SAH	N3-C4-N9	2.66	131.70	127.17
3	B	505	SAH	C5'-SD-CG	2.52	109.74	102.26
3	B	505	SAH	C4-C5-N7	-2.50	107.73	110.58
3	A	505	SAH	C4-C5-N7	-2.27	107.98	110.58
3	A	505	SAH	C6-C5-C4	2.13	120.09	117.18
3	B	505	SAH	C4-N9-C8	2.11	107.95	105.74
3	A	505	SAH	C4-N9-C8	2.08	107.92	105.74
3	B	505	SAH	C5'-C4'-C3'	-2.07	109.89	115.06
3	B	505	SAH	C5-C4-N9	2.03	108.03	105.81

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	505	SAH	N-CA-CB-CG
3	A	505	SAH	C-CA-CB-CG
3	A	505	SAH	OXT-C-CA-N
3	A	505	SAH	O-C-CA-N
3	A	505	SAH	C3'-C4'-C5'-SD

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	505	SAH	CB-CG-SD-C5'
3	A	505	SAH	CB-CG-SD-C5'
3	B	505	SAH	O-C-CA-CB
3	B	505	SAH	OXT-C-CA-CB


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	505	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/303 (89%)	1.05	37 (13%) 6 5	28, 46, 81, 172	0
1	B	267/303 (88%)	0.94	34 (12%) 8 6	27, 47, 83, 155	0
All	All	537/606 (88%)	1.00	71 (13%) 7 5	27, 46, 83, 172	0

All (71) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	379	ILE	6.9
1	A	263	SER	6.6
1	B	284	GLU	6.4
1	A	485	ARG	6.3
1	B	380	THR	5.6
1	A	266	GLY	5.6
1	A	265	LEU	5.5
1	B	488	LEU	5.3
1	A	275	ARG	5.3
1	A	264	SER	5.2
1	B	381	TYR	5.1
1	A	284	GLU	4.9
1	A	380	THR	4.9
1	A	487	TRP	4.4
1	A	389	ASP	4.3
1	A	381	TYR	4.1
1	B	391	ALA	3.9
1	B	265	LEU	3.9
1	A	427	PHE	3.9
1	B	308	ASN	3.9
1	B	274	SER	3.9
1	B	285	PRO	3.7
1	A	195	ASP	3.7
1	B	286	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	266	GLY	3.6
1	A	308	ASN	3.4
1	B	196	SER	3.3
1	B	371	ASP	3.3
1	A	261	ASN	3.3
1	B	261	ASN	3.2
1	B	320	ARG	3.2
1	B	267	GLY	3.2
1	B	262	CYS	3.2
1	A	196	SER	3.2
1	B	418	ILE	3.2
1	A	262	CYS	3.1
1	A	474	ARG	3.1
1	A	197	TYR	3.0
1	A	475	ARG	3.0
1	A	283	ASP	3.0
1	B	453	GLY	3.0
1	B	487	TRP	2.9
1	A	473	LEU	2.9
1	B	283	ASP	2.9
1	B	337	GLU	2.8
1	B	382	LEU	2.8
1	B	389	ASP	2.7
1	B	474	ARG	2.7
1	B	263	SER	2.7
1	A	318	PRO	2.6
1	A	236	ASP	2.6
1	A	386	ASP	2.6
1	A	453	GLY	2.5
1	A	267	GLY	2.5
1	A	337	GLU	2.5
1	B	344	SER	2.5
1	A	198	THR	2.5
1	A	486	GLY	2.4
1	B	275	ARG	2.4
1	A	368	ALA	2.4
1	A	286	THR	2.3
1	A	483	ASN	2.3
1	B	383	PHE	2.3
1	B	197	TYR	2.2
1	A	274	SER	2.2
1	B	388	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	313	CYS	2.1
1	A	215	GLU	2.1
1	A	207	GLU	2.1
1	A	273	PRO	2.1
1	B	392	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SAH	A	505	26/26	0.89	0.12	37,48,61,63	0
3	SAH	B	505	26/26	0.93	0.09	29,44,60,77	0
2	ZN	A	502	1/1	0.96	0.04	31,31,31,31	0
2	ZN	B	502	1/1	0.97	0.06	38,38,38,38	0
2	ZN	B	504	1/1	0.98	0.04	40,40,40,40	0
2	ZN	B	501	1/1	0.98	0.04	41,41,41,41	0
2	ZN	B	503	1/1	0.98	0.04	34,34,34,34	0
2	ZN	A	501	1/1	0.99	0.03	49,49,49,49	0
2	ZN	A	503	1/1	0.99	0.05	37,37,37,37	0
2	ZN	A	504	1/1	0.99	0.05	44,44,44,44	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SAH A 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 505:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.