



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 02:44 PM UTC

PDB ID : 1BPR / pdb_00001bpr
Title : NMR STRUCTURE OF THE SUBSTRATE BINDING DOMAIN OF DNAK,
MINIMIZED AVERAGE STRUCTURE
Authors : Wang, H.; Kurochkin, A.V.; Pang, Y.; Hu, W.; Flynn, G.C.; Zuiderweg, E.R.P.
Deposited on : 1998-08-11

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	191	 63% 23% . . 9%

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2655 atoms, of which 1330 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNAK.

Mol	Chain	Residues	Atoms						Trace
1	A	173	Total	C	H	N	O	S	0
			2655	812	1330	239	270	4	

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	381	SER	VAL	conflict	UNP P0A6Y8
A	382	ILE	LEU	conflict	UNP P0A6Y8
A	383	GLU	THR	conflict	UNP P0A6Y8
A	385	ARG	ASP	conflict	UNP P0A6Y8

4 Residue-property plots

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: DNAK

Chain A: 



5 Refinement protocol and experimental data overview

The models were refined using the following method: *DISTANCE GEOMETRY AND RESTRAINED MOLECULAR DYNAMICS WITH SIMULATED ANNEALING*.

Of the 100 calculated structures, 1 were deposited, based on the following criterion: *TOTAL ENERGY*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Discover	refinement	
BIOSYM	structure solution	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.87	0/1338 (0.0%)	1.81	33/1803 (1.8%)
All	All	0.87	0/1338 (0.0%)	1.81	33/1803 (1.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0	1
All	All	0	1

There are no bond-length outliers.

All angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	511	GLU	N-CA-C	-14.41	92.42	112.45
1	A	485	HIS	ND1-CE1-NE2	12.60	121.00	108.40
1	A	544	HIS	ND1-CE1-NE2	12.57	120.97	108.40
1	A	422	HIS	ND1-CE1-NE2	12.54	120.94	108.40
1	A	541	HIS	ND1-CE1-NE2	12.54	120.94	108.40
1	A	439	HIS	ND1-CE1-NE2	12.50	120.90	108.40
1	A	449	ALA	N-CA-C	-11.37	96.84	112.30
1	A	520	GLU	N-CA-C	-9.81	99.87	112.23
1	A	439	HIS	CE1-NE2-CD2	-9.60	99.40	109.00
1	A	485	HIS	CE1-NE2-CD2	-9.55	99.45	109.00
1	A	422	HIS	CE1-NE2-CD2	-9.54	99.46	109.00
1	A	541	HIS	CE1-NE2-CD2	-9.54	99.46	109.00
1	A	544	HIS	CE1-NE2-CD2	-9.54	99.46	109.00
1	A	454	LEU	N-CA-C	-9.33	98.68	111.56
1	A	520	GLU	N-CA-CB	6.77	120.73	110.30
1	A	516	VAL	N-CA-C	-6.45	103.97	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	440	VAL	CB-CA-C	6.41	118.85	110.91
1	A	544	HIS	CG-ND1-CE1	-6.39	98.43	109.30
1	A	485	HIS	CG-ND1-CE1	-6.39	98.43	109.30
1	A	541	HIS	CG-ND1-CE1	-6.39	98.44	109.30
1	A	422	HIS	CG-ND1-CE1	-6.37	98.48	109.30
1	A	492	ASN	N-CA-C	-6.31	105.23	113.12
1	A	439	HIS	CG-ND1-CE1	-6.28	98.63	109.30
1	A	411	LEU	N-CA-C	-6.13	103.31	111.96
1	A	411	LEU	N-CA-CB	5.38	119.90	110.27
1	A	531	GLU	N-CA-C	-5.30	105.56	112.23
1	A	537	ASN	N-CA-CB	5.15	119.19	110.49
1	A	422	HIS	CB-CG-CD2	-5.12	124.54	131.20
1	A	454	LEU	CB-CA-C	5.08	119.11	111.18
1	A	485	HIS	CB-CG-CD2	-5.04	124.65	131.20
1	A	541	HIS	CB-CG-CD2	-5.03	124.66	131.20
1	A	455	GLY	N-CA-C	5.02	117.41	111.63
1	A	544	HIS	CB-CG-CD2	-5.01	124.68	131.20

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
1	A	395	THR	Peptide

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1325	1330	1330	16
All	All	1325	1330	1330	16

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:442:GLN:O	1:A:452:LYS:HB3	1.26	1.30

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:510:ASP:O	1:A:514:LYS:HB2	0.92	1.64
1:A:510:ASP:O	1:A:514:LYS:CB	0.91	2.17
1:A:442:GLN:O	1:A:452:LYS:CB	0.89	2.19
1:A:471:GLN:O	1:A:491:LYS:HB2	0.82	1.75
1:A:516:VAL:HA	1:A:520:GLU:HB2	0.66	1.66
1:A:471:GLN:O	1:A:491:LYS:CB	0.61	2.48
1:A:510:ASP:O	1:A:514:LYS:HB3	0.60	1.95
1:A:400:GLY:O	1:A:440:VAL:HA	0.49	2.08
1:A:508:ASN:O	1:A:511:GLU:HB3	0.48	2.09
1:A:516:VAL:CA	1:A:520:GLU:HB2	0.46	2.40
1:A:552:GLU:O	1:A:553:ALA:HB2	0.44	2.13
1:A:471:GLN:O	1:A:491:LYS:CG	0.44	2.66
1:A:529:PHE:O	1:A:533:VAL:HG23	0.43	2.13
1:A:471:GLN:O	1:A:491:LYS:HG3	0.42	2.14
1:A:524:GLU:O	1:A:527:ARG:HB3	0.41	2.15

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	171/191 (90%)	118 (69%)	35 (20%)	18 (11%)	1	9
All	All	171/191 (90%)	118 (69%)	35 (20%)	18 (11%)	1	9

All 18 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	383	GLU
1	A	384	GLY
1	A	397	LEU
1	A	398	SER
1	A	404	MET
1	A	431	ASP
1	A	445	ARG
1	A	456	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	467	ARG
1	A	468	GLY
1	A	469	MET
1	A	470	PRO
1	A	505	SER
1	A	534	GLN
1	A	535	THR
1	A	536	ARG
1	A	537	ASN
1	A	538	GLN

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/161 (91%)	136 (93%)	11 (7%)	14	62
All	All	147/161 (91%)	136 (93%)	11 (7%)	14	62

All 11 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	383	GLU
1	A	397	LEU
1	A	399	LEU
1	A	440	VAL
1	A	454	LEU
1	A	472	ILE
1	A	475	THR
1	A	516	VAL
1	A	535	THR
1	A	537	ASN
1	A	548	LYS

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided