



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:43 PM UTC

PDB ID : 4BPK / pdb_00004bpk
Title : Bcl-xL bound to alpha beta Puma BH3 peptide 5
Authors : Smith, B.J.; Lee, E.F.; Checco, J.W.; Gellman, S.H.; Fairlie, W.D.
Deposited on : 2013-05-27
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	157	31% 73% 14% • 11%
1	B	157	27% 75% 13% • 12%
2	C	22	23% 55% 32% 5% 5% 5%
2	D	22	9% 68% 14% 14% 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BCL-2-LIKE PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	6	15	0
			1262	809	216	232	5			
1	B	138	Total	C	N	O	S	15	8	0
			1176	751	197	224	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q07817
A	-2	PRO	-	expression tag	UNP Q07817
A	-1	LEU	-	expression tag	UNP Q07817
A	0	GLY	-	expression tag	UNP Q07817
B	-3	GLY	-	expression tag	UNP Q07817
B	-2	PRO	-	expression tag	UNP Q07817
B	-1	LEU	-	expression tag	UNP Q07817
B	0	GLY	-	expression tag	UNP Q07817


- Molecule 2 is a protein called ALPHA BETA BH3-PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	21	Total	C	N	O	S	0	1	1
			180	114	35	30	1			
2	D	22	Total	C	N	O	S	0	1	1
			184	116	37	30	1			

- Molecule 3 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	Cd	0	2
			8	8		
3	B	9	Total	Cd	0	2
			11	11		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	67	Total	O	0	4
			72	72		
5	B	76	Total	O	0	2
			78	78		
5	C	5	Total	O	0	0
			5	5		
5	D	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BCL-2-LIKE PROTEIN 1




• Molecule 1: BCL-2-LIKE PROTEIN 1



• Molecule 2: ALPHA BETA BH3-PEPTIDE



• Molecule 2: ALPHA BETA BH3-PEPTIDE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.17Å 72.13Å 80.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.07 – 1.76 36.07 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.0 (36.07-1.76) 98.7 (36.07-1.76)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.76Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.2_1309)	Depositor
R, R_{free}	0.219 , 0.252 0.237 , 0.267	Depositor DCC
R_{free} test set	1920 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2987	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, B3Q, CD, ACE, HT7, AHP, B3E, B3A, NH2, HR7, B3D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1300	0.70	0/1759
1	B	0.49	0/1210	0.73	0/1637
2	C	0.35	0/97	0.64	0/120
2	D	0.29	0/107	0.58	0/131
All	All	0.47	0/2714	0.71	0/3647

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	7
2	D	0	5
All	All	0	12

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	18	B3A	Mainchain
2	C	4	B3E	Mainchain,Peptide
2	C	8[A]	B3Q	Mainchain,Peptide
2	C	8[B]	B3Q	Mainchain,Peptide
2	D	18	B3A	Mainchain
2	D	4	B3E	Mainchain,Peptide
2	D	8	B3Q	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1262	0	1207	23	0
1	B	1176	0	1120	16	0
2	C	180	0	158	11	0
2	D	184	0	166	6	0
3	A	8	0	0	0	0
3	B	11	0	0	0	0
4	A	4	0	6	1	0
4	B	4	0	6	1	0
5	A	72	0	0	1	0
5	B	78	0	0	2	0
5	C	5	0	0	0	0
5	D	3	0	0	1	0
All	All	2987	0	2663	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:ASN:OD1	1:B:5:ASN:ND2	2.12	0.76
2:C:9:AHP:HA	2:C:12:MET:HE2	1.72	0.71
2:D:4:B3E:OF1	5:D:2001:HOH:O	2.11	0.68
1:A:157:LYS:HB2	1:A:159:MET:HE2	1.79	0.63
1:A:96:GLU:OE2	1:A:100:ARG:NH2	2.34	0.61
1:A:114:ILE:HB	1:A:159:MET:HE1	1.82	0.60
1:A:129:GLU:O	2:C:10:ARG:NH1	2.35	0.60
1:A:129:GLU:OE2	2:C:10:ARG:NH2	2.39	0.55
1:A:125[B]:GLN:OE1	2:C:3:ARG:NH1	2.42	0.53
1:A:16[A]:LYS:NZ	1:B:98:GLU:OE1	2.41	0.52
1:B:136:ASN:HB3	4:B:1205:EDO:H11	1.91	0.52
1:A:97:PHE:HZ	2:C:12:MET:HE3	1.74	0.52
5:A:2059:HOH:O	2:C:4:B3E:HD3	2.10	0.51
1:B:154[B]:SER:OG	1:B:159:MET:O	2.29	0.50
4:A:1202:EDO:H22	1:B:91:ARG:HH22	1.77	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:129:GLU:OE1	2:D:3[A]:ARG:NH1	2.45	0.49
1:B:112:LEU:HD23	2:D:5:ILE:HD13	1.95	0.49
1:A:175:ASN:HD21	1:B:2:SER:HA	1.77	0.48
1:A:1:MET:HE3	1:A:1:MET:HB3	1.71	0.47
1:B:108:LEU:HD11	2:D:9:AHP:HE2	1.97	0.47
1:B:193:GLU:HG3	5:B:2072:HOH:O	2.13	0.47
1:A:121:GLN:O	1:A:125[A]:GLN:HG2	2.15	0.46
1:A:99:LEU:HD12	1:A:102[B]:ARG:HE	1.81	0.46
1:A:97:PHE:CZ	2:C:12:MET:HE3	2.51	0.46
1:A:132[A]:ARG:HB3	2:C:10:ARG:HH12	1.80	0.46
1:A:158:GLU:CD	1:A:160[A]:GLN:HE22	2.23	0.46
1:A:5:ASN:HB3	1:B:174:LEU:HD23	1.99	0.45
1:B:97:PHE:CZ	2:D:12:MET:HE2	2.52	0.45
1:B:146:PHE:HB2	2:D:9:AHP:HZ2	1.99	0.44
1:A:160[B]:GLN:O	1:A:160[B]:GLN:HG3	2.19	0.43
2:C:9:AHP:HD2	2:C:12:MET:CE	2.49	0.43
1:B:92:GLU:HB3	1:B:195:TYR:OH	2.19	0.43
1:A:16[B]:LYS:HG3	1:B:148:GLY:HA3	2.01	0.43
1:A:113[B]:HIS:HB2	2:C:1:HT7:HH2	2.01	0.43
1:A:129:GLU:HB2	1:A:132[B]:ARG:HH21	1.83	0.42
1:A:99:LEU:HA	1:A:102[B]:ARG:HG3	2.00	0.42
1:B:112:LEU:HD22	5:B:2037:HOH:O	2.20	0.42
2:C:10:ARG:HD2	2:C:11:HR7:HD3	2.01	0.41
1:A:3:GLN:O	1:A:7:GLU:HG3	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	152/157 (97%)	146 (96%)	6 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	144/157 (92%)	143 (99%)	1 (1%)	0	100	100
2	C	13/22 (59%)	13 (100%)	0	0	100	100
2	D	14/22 (64%)	14 (100%)	0	0	100	100
All	All	323/358 (90%)	316 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/130 (103%)	128 (96%)	6 (4%)	24	7
1	B	126/130 (97%)	121 (96%)	5 (4%)	28	9
2	C	9/9 (100%)	9 (100%)	0	100	100
2	D	10/9 (111%)	8 (80%)	2 (20%)	1	0
All	All	279/278 (100%)	266 (95%)	13 (5%)	26	6

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	132[A]	ARG
1	A	132[B]	ARG
1	A	133	ASP
1	A	161	VAL
1	A	174	LEU
1	B	108	LEU
1	B	124	GLU
1	B	157	LYS
1	B	160	GLN
1	B	175	ASN
2	D	3[A]	ARG
2	D	3[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	GLN
1	B	160	GLN
1	B	183	GLN
2	D	17	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

15 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HT7	C	1	2	16,16,17	0.52	0	18,21,23	0.64	0
2	B3E	C	4	2	9,9,10	0.82	0	8,10,12	1.09	1 (12%)
2	HR7	D	11	2	11,11,12	0.74	0	9,12,14	1.05	0
2	B3A	C	18	2	5,5,6	0.69	0	5,5,7	1.16	0
2	HT7	D	1	2	16,16,17	0.53	0	18,21,23	0.58	0
2	B3D	C	15	2	7,7,9	0.79	0	5,8,11	1.08	0
2	AHP	D	9	2	7,8,9	0.41	0	3,8,10	0.47	0
2	AHP	C	9	2	7,8,9	0.44	0	3,8,10	0.34	0
2	HR7	C	11	2	11,11,12	0.86	1 (9%)	9,12,14	1.42	2 (22%)
2	B3Q	C	8[A]	-	9,9,10	0.67	0	8,10,12	1.04	0
2	B3Q	C	8[B]	-	9,9,10	0.57	0	8,10,12	0.86	0
2	B3D	D	15	2	7,7,9	0.80	0	5,8,11	0.80	0
2	B3E	D	4	3,2	9,9,10	0.83	0	8,10,12	1.18	1 (12%)
2	B3Q	D	8	2	9,9,10	0.63	0	8,10,12	0.95	1 (12%)
2	B3A	D	18	2	5,5,6	0.79	0	5,5,7	1.23	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HT7	C	1	2	-	0/7/7/8	0/2/2/2
2	B3E	C	4	2	-	5/8/8/9	-
2	HR7	D	11	2	-	3/8/10/11	-
2	B3A	C	18	2	-	1/3/3/4	-
2	HT7	D	1	2	-	0/7/7/8	0/2/2/2
2	B3D	C	15	2	-	2/6/6/8	-
2	AHP	D	9	2	-	1/6/7/9	-
2	AHP	C	9	2	-	0/6/7/9	-
2	HR7	C	11	2	-	3/8/10/11	-
2	B3Q	C	8[A]	-	-	1/8/8/9	-
2	B3Q	C	8[B]	-	-	5/8/8/9	-
2	B3D	D	15	2	-	0/6/6/8	-
2	B3E	D	4	3,2	-	1/8/8/9	-
2	B3Q	D	8	2	-	3/8/8/9	-
2	B3A	D	18	2	-	1/3/3/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	11	HR7	CH-NZ	2.23	1.34	1.28

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	11	HR7	O-C-CA	-2.34	118.58	125.38
2	D	18	B3A	CG-CA-CB	-2.14	109.12	111.86
2	D	8	B3Q	O-C-CB	-2.12	119.22	125.38
2	C	11	HR7	CB-CA-C	-2.09	108.80	112.17
2	C	4	B3E	O-C-CB	-2.07	119.37	125.38
2	D	4	B3E	O-C-CB	-2.02	119.49	125.38

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	4	B3E	N-CA-CG-CD

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	4	B3E	CB-CA-CG-CD
2	C	4	B3E	O-C-CB-CA
2	C	15	B3D	CB-CA-CG-CD
2	D	4	B3E	O-C-CB-CA
2	D	11	HR7	CG-CD-CE-NZ
2	C	8[A]	B3Q	O-C-CB-CA
2	C	8[B]	B3Q	O-C-CB-CA
2	D	8	B3Q	O-C-CB-CA
2	C	11	HR7	CE-CD-CG-CB
2	C	8[B]	B3Q	CG-CD-CE-OF1
2	C	8[B]	B3Q	CG-CD-CE-NF2
2	D	11	HR7	CE-CD-CG-CB
2	C	11	HR7	CG-CD-CE-NZ
2	C	15	B3D	N-CA-CG-CD
2	D	8	B3Q	CG-CD-CE-OF1
2	C	18	B3A	O-C-CB-CA
2	D	11	HR7	O-C-CA-CB
2	D	18	B3A	O-C-CB-CA
2	C	8[B]	B3Q	N-CA-CG-CD
2	C	8[B]	B3Q	CB-CA-CG-CD
2	D	8	B3Q	CG-CD-CE-NF2
2	C	4	B3E	CG-CD-CE-OF1
2	C	11	HR7	O-C-CA-CB
2	C	4	B3E	CG-CD-CE-OF2
2	D	9	AHP	C-CA-CB-CG

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	HT7	1	0
2	C	4	B3E	1	0
2	D	9	AHP	2	0
2	C	9	AHP	2	0
2	C	11	HR7	1	0
2	D	4	B3E	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 21 ligands modelled in this entry, 19 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	1205	-	3,3,3	0.44	0	2,2,2	0.39	0
4	EDO	A	1202	-	3,3,3	0.54	0	2,2,2	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	1205	-	-	0/1/1/1	-
4	EDO	A	1202	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1202	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1205	EDO	1	0
4	A	1202	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/157 (88%)	1.58	49 (35%) 1 1	17, 41, 69, 85	17 (12%)
1	B	138/157 (87%)	1.33	42 (30%) 1 1	13, 36, 78, 94	12 (8%)
2	C	13/22 (59%)	1.83	5 (38%) 1 1	36, 48, 71, 86	0
2	D	13/22 (59%)	1.44	2 (15%) 5 5	35, 46, 55, 70	1 (7%)
All	All	303/358 (84%)	1.47	98 (32%) 1 1	13, 41, 73, 94	30 (9%)

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	SER	6.6
1	A	195	TYR	6.5
1	A	113[A]	HIS	6.2
1	A	186	GLY	6.0
1	B	110	SER	5.2
1	B	155	VAL	4.9
1	A	189	ASP	4.8
1	B	108	LEU	4.5
1	A	114	ILE	4.4
1	A	181	TRP	4.3
1	B	107	ASP	4.2
1	A	180	PRO	4.2
1	B	116	PRO	4.0
1	A	117	GLY	4.0
1	A	187	GLY	3.8
1	B	114	ILE	3.7
1	B	117	GLY	3.7
1	B	113	HIS	3.7
1	B	101	TYR	3.6
1	A	183	GLN	3.5
1	A	101	TYR	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176[A]	ASP	3.4
1	B	106[A]	SER	3.4
1	A	184	GLU	3.3
1	A	96	GLU	3.3
1	B	153	GLU	3.3
1	B	100	ARG	3.3
1	B	186	GLY	3.2
1	B	112	LEU	3.2
1	B	156	ASP	3.2
1	A	175	ASN	3.2
1	A	160[A]	GLN	3.1
1	A	190	THR	3.1
1	B	154[A]	SER	3.1
1	B	157	LYS	3.0
2	C	20	TYR	3.0
1	A	194	LEU	3.0
1	B	158	GLU	3.0
1	A	133	ASP	2.9
1	B	193	GLU	2.9
1	A	161	VAL	2.9
1	A	182	ILE	2.9
1	B	159	MET	2.9
1	A	177[A]	HIS	2.9
1	B	109	THR	2.9
2	D	7	ALA	2.8
1	B	195	TYR	2.8
2	C	17	ASN	2.8
1	A	26	GLN	2.8
1	A	132[A]	ARG	2.8
1	A	118	THR	2.8
2	D	3[A]	ARG	2.8
1	A	134	GLY	2.8
1	A	185	ASN	2.7
1	A	125[A]	GLN	2.7
1	A	172	THR	2.7
1	A	103	ARG	2.6
1	B	122	SER	2.6
1	B	118	THR	2.6
1	B	103	ARG	2.6
1	A	193	GLU	2.6
1	B	152	VAL	2.6
1	B	115	THR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	110	SER	2.5
1	B	99	LEU	2.5
1	A	22[A]	TYR	2.5
1	B	105	PHE	2.4
1	A	106	SER	2.4
1	A	100	ARG	2.4
2	C	16	LEU	2.4
1	A	6	ARG	2.4
1	B	111	GLN	2.3
1	B	4[A]	SER	2.3
1	B	184	GLU	2.3
1	A	112	LEU	2.3
1	B	150	LEU	2.3
1	A	1	MET	2.3
1	A	155	VAL	2.3
1	A	128	ASN	2.3
1	B	165	ARG	2.2
2	C	3	ARG	2.2
1	A	20	LYS	2.2
1	A	4	SER	2.2
1	A	99	LEU	2.2
1	A	3	GLN	2.2
1	B	102	ARG	2.2
1	A	104	ALA	2.2
1	B	119	ALA	2.2
2	C	12	MET	2.2
1	A	174	LEU	2.2
1	A	21	GLY	2.1
1	A	120	TYR	2.1
1	B	3	GLN	2.1
1	B	190	THR	2.1
1	B	104	ALA	2.1
1	B	164	SER	2.0
1	A	116	PRO	2.0
1	B	135	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	B3D	C	15	8/10	0.60	0.24	57,64,77,88	0
2	B3D	D	15	8/10	0.71	0.20	54,59,86,90	0
2	B3E	C	4	10/11	0.74	0.20	41,51,78,81	0
2	B3A	D	18	6/7	0.74	0.22	62,69,72,74	0
2	HR7	C	11	12/13	0.78	0.23	50,61,83,84	0
2	HT7	C	1	15/16	0.80	0.17	47,55,62,62	0
2	B3A	C	18	6/7	0.83	0.20	67,75,84,88	0
2	B3E	D	4	10/11	0.84	0.20	50,55,73,75	0
2	B3Q	D	8	10/11	0.86	0.15	47,52,66,72	0
2	HR7	D	11	12/13	0.88	0.16	46,55,70,73	0
2	HT7	D	1	15/16	0.88	0.16	56,69,79,82	0
2	B3Q	C	8[B]	10/11	0.89	0.13	40,46,53,57	7
2	B3Q	C	8[A]	10/11	0.89	0.13	40,46,53,57	7
2	AHP	C	9	9/10	0.91	0.11	34,36,46,46	0
2	AHP	D	9	9/10	0.95	0.12	39,43,56,56	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CD	B	1204[A]	1/1	0.65	0.26	118,118,118,118	1
3	CD	B	1204[B]	1/1	0.65	0.26	120,120,120,120	1
4	EDO	A	1202	4/4	0.78	0.18	39,40,46,57	0
3	CD	B	1203	1/1	0.85	0.23	106,106,106,106	0
3	CD	B	1202	1/1	0.87	0.18	97,97,97,97	0
4	EDO	B	1205	4/4	0.91	0.15	43,50,52,54	0
3	CD	A	1200[A]	1/1	0.92	0.10	62,62,62,62	1
3	CD	A	1200[B]	1/1	0.92	0.10	64,64,64,64	1
3	CD	B	1201	1/1	0.92	0.26	74,74,74,74	0
3	CD	B	1196	1/1	0.96	0.11	47,47,47,47	0
3	CD	B	1199	1/1	0.96	0.25	54,54,54,54	0
3	CD	A	1197	1/1	0.96	0.22	56,56,56,56	0
3	CD	B	1200[B]	1/1	0.97	0.14	88,88,88,88	1
3	CD	B	1197	1/1	0.97	0.23	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CD	A	1198	1/1	0.97	0.34	68,68,68,68	0
3	CD	B	1200[A]	1/1	0.97	0.14	46,46,46,46	1
3	CD	A	1201	1/1	0.98	0.10	53,53,53,53	0
3	CD	B	1198	1/1	0.98	0.21	45,45,45,45	0
3	CD	A	1196[B]	1/1	0.99	0.18	45,45,45,45	1
3	CD	A	1199	1/1	0.99	0.12	46,46,46,46	0
3	CD	A	1196[A]	1/1	0.99	0.18	44,44,44,44	1

6.5 Other polymers [i](#)

There are no such residues in this entry.