



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 05:08 PM UTC

PDB ID : 5BQK / pdb_00005bqk
Title : CRYSTAL STRUCTURE OF C-TERMINAL DOMAIN OF ICP27 PROTEIN FROM HSV-1
Authors : Patel, V.; Rajakannan, V.; Dahlroth, S.; Nordlund, P.
Deposited on : 2015-05-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	3% 88% 10% ..
1	B	271	5% 86% 12% .
1	C	271	6% 82% 16% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6369 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ICP27.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	Se	0	0	0
			2025	1288	362	360	10	5			
1	B	266	Total	C	N	O	S	Se	0	0	0
			2028	1289	362	362	10	5			
1	C	267	Total	C	N	O	S	Se	0	0	0
			2030	1289	360	366	10	5			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		
3	B	92	Total	O	0	0
			92	92		
3	C	76	Total	O	0	0
			76	76		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ICP27



• Molecule 1: ICP27



• Molecule 1: ICP27



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	156.98Å 120.33Å 79.04Å 90.00° 100.46° 90.00°	Depositor
Resolution (Å)	23.80 – 2.00 23.80 – 2.00	Depositor EDS
% Data completeness (in resolution range)	91.5 (23.80-2.00) 91.8 (23.80-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.99Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.217 , 0.249 0.221 , 0.250	Depositor DCC
R_{free} test set	4475 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 30.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6369	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	7/2066 (0.3%)	1.26	12/2800 (0.4%)
1	B	1.32	4/2069 (0.2%)	1.19	8/2803 (0.3%)
1	C	1.26	2/2071 (0.1%)	1.27	9/2807 (0.3%)
All	All	1.34	13/6206 (0.2%)	1.24	29/8410 (0.3%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	333	ALA	CA-CB	8.16	1.66	1.53
1	A	456	VAL	CA-CB	6.62	1.61	1.54
1	A	338	ALA	C-O	-6.50	1.16	1.24
1	B	343	VAL	CA-CB	6.40	1.61	1.54
1	B	462	PHE	CA-C	5.95	1.61	1.52
1	B	354	ALA	CA-CB	5.44	1.61	1.53
1	A	391	ALA	N-CA	5.39	1.52	1.46
1	C	454	VAL	CA-CB	5.39	1.60	1.54
1	C	498	PRO	CA-C	5.17	1.56	1.52
1	A	313	GLU	CG-CD	5.17	1.65	1.52
1	A	359	THR	CA-CB	5.07	1.61	1.53
1	B	324	ARG	N-CA	5.06	1.52	1.46
1	A	310	VAL	C-O	5.04	1.29	1.24

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	379	ILE	N-CA-C	-8.02	103.04	110.82
1	A	498	PRO	N-CA-C	7.62	120.00	110.70
1	A	498	PRO	CA-C-N	-7.35	110.65	119.84
1	A	498	PRO	C-N-CA	-7.35	110.65	119.84
1	C	498	PRO	CA-C-N	-7.35	112.07	119.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	498	PRO	C-N-CA	-7.35	112.07	119.93
1	C	498	PRO	N-CA-C	7.23	119.52	110.70
1	B	498	PRO	CA-C-N	-7.06	112.70	119.90
1	B	498	PRO	C-N-CA	-7.06	112.70	119.90
1	C	386	VAL	CB-CA-C	-7.00	102.56	112.22
1	C	477	ASP	N-CA-C	6.92	118.48	111.07
1	A	371	LEU	CA-C-N	-5.95	114.14	120.03
1	A	371	LEU	C-N-CA	-5.95	114.14	120.03
1	B	386	VAL	CB-CA-C	-5.85	104.36	112.02
1	A	386	VAL	CB-CA-C	-5.78	104.57	111.97
1	B	468	CYS	N-CA-C	-5.76	105.08	111.36
1	A	386	VAL	N-CA-CB	5.53	117.02	110.55
1	B	498	PRO	N-CA-C	5.53	117.45	110.70
1	C	380	ILE	CB-CA-C	-5.52	104.69	112.14
1	B	475	ILE	N-CA-C	5.43	116.20	110.72
1	A	381	GLY	CA-C-N	-5.36	114.15	122.67
1	A	381	GLY	C-N-CA	-5.36	114.15	122.67
1	A	477	ASP	N-CA-C	5.25	116.81	111.14
1	A	283	GLN	N-CA-C	5.18	118.07	109.79
1	A	380	ILE	CB-CA-C	-5.13	105.13	112.22
1	C	510	SER	N-CA-C	-5.10	106.56	112.89
1	B	311	SER	N-CA-C	-5.08	103.69	110.55
1	C	361	ALA	N-CA-C	-5.06	105.84	111.36
1	B	386	VAL	N-CA-C	-5.01	105.82	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2025	0	1992	11	0
1	B	2028	0	1998	20	0
1	C	2030	0	1994	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
3	A	115	0	0	2	0
3	B	92	0	0	1	0
3	C	76	0	0	1	0
All	All	6369	0	5984	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:405:ARG:HG2	1:C:405:ARG:HH11	1.34	0.90
1:C:405:ARG:HG2	1:C:405:ARG:NH1	1.96	0.78
3:B:728:HOH:O	1:C:396:PRO:HG3	1.94	0.68
1:B:254:ARG:HD2	1:C:461:HIS:HD1	1.60	0.67
1:B:254:ARG:NH2	1:C:461:HIS:O	2.29	0.66
1:B:250:ARG:O	1:B:254:ARG:HG3	1.97	0.65
1:A:313:GLU:H	1:A:313:GLU:CD	2.05	0.64
1:A:335:THR:HG22	1:A:339:MSE:HE2	1.80	0.61
1:B:276:MSE:HE1	1:B:339:MSE:HE3	1.82	0.61
1:C:393:ARG:HG2	3:C:726:HOH:O	2.06	0.56
1:B:408:CYS:HB3	1:B:412:GLU:HG2	1.88	0.55
1:A:498:PRO:O	1:A:499:PRO:C	2.46	0.54
1:C:498:PRO:O	1:C:499:PRO:C	2.48	0.54
1:C:498:PRO:HD2	1:C:499:PRO:HD3	1.90	0.54
1:A:280:PHE:CZ	1:A:339:MSE:HE1	2.44	0.53
1:B:502:HIS:HE1	1:B:505:TYR:OH	1.91	0.53
1:C:316:VAL:O	1:C:369:HIS:HE1	1.92	0.52
1:C:276:MSE:HE1	1:C:339:MSE:HE3	1.90	0.52
1:C:494:HIS:HD2	1:C:497:ALA:O	1.94	0.51
1:C:425:ILE:O	1:C:429:VAL:HG23	2.11	0.51
1:B:254:ARG:HD2	1:C:461:HIS:ND1	2.25	0.50
1:A:349:PHE:CE2	1:A:353:LEU:HD11	2.47	0.49
1:C:405:ARG:HH11	1:C:405:ARG:CG	2.13	0.49
1:B:405:ARG:HB2	1:B:407:LEU:HG	1.95	0.49
1:C:265:ILE:O	1:C:268:SER:HB2	2.13	0.48
1:C:479:HIS:O	1:C:482:GLU:HG2	2.13	0.47
1:B:408:CYS:HB2	1:B:413:LEU:HG	1.95	0.47
1:C:364:LYS:HG3	1:C:460:MSE:HB3	1.95	0.47
1:C:422:ILE:O	1:C:423:LYS:HD3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:PHE:CE2	1:A:339:MSE:HE1	2.50	0.46
1:B:254:ARG:HG2	1:C:462:PHE:HA	1.98	0.45
1:A:494:HIS:HE1	3:A:755:HOH:O	2.00	0.45
1:B:489:GLU:O	1:B:489:GLU:HG2	2.16	0.45
1:B:502:HIS:CE1	1:B:505:TYR:OH	2.69	0.45
1:C:279:PRO:HG3	1:C:339:MSE:HE1	1.98	0.45
1:B:303:PHE:CE2	1:B:308:ARG:HG3	2.50	0.45
1:B:405:ARG:HA	1:B:405:ARG:NE	2.32	0.44
1:A:256:ILE:HG23	1:A:257:SER:N	2.32	0.44
1:C:405:ARG:HB3	1:C:407:LEU:HG	2.00	0.44
1:C:276:MSE:O	1:C:279:PRO:HD3	2.17	0.44
1:A:494:HIS:CE1	3:A:755:HOH:O	2.71	0.43
1:A:502:HIS:HE1	1:A:505:TYR:OH	2.03	0.42
1:C:293:ALA:N	1:C:294:PRO:CD	2.82	0.42
1:B:319:GLY:N	1:B:320:PRO:CD	2.83	0.42
1:B:498:PRO:O	1:B:499:PRO:C	2.62	0.42
1:C:401:TYR:CD2	1:C:401:TYR:C	2.98	0.42
1:B:411:ASP:O	1:B:415:SER:HB2	2.20	0.42
1:C:374:ARG:O	1:C:377:ASP:HB3	2.20	0.41
1:B:323:TYR:C	1:B:323:TYR:CD1	2.99	0.41
1:A:347:GLU:O	1:A:393:ARG:NH1	2.54	0.41
1:B:246:ASP:OD2	1:C:423:LYS:NZ	2.48	0.40
1:B:405:ARG:CB	1:B:407:LEU:HG	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/271 (97%)	262 (100%)	1 (0%)	0	100	100
1	B	262/271 (97%)	258 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	263/271 (97%)	255 (97%)	6 (2%)	2 (1%)	16	11
All	All	788/813 (97%)	775 (98%)	11 (1%)	2 (0%)	36	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	288	ALA
1	C	444	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/213 (96%)	204 (100%)	1 (0%)	81	87
1	B	207/213 (97%)	204 (99%)	3 (1%)	59	66
1	C	208/213 (98%)	199 (96%)	9 (4%)	26	25
All	All	620/639 (97%)	607 (98%)	13 (2%)	47	52

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	GLU
1	B	266	SER
1	B	307	THR
1	B	424	ASP
1	C	263	ASP
1	C	283	GLN
1	C	320	PRO
1	C	376	GLN
1	C	403	LYS
1	C	405	ARG
1	C	412	GLU
1	C	424	ASP
1	C	454	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	329	ASN
1	A	369	HIS
1	A	481	GLN
1	A	494	HIS
1	A	502	HIS
1	B	318	HIS
1	B	329	ASN
1	B	481	GLN
1	B	502	HIS
1	C	277	HIS
1	C	369	HIS
1	C	376	GLN
1	C	481	GLN
1	C	494	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/271 (96%)	0.20	7 (2%) 56 55	22, 32, 47, 57	0
1	B	261/271 (96%)	0.44	13 (4%) 34 33	26, 35, 53, 64	0
1	C	262/271 (96%)	0.68	16 (6%) 27 26	26, 40, 55, 62	0
All	All	785/813 (96%)	0.44	36 (4%) 37 36	22, 35, 53, 64	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	289	ASN	4.3
1	C	287	ALA	3.9
1	C	297	ALA	3.6
1	B	302	PRO	3.4
1	B	281	GLY	3.4
1	C	291	PRO	3.3
1	B	269	PHE	3.2
1	B	297	ALA	3.2
1	B	304	ASP	3.2
1	C	242	ALA	3.1
1	A	269	PHE	3.1
1	A	297	ALA	2.9
1	C	288	ALA	2.9
1	B	289	ASN	2.9
1	C	302	PRO	2.9
1	C	290	SER	2.8
1	C	442	ARG	2.7
1	A	242	ALA	2.7
1	C	269	PHE	2.6
1	B	408	CYS	2.6
1	B	282	GLY	2.6
1	A	282	GLY	2.5
1	C	367	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	307	THR	2.4
1	C	404	ALA	2.4
1	B	303	PHE	2.4
1	A	287	ALA	2.3
1	C	461	HIS	2.3
1	A	288	ALA	2.3
1	C	274	GLN	2.2
1	A	302	PRO	2.1
1	C	455	GLY	2.1
1	B	262	VAL	2.1
1	B	306	GLU	2.0
1	C	334	SER	2.1
1	B	245	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	601	1/1	0.99	0.03	35,35,35,35	0
2	ZN	A	601	1/1	1.00	0.02	33,33,33,33	0
2	ZN	C	601	1/1	1.00	0.01	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.