



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:45 AM UTC

PDB ID : 5BTN / pdb_00005btn
Title : Crystal structure of a topoisomerase II complex
Authors : Blower, T.R.; Williamson, B.H.; Kerns, R.J.; Berger, J.M.
Deposited on : 2015-06-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	8% 93% 5% •
1	C	503	6% 91% 5% •
2	B	253	11% 92% 5% •
2	D	253	11% 90% 8% •
3	E	24	38% 58% 29% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	24	
4	F	24	
4	G	24	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	PTR	A	129	-	-	X	-
1	PTR	C	129	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 26010 atoms, of which 12525 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit A.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	487	Total	C	H	N	O	P	S	0	1	0
			7693	2388	3854	704	733	1	13			
1	C	487	Total	C	H	N	O	P	S	0	1	0
			7690	2388	3851	704	733	1	13			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	SER	ALA	engineered mutation	UNP P9WG47
A	501	ILE	-	expression tag	UNP P9WG47
A	502	GLY	-	expression tag	UNP P9WG47
A	503	SER	-	expression tag	UNP P9WG47
A	504	GLY	-	expression tag	UNP P9WG47
C	90	SER	ALA	engineered mutation	UNP P9WG47
C	501	ILE	-	expression tag	UNP P9WG47
C	502	GLY	-	expression tag	UNP P9WG47
C	503	SER	-	expression tag	UNP P9WG47
C	504	GLY	-	expression tag	UNP P9WG47

- Molecule 2 is a protein called DNA gyrase subunit B.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	245	Total	C	H	N	O	S	0	0	0
			3906	1217	1974	348	360	7			
2	D	247	Total	C	H	N	O	S	0	0	0
			3930	1224	1984	351	364	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	423	SER	-	expression tag	UNP P9WG45
B	424	ASN	-	expression tag	UNP P9WG45

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	425	ALA	-	expression tag	UNP P9WG45
D	423	SER	-	expression tag	UNP P9WG45
D	424	ASN	-	expression tag	UNP P9WG45
D	425	ALA	-	expression tag	UNP P9WG45

- Molecule 3 is a DNA chain called DNA substrate 24-mer GGTCATGAATGACTATGCAC GTAA.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	21	Total	C	H	N	O	P	0	21	0
			620	196	210	74	120	20			
3	H	21	Total	C	H	N	O	P	0	21	0
			616	196	206	74	120	20			


- Molecule 4 is a DNA chain called DNA substrate 24-mer TTACGTGCATAGTCATTCAT GACC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	F	21	Total	C	H	N	O	P	0	21	0
			612	196	202	74	120	20			
4	G	21	Total	C	H	N	O	P	0	21	0
			608	196	198	74	120	20			

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		
5	E	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 1-cyclopropyl-6-fluoro-8-methyl-7-[(4aS,7aS)-octahydro-6H-pyrrolo[3,4-b]pyridin-6-yl]-4-oxo-1,4-dihydroquinoline-3-carboxylic acid (CCD ID: 8MX) (formula: C₂₁H₂₄FN₃O₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	G	1	Total	C	F	H	N	O	0	0
			51	21	1	23	3	3		
6	H	1	Total	C	F	H	N	O	0	0
			51	21	1	23	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	100	Total	O	0	0
			100	100		
7	B	31	Total	O	0	0
			31	31		
7	C	41	Total	O	0	0
			41	41		
7	D	29	Total	O	0	0
			29	29		
7	E	6	Total	O	0	0
			6	6		
7	F	5	Total	O	0	0
			5	5		
7	G	7	Total	O	0	0
			7	7		
7	H	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA gyrase subunit A



• Molecule 1: DNA gyrase subunit A




• Molecule 2: DNA gyrase subunit B





• Molecule 2: DNA gyrase subunit B







- Molecule 3: DNA substrate 24-mer GGTCATGAATGACTATGCACGTAA




- Molecule 3: DNA substrate 24-mer GGTCATGAATGACTATGCACGTAA



- Molecule 4: DNA substrate 24-mer TTACGTGCATAGTCATTCATGACC



- Molecule 4: DNA substrate 24-mer TTACGTGCATAGTCATTCATGACC



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	108.34Å 82.91Å 129.08Å 90.00° 108.78° 90.00°	Depositor
Resolution (Å)	49.65 – 2.50 49.65 – 2.50	Depositor EDS
% Data completeness (in resolution range)	95.6 (49.65-2.50) 94.3 (49.65-2.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.51Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.223 , 0.251 0.231 , 0.254	Depositor DCC
R_{free} test set	3566 reflections (4.18%)	wwPDB-VP
Wilson B-factor (Å ²)	47.7	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	26010	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 8MX, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/3882	0.66	2/5254 (0.0%)
1	C	0.30	0/3882	0.64	1/5254 (0.0%)
2	B	0.29	0/1960	0.60	0/2634
2	D	0.29	0/1974	0.60	0/2653
3	E	0.20	0/458	0.47	0/704
3	H	0.19	0/458	0.47	0/704
4	F	0.18	0/458	0.45	0/704
4	G	0.18	0/458	0.45	0/704
All	All	0.28	0/13530	0.61	3/18611 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	293	ASP	N-CA-C	-6.82	104.23	112.54
1	C	293	ASP	N-CA-C	-5.46	104.72	111.33
1	A	292	ARG	N-CA-C	5.12	116.55	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3839	3854	3853	24	0
1	C	3839	3851	3853	23	0
2	B	1932	1974	1973	7	0
2	D	1946	1984	1984	14	0
3	E	410	210	210	11	0
3	H	410	206	206	5	0
4	F	410	202	202	7	0
4	G	410	198	198	7	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
6	G	28	23	0	1	0
6	H	28	23	0	2	0
7	A	100	0	0	2	0
7	B	31	0	0	0	0
7	C	41	0	0	0	0
7	D	29	0	0	0	0
7	E	6	0	0	1	0
7	F	5	0	0	0	0
7	G	7	0	0	0	0
7	H	10	0	0	0	0
All	All	13485	12525	12479	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:GLU:OE1	1:A:267:THR:HG22	1.59	1.02
1:C:129:PTR:O3P	4:G:11[B]:DA:C5'	2.12	0.97
1:A:261:GLU:OE1	1:A:267:THR:CG2	2.17	0.92
1:C:129:PTR:O3P	3:E:11[A]:DG:H4'	1.73	0.88
1:A:292:ARG:NH2	2:D:431:SER:O	2.08	0.85
1:C:98:ARG:NH1	4:F:7[A]:DG:OP1	2.12	0.81
3:E:21[A]:DG:N2	4:F:4[A]:DC:O2	2.12	0.80
2:D:653:GLY:O	2:D:659:ARG:NH2	2.14	0.80
1:A:129:PTR:O2P	3:H:11[B]:DG:H4'	1.85	0.77
1:A:129:PTR:O2P	4:F:11[A]:DA:C5'	2.32	0.77
1:C:129:PTR:O3P	4:G:11[B]:DA:C4'	2.34	0.75
4:G:7[B]:DG:N2	3:H:18[B]:DC:O2	2.20	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:129:PTR:O3P	3:E:11[A]:DG:C4'	2.34	0.74
1:A:129:PTR:O2P	3:H:11[B]:DG:C5'	2.37	0.73
1:C:129:PTR:O3P	3:E:11[A]:DG:C5'	2.37	0.72
1:C:129:PTR:O3P	4:G:11[B]:DA:H4'	1.88	0.72
1:C:357:ARG:NH1	1:C:359:ASP:OD1	2.26	0.68
2:D:429:ARG:NH1	2:D:439:PRO:O	2.28	0.67
2:D:659:ARG:NH1	3:E:19[A]:DA:OP1	2.28	0.66
1:A:129:PTR:O2P	3:H:11[B]:DG:C4'	2.44	0.66
1:A:129:PTR:O2P	4:F:11[A]:DA:H4'	1.96	0.65
2:B:429:ARG:NH1	2:B:439:PRO:O	2.30	0.65
1:C:262:ASP:OD1	1:C:263:SER:N	2.37	0.58
1:A:441:ARG:NE	1:C:437:ASP:OD1	2.38	0.56
1:A:98:ARG:NH2	3:E:7[A]:DG:OP1	2.38	0.56
1:A:261:GLU:OE1	1:A:267:THR:HG21	2.05	0.56
1:A:261:GLU:HG2	7:A:696:HOH:O	2.07	0.55
1:C:389:HIS:O	1:C:431:GLN:NE2	2.40	0.54
1:A:98:ARG:CZ	3:E:7[A]:DG:OP1	2.58	0.51
1:C:31:TYR:HA	2:D:535:VAL:CG1	2.40	0.51
4:G:5[B]:DG:N2	3:H:20[B]:DC:O2	2.43	0.51
1:A:98:ARG:NH1	3:E:7[A]:DG:OP1	2.44	0.50
6:H:101:8MX:C01	6:H:101:8MX:C09	2.90	0.50
2:B:446:ARG:NH1	2:B:473:SER:OG	2.45	0.50
2:D:484:LYS:NZ	2:D:536:ASP:O	2.45	0.49
1:A:226:PRO:O	1:A:494:ARG:NH2	2.46	0.49
1:C:428:ASP:OD1	1:C:429:GLU:N	2.43	0.49
3:E:12[A]:DA:OP1	7:E:201:HOH:O	2.16	0.48
1:A:129:PTR:O2P	4:F:11[A]:DA:C4'	2.51	0.46
1:A:427:ILE:CG2	1:A:431:GLN:HB2	2.45	0.46
1:A:23:GLU:OE1	2:B:634:ARG:NH1	2.49	0.45
1:C:60:PHE:HB2	1:C:142:LEU:HD13	1.99	0.45
1:A:51:VAL:HG11	3:E:8[A]:DA:H5'	1.99	0.45
1:A:275:PRO:HG2	1:A:278:VAL:HG21	1.99	0.45
1:A:15:ILE:HD12	2:B:633:LEU:HB2	1.99	0.44
1:C:275:PRO:HG2	1:C:278:VAL:HG21	1.99	0.44
4:F:16[A]:DT:C6	4:F:17[A]:DT:H72	2.53	0.44
1:C:31:TYR:HA	2:D:535:VAL:HG13	2.00	0.43
1:C:186:ALA:O	1:C:344:ASN:ND2	2.52	0.43
1:C:343:ALA:HB1	1:C:345:MET:HE3	1.99	0.43
2:D:446:ARG:NH2	2:D:473:SER:OG	2.51	0.43
2:D:627:ASP:HB3	2:D:630:VAL:HG22	1.99	0.43
1:A:98:ARG:NH1	4:G:7[B]:DG:OP1	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:GLU:OE1	2:D:634:ARG:NH1	2.51	0.43
1:C:34:SER:OG	2:D:535:VAL:HG11	2.19	0.43
2:B:511:THR:HG21	2:B:517:PHE:CE2	2.53	0.43
6:H:101:8MX:C01	6:H:101:8MX:C23	2.97	0.43
6:G:101:8MX:C23	6:G:101:8MX:C01	2.97	0.42
2:B:553:ARG:N	2:B:554:PRO:CD	2.83	0.42
2:D:497:LEU:O	2:D:503:GLN:NE2	2.51	0.42
3:E:20[A]:DC:O2	4:F:5[A]:DG:N2	2.51	0.42
1:C:65:ARG:HB3	1:C:66:PRO:HD2	2.01	0.41
2:D:553:ARG:N	2:D:554:PRO:CD	2.83	0.41
4:G:16[B]:DT:C6	4:G:17[B]:DT:H72	2.55	0.41
1:A:402:ASP:OD1	1:C:409:ARG:NH1	2.52	0.41
2:B:601:ILE:HG22	2:B:606:GLY:HA2	2.03	0.41
1:A:261:GLU:CG	7:A:696:HOH:O	2.65	0.41
1:C:292:ARG:O	1:C:293:ASP:C	2.62	0.41
2:D:511:THR:HG21	2:D:517:PHE:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/503 (96%)	467 (96%)	18 (4%)	0	100	100
1	C	485/503 (96%)	468 (96%)	17 (4%)	0	100	100
2	B	241/253 (95%)	232 (96%)	9 (4%)	0	100	100
2	D	243/253 (96%)	235 (97%)	8 (3%)	0	100	100
All	All	1454/1512 (96%)	1402 (96%)	52 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/421 (97%)	407 (100%)	1 (0%)	87	95
1	C	408/421 (97%)	407 (100%)	1 (0%)	87	95
2	B	205/211 (97%)	205 (100%)	0	100	100
2	D	207/211 (98%)	207 (100%)	0	100	100
All	All	1228/1264 (97%)	1226 (100%)	2 (0%)	87	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	422	ILE
1	C	422	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	GLN
1	A	207	ASN
1	A	277	GLN
1	A	449	GLN
2	B	514	HIS
1	C	161	GLN
1	C	207	ASN
1	C	238	GLN
1	C	277	GLN
1	C	344	ASN
1	C	449	GLN
2	D	525	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	129	1	15,16,17	0.78	0	17,22,24	0.75	0
1	PTR	C	129	1	15,16,17	0.78	0	17,22,24	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	129	1	-	0/10/11/13	0/1/1/1
1	PTR	C	129	1	-	1/10/11/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	129	PTR	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	129	PTR	6	0
1	C	129	PTR	6	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	8MX	G	101	5	32,32,32	5.07	12 (37%)	41,49,49	2.68	16 (39%)
6	8MX	H	101	5	32,32,32	5.13	9 (28%)	41,49,49	2.95	20 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	8MX	G	101	5	-	6/12/33/33	1/5/5/5
6	8MX	H	101	5	-	2/12/33/33	0/5/5/5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	H	101	8MX	C16-C15	-19.61	1.20	1.52
6	G	101	8MX	C16-C15	-17.72	1.23	1.52
6	H	101	8MX	C09-C10	-16.73	1.28	1.53
6	G	101	8MX	C09-C10	-15.83	1.30	1.53
6	G	101	8MX	C16-N08	10.03	1.62	1.46
6	G	101	8MX	C10-C15	8.47	1.67	1.53
6	H	101	8MX	C16-N08	7.90	1.58	1.46
6	H	101	8MX	C10-C15	6.68	1.64	1.53
6	G	101	8MX	C12-C13	3.96	1.68	1.49
6	H	101	8MX	C12-C13	3.34	1.65	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	101	8MX	C04-C18	-3.28	1.41	1.48
6	H	101	8MX	C04-C18	-3.17	1.42	1.48
6	H	101	8MX	C03-C02	2.89	1.46	1.40
6	G	101	8MX	C03-C02	2.76	1.45	1.40
6	H	101	8MX	C12-C11	-2.61	1.46	1.53
6	G	101	8MX	C09-N08	2.43	1.50	1.46
6	G	101	8MX	C15-N14	2.40	1.51	1.47
6	G	101	8MX	C20-C18	-2.34	1.39	1.44
6	H	101	8MX	C20-C18	-2.25	1.39	1.44
6	G	101	8MX	C12-C11	-2.22	1.47	1.53
6	G	101	8MX	C07-N08	2.14	1.47	1.40

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	101	8MX	C24-C23-N22	9.17	133.37	118.87
6	G	101	8MX	C24-C23-N22	8.76	132.72	118.87
6	H	101	8MX	C16-N08-C09	-6.60	102.06	111.64
6	H	101	8MX	C09-N08-C07	-6.23	113.67	123.36
6	H	101	8MX	C13-N14-C15	-5.78	108.02	111.62
6	G	101	8MX	C23-N22-C21	-5.72	111.36	119.74
6	H	101	8MX	C23-N22-C21	-5.27	112.02	119.74
6	G	101	8MX	C09-N08-C07	-5.24	115.21	123.36
6	G	101	8MX	C16-N08-C07	-4.68	116.08	123.36
6	G	101	8MX	C25-C23-N22	4.36	125.76	118.87
6	H	101	8MX	C03-N22-C21	3.63	122.76	119.82
6	G	101	8MX	C03-N22-C23	3.61	126.13	122.35
6	G	101	8MX	C13-N14-C15	-3.55	109.40	111.62
6	G	101	8MX	C03-N22-C21	3.32	122.50	119.82
6	H	101	8MX	C01-C02-C07	-3.12	115.12	121.20
6	G	101	8MX	C04-C03-N22	-3.03	115.25	119.00
6	H	101	8MX	C16-N08-C07	-3.01	118.67	123.36
6	H	101	8MX	C04-C03-N22	-3.01	115.28	119.00
6	H	101	8MX	C25-C23-N22	2.97	123.57	118.87
6	H	101	8MX	F17-C06-C07	2.90	123.05	118.28
6	G	101	8MX	C05-C04-C18	-2.79	115.25	119.94
6	H	101	8MX	C03-N22-C23	2.71	125.19	122.35
6	G	101	8MX	C09-C10-C15	2.68	108.61	103.97
6	G	101	8MX	C06-C07-N08	-2.68	114.38	121.93
6	H	101	8MX	C05-C04-C18	-2.42	115.88	119.94
6	G	101	8MX	C16-N08-C09	-2.38	108.18	111.64
6	H	101	8MX	C04-C03-C02	-2.34	117.91	120.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	101	8MX	C20-C21-N22	-2.20	121.78	124.42
6	G	101	8MX	C04-C03-C02	-2.19	118.09	120.78
6	H	101	8MX	O27-C26-C20	2.14	120.70	115.69
6	H	101	8MX	C05-C06-C07	-2.10	120.22	123.34
6	G	101	8MX	C04-C18-C20	2.09	118.28	115.64
6	H	101	8MX	C04-C18-C20	2.09	118.28	115.64
6	H	101	8MX	C12-C13-N14	-2.03	105.65	110.84
6	G	101	8MX	C01-C02-C07	-2.01	117.29	121.20
6	H	101	8MX	O28-C26-C20	-2.00	118.53	122.67

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	G	101	8MX	C06-C07-N08-C09
6	H	101	8MX	C02-C07-N08-C16
6	G	101	8MX	C18-C20-C26-O28
6	G	101	8MX	C18-C20-C26-O27
6	G	101	8MX	C21-C20-C26-O27
6	G	101	8MX	C21-C20-C26-O28
6	G	101	8MX	C06-C07-N08-C16
6	H	101	8MX	C06-C07-N08-C09

All (1) ring outliers are listed below:



Mol	Chain	Res	Type	Atoms
6	G	101	8MX	C10-C11-C12-C13-C15-N14

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	G	101	8MX	1	0
6	H	101	8MX	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	486/503 (96%)	0.58	39 (8%)	18	16	25, 61, 116, 185	1 (0%)
1	C	486/503 (96%)	0.63	28 (5%)	29	25	31, 64, 115, 186	1 (0%)
2	B	245/253 (96%)	0.76	27 (11%)	10	8	47, 71, 105, 142	0
2	D	247/253 (97%)	0.76	27 (10%)	10	9	42, 67, 104, 161	0
3	E	21/24 (87%)	1.78	9 (42%)	0	0	29, 52, 97, 106	21 (100%)
3	H	21/24 (87%)	0.92	4 (19%)	3	2	15, 31, 68, 91	21 (100%)
4	F	21/24 (87%)	1.88	8 (38%)	1	0	33, 46, 94, 100	21 (100%)
4	G	21/24 (87%)	0.66	1 (4%)	35	31	14, 27, 76, 91	21 (100%)
All	All	1548/1608 (96%)	0.69	143 (9%)	14	13	14, 64, 114, 186	86 (5%)

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	501	ILE	8.0
2	D	426	LEU	6.4
1	A	501	ILE	6.0
2	D	431	SER	5.2
1	A	293	ASP	4.8
2	B	437	GLY	4.6
2	B	427	VAL	4.6
2	B	426	LEU	4.5
2	D	437	GLY	4.4
1	A	297	ALA	4.3
2	D	605	ASP	4.3
2	B	425	ALA	4.2
2	B	430	LYS	4.1
1	A	414	VAL	4.1
1	A	500	ALA	4.0
2	D	601	ILE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	412	GLU	3.9
3	E	22[A]	DT	3.9
1	C	500	ALA	3.8
2	B	601	ILE	3.8
2	D	484	LYS	3.8
1	A	408	ILE	3.8
2	B	605	ASP	3.8
2	D	675	VAL	3.7
3	H	3[B]	DT	3.7
1	A	387	ARG	3.5
2	D	602	ASN	3.5
2	D	609	ARG	3.4
2	D	674	ASP	3.4
1	A	413	THR	3.3
2	B	606	GLY	3.2
1	A	261	GLU	3.2
1	A	406	ALA	3.2
2	D	424	ASN	3.2
2	B	609	ARG	3.1
2	D	533	ALA	3.1
1	C	383	LYS	3.1
3	E	3[A]	DT	3.1
1	A	296	LEU	3.1
1	C	209	ASP	3.0
2	D	606	GLY	3.0
2	B	533	ALA	3.0
3	E	6[A]	DT	3.0
1	A	458	ILE	2.9
2	B	511	THR	2.9
1	A	410	ALA	2.9
3	E	11[A]	DG	2.9
4	F	5[A]	DG	2.9
1	C	387	ARG	2.9
1	A	436	LEU	2.9
2	D	494	ASP	2.9
1	A	441	ARG	2.9
4	F	21[A]	DG	2.9
4	F	6[A]	DT	2.9
2	B	602	ASN	2.9
1	C	458	ILE	2.9
3	H	11[B]	DG	2.9
4	G	23[B]	DC	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	F	22[A]	DA	2.9
2	D	427	VAL	2.8
1	C	267	THR	2.8
1	C	297	ALA	2.8
2	D	565	GLN	2.8
3	E	5[A]	DA	2.8
1	C	296	LEU	2.8
1	C	290	GLN	2.8
1	A	427	ILE	2.8
1	A	263	SER	2.8
1	C	16	GLU	2.8
1	C	412	GLU	2.8
1	A	15	ILE	2.7
2	B	674	ASP	2.7
1	A	409	ARG	2.7
1	A	122	ASP	2.7
3	E	4[A]	DC	2.7
2	D	511	THR	2.7
1	A	430	ILE	2.7
1	C	408	ILE	2.7
1	A	67	ASP	2.6
2	B	675	VAL	2.6
3	E	23[A]	DA	2.6
4	F	4[A]	DC	2.6
3	H	21[B]	DG	2.6
1	C	15	ILE	2.6
1	C	427	ILE	2.6
2	D	493	ILE	2.6
4	F	8[A]	DC	2.6
2	B	493	ILE	2.5
4	F	15[A]	DA	2.5
1	C	442	ARG	2.5
2	D	607	ILE	2.5
1	A	440	LEU	2.5
1	A	262	ASP	2.5
1	C	293	ASP	2.5
1	C	422	ILE	2.5
1	A	292	ARG	2.5
2	D	664	THR	2.5
2	B	494	ASP	2.4
1	A	266	ARG	2.4
1	C	292	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	450	ARG	2.4
3	E	13[A]	DC	2.4
2	D	668	LYS	2.3
1	A	404	VAL	2.3
1	C	404	VAL	2.3
2	B	565	GLN	2.3
1	C	424	LEU	2.3
2	B	671	ARG	2.3
1	A	405	ILE	2.3
2	B	603	LYS	2.3
2	B	604	GLU	2.3
1	A	265	GLY	2.2
1	A	431	GLN	2.2
1	A	428	ASP	2.2
3	E	12[A]	DA	2.2
2	D	671	ARG	2.2
1	C	259	VAL	2.2
2	B	664	THR	2.2
2	B	485	ILE	2.2
1	C	258	GLU	2.2
1	A	68	ARG	2.2
2	B	585	ARG	2.2
4	F	3[A]	DA	2.2
2	B	607	ILE	2.2
1	A	246	THR	2.1
1	C	265	GLY	2.1
1	C	321	ASP	2.1
2	D	603	LYS	2.1
2	D	425	ALA	2.1
2	D	438	LEU	2.1
1	C	418	ARG	2.1
2	B	492	ARG	2.1
1	A	400	ALA	2.1
1	A	401	LEU	2.1
1	C	264	ARG	2.1
1	A	264	ARG	2.1
1	A	298	GLY	2.1
2	B	495	ARG	2.1
2	B	668	LYS	2.1
1	A	403	GLU	2.0
2	D	654	GLU	2.0
2	D	592	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	4[B]	DC	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PTR	C	129	16/17	0.94	0.11	47,51,74,78	0
1	PTR	A	129	16/17	0.96	0.09	43,48,68,70	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	8MX	G	101	28/28	0.86	0.22	56,78,113,113	0
6	8MX	H	101	28/28	0.90	0.16	47,65,141,141	0
5	MG	E	101	1/1	0.95	0.08	74,74,74,74	0
5	MG	C	601	1/1	0.96	0.05	46,46,46,46	0
5	MG	B	701	1/1	0.96	0.06	37,37,37,37	0
5	MG	D	701	1/1	0.97	0.05	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 8MX G 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 8MX H 101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.