



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 11:36 AM EDT

PDB ID : 8BTL / pdb_00008btl
Title : Crystal structure of a complex between the E2 conjugating enzyme UBE2A and the E3 ligase module from UBR4
Authors : Virdee, S.; Mabbitt, P.D.; Barnsby-Greer, L.
Deposited on : 2022-11-29
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	459	
2	C	158	
2	D	158	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5187 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cDNA FLJ12511 fis, clone NT2RM2001727, highly similar to Homo sapiens ubiquitin protein ligase E3 component n-recogin 4 (UBR4), mRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2795	1794	468	521	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4725	GLY	-	expression tag	UNP B3KMT2
A	4726	PRO	-	expression tag	UNP B3KMT2
A	4727	LEU	-	expression tag	UNP B3KMT2
A	4729	SER	LEU	conflict	UNP B3KMT2

- Molecule 2 is a protein called Ubiquitin conjugating enzyme E2 A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	153	Total	C	N	O	S	0	0	0
			1211	762	207	238	4			
2	D	150	Total	C	N	O	S	0	0	0
			1180	746	200	231	3			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-5	GLY	-	expression tag	UNP A0A7N4PN18
C	-4	HIS	-	expression tag	UNP A0A7N4PN18
C	-3	MET	VAL	conflict	UNP A0A7N4PN18
C	0	SER	ASP	conflict	UNP A0A7N4PN18
C	88	LYS	CYS	conflict	UNP A0A7N4PN18
D	-5	GLY	-	expression tag	UNP A0A7N4PN18
D	-4	HIS	-	expression tag	UNP A0A7N4PN18
D	-3	MET	VAL	conflict	UNP A0A7N4PN18
D	0	SER	ASP	conflict	UNP A0A7N4PN18

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	88	LYS	CYS	conflict	UNP A0A7N4PN18


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cDNA FLJ12511 fis, clone NT2RM2001727, highly similar to Homo sapiens ubiquitin protein ligase E3 component n-recognin 4 (UBR4), mRNA



- Molecule 2: Ubiquitin conjugating enzyme E2 A



- Molecule 2: Ubiquitin conjugating enzyme E2 A



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.16Å 89.16Å 263.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.83 – 3.20 65.83 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (65.83-3.20) 99.6 (65.83-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 3.19Å)	Xtriage
Refinement program	PHENIX 1.17.1	Depositor
R, R_{free}	0.228 , 0.263 0.232 , 0.255	Depositor DCC
R_{free} test set	972 reflections (5.29%)	wwPDB-VP
Wilson B-factor (Å ²)	84.0	Xtriage
Anisotropy	0.538	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5187	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/2866	0.26	0/3900
2	C	0.09	0/1243	0.27	0/1694
2	D	0.08	0/1212	0.25	0/1655
All	All	0.09	0/5321	0.26	0/7249

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2795	0	2720	21	0
2	C	1211	0	1154	14	0
2	D	1180	0	1115	12	0
3	A	1	0	0	0	0
All	All	5187	0	4989	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:80:ASN:HD21	2:D:117:ASN:HB2	1.57	0.69
2:D:88:LYS:HA	2:D:93:GLN:HE22	1.67	0.58
1:A:5047:PRO:HD2	1:A:5050:GLN:HG3	1.85	0.57
2:D:89:LEU:HD13	2:D:112:LEU:HD22	1.89	0.55
2:C:8:ARG:HE	2:C:64:PRO:HG3	1.73	0.54
1:A:5054:THR:HG22	1:A:5058:ILE:HG13	1.89	0.54
2:C:6:ARG:HA	2:C:9:LEU:HB2	1.90	0.54
1:A:5042:ALA:HB1	1:A:5046:LEU:HD12	1.92	0.52
2:D:43:PRO:O	2:D:46:THR:OG1	2.27	0.51
1:A:4833:GLU:HB3	1:A:4837:THR:HG22	1.93	0.51
1:A:4884:ASN:ND2	1:A:4911:ASN:OD1	2.36	0.51
2:D:28:PRO:HG3	2:D:36:TRP:CE2	2.46	0.51
2:C:39:VAL:HG22	2:C:54:LYS:HG2	1.93	0.50
1:A:4930:SER:HB2	2:D:93:GLN:HE21	1.77	0.50
1:A:4965:PHE:HE1	1:A:4990:ILE:HD12	1.77	0.50
1:A:4868:GLU:O	1:A:4873:LYS:NZ	2.35	0.49
2:D:8:ARG:NH2	2:D:100:TYR:O	2.46	0.49
1:A:5112:THR:HG23	1:A:5113:SER:H	1.78	0.48
2:C:42:GLY:HA3	2:C:48:PHE:O	2.14	0.48
2:C:57:ILE:HG23	2:C:70:VAL:HG22	1.95	0.47
2:D:29:SER:OG	2:D:32:ASN:O	2.32	0.47
1:A:5110:VAL:HG22	1:A:5126:TYR:CG	2.49	0.47
1:A:5159:LEU:HD12	1:A:5171:PRO:HB3	1.97	0.47
2:C:46:THR:HG21	2:C:114:ASP:HA	1.97	0.47
2:D:79:PRO:HB3	2:D:127:ALA:HB2	1.97	0.47
1:A:4987:ILE:HG12	1:A:5142:LEU:HB2	1.97	0.47
2:C:6:ARG:NH1	2:C:61:GLU:OE2	2.48	0.46
2:C:8:ARG:HH12	2:C:101:ASP:HB3	1.81	0.46
1:A:4901:ARG:HH21	1:A:4906:SER:HA	1.82	0.44
1:A:4957:ASN:O	1:A:4961:ILE:HG13	2.17	0.44
1:A:5079:LEU:HD23	1:A:5079:LEU:HA	1.83	0.43
1:A:5028:SER:OG	1:A:5061:ARG:NH2	2.52	0.43
2:D:42:GLY:HA3	2:D:48:PHE:O	2.18	0.42
1:A:4882:HIS:HD2	1:A:4914:THR:HG21	1.84	0.42
1:A:4842:ARG:HH21	2:C:101:ASP:HB3	1.85	0.42
2:D:63:TYR:CD1	2:D:64:PRO:HA	2.55	0.41
2:D:88:LYS:HA	2:D:93:GLN:NE2	2.32	0.41
2:C:91:ILE:HG23	2:C:100:TYR:CE2	2.56	0.41
1:A:4901:ARG:NH2	1:A:4906:SER:HA	2.36	0.41
2:C:5:ALA:O	2:C:6:ARG:HB2	2.21	0.41
2:C:92:LEU:HD21	2:C:105:ILE:HG23	2.03	0.41
1:A:5055:ARG:HB3	1:A:5097:TRP:HZ2	1.86	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:28:PRO:HG3	2:C:36:TRP:CE2	2.56	0.40
2:C:94:ASN:OD1	2:C:95:ARG:HG2	2.20	0.40
1:A:5048:PRO:O	1:A:5052:ARG:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/459 (77%)	336 (96%)	16 (4%)	0	100	100
2	C	151/158 (96%)	144 (95%)	7 (5%)	0	100	100
2	D	148/158 (94%)	143 (97%)	5 (3%)	0	100	100
All	All	651/775 (84%)	623 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/393 (76%)	298 (100%)	0	100	100
2	C	134/141 (95%)	134 (100%)	0	100	100
2	D	129/141 (92%)	129 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	561/675 (83%)	561 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4995	HIS
1	A	5002	ASN
2	C	110	GLN
2	D	80	ASN
2	D	93	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/459 (77%)	-0.02	6 (1%) 69 49	55, 74, 101, 142	0
2	C	153/158 (96%)	-0.05	0 100 100	63, 90, 119, 130	0
2	D	150/158 (94%)	-0.16	0 100 100	66, 81, 105, 116	0
All	All	657/775 (84%)	-0.06	6 (0%) 81 64	55, 79, 111, 142	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5112	THR	2.5
1	A	5071	VAL	2.3
1	A	4888	TYR	2.3
1	A	5116	GLU	2.3
1	A	5181	SER	2.1
1	A	5114	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	$Q < 0.9$
3	ZN	A	5201	1/1	1.00	0.04	63,63,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.