



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 03:04 PM UTC

PDB ID : 4BUO / pdb_00004buo
Title : High Resolution Structure of Thermostable Agonist-bound Neurotensin Receptor 1 Mutant without Lysozyme Fusion
Authors : Egloff, P.; Hillenbrand, M.; Schlinkmann, K.M.; Batyuk, A.; Mittl, P.; Plueckthun, A.
Deposited on : 2013-06-21
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	
1	B	335	
2	C	10	
2	D	10	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLY	B	1389	-	X	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEUROTENSIN RECEPTOR TYPE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	0	0
			2400	1589	387	409	15			
1	B	314	Total	C	N	O	S	0	0	0
			2463	1628	394	424	17			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	PRO	-	expression tag	UNP P20789
A	48	GLY	-	expression tag	UNP P20789
A	49	SER	-	expression tag	UNP P20789
A	391	THR	-	expression tag	UNP P20789
A	392	ARG	-	expression tag	UNP P20789
A	393	GLU	-	expression tag	UNP P20789
A	394	LEU	-	expression tag	UNP P20789
A	395	GLU	-	expression tag	UNP P20789
A	396	VAL	-	expression tag	UNP P20789
A	397	LEU	-	expression tag	UNP P20789
A	398	PHE	-	expression tag	UNP P20789
A	399	GLN	-	expression tag	UNP P20789
A	86	LEU	ALA	engineered mutation	UNP P20789
A	103	ASP	HIS	engineered mutation	UNP P20789
A	105	TYR	HIS	engineered mutation	UNP P20789
A	161	VAL	ALA	engineered mutation	UNP P20789
A	167	LEU	ARG	engineered mutation	UNP P20789
A	213	LEU	ARG	engineered mutation	UNP P20789
A	234	LEU	VAL	engineered mutation	UNP P20789
A	253	ALA	ILE	engineered mutation	UNP P20789
A	305	ARG	HIS	engineered mutation	UNP P20789
A	358	VAL	PHE	engineered mutation	UNP P20789
A	362	ALA	SER	engineered mutation	UNP P20789
B	47	PRO	-	expression tag	UNP P20789
B	48	GLY	-	expression tag	UNP P20789

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	49	SER	-	expression tag	UNP P20789
B	391	THR	-	expression tag	UNP P20789
B	392	ARG	-	expression tag	UNP P20789
B	393	GLU	-	expression tag	UNP P20789
B	394	LEU	-	expression tag	UNP P20789
B	395	GLU	-	expression tag	UNP P20789
B	396	VAL	-	expression tag	UNP P20789
B	397	LEU	-	expression tag	UNP P20789
B	398	PHE	-	expression tag	UNP P20789
B	399	GLN	-	expression tag	UNP P20789
B	86	LEU	ALA	engineered mutation	UNP P20789
B	103	ASP	HIS	engineered mutation	UNP P20789
B	105	TYR	HIS	engineered mutation	UNP P20789
B	161	VAL	ALA	engineered mutation	UNP P20789
B	167	LEU	ARG	engineered mutation	UNP P20789
B	213	LEU	ARG	engineered mutation	UNP P20789
B	234	LEU	VAL	engineered mutation	UNP P20789
B	253	ALA	ILE	engineered mutation	UNP P20789
B	305	ARG	HIS	engineered mutation	UNP P20789
B	358	VAL	PHE	engineered mutation	UNP P20789
B	362	ALA	SER	engineered mutation	UNP P20789


- Molecule 2 is a protein called NEUROTENSIN/NEUROMEDIN N.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			58	38	12	8			
2	D	6	Total	C	N	O	0	0	0
			58	38	12	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	4	GLY	-	SEE REMARK 999	UNP P20068
C	5	PRO	-	SEE REMARK 999	UNP P20068
C	6	GLY	-	SEE REMARK 999	UNP P20068
C	7	GLY	-	SEE REMARK 999	UNP P20068
D	4	GLY	-	SEE REMARK 999	UNP P20068
D	5	PRO	-	SEE REMARK 999	UNP P20068
D	6	GLY	-	SEE REMARK 999	UNP P20068
D	7	GLY	-	SEE REMARK 999	UNP P20068

- Molecule 3 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NEUROTENSIN RECEPTOR TYPE 1




• Molecule 1: NEUROTENSIN RECEPTOR TYPE 1



• Molecule 2: NEUROTENSIN/NEUROMEDIN N



• Molecule 2: NEUROTENSIN/NEUROMEDIN N



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.31Å 89.40Å 212.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.82 – 2.75 19.82 – 2.75	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.82-2.75) 98.4 (19.82-2.75)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 2.77Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.248 , 0.273 0.272 , 0.301	Depositor DCC
R_{free} test set	1588 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	110.0	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5034	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2459	0.73	0/3355
1	B	0.29	0/2525	0.72	0/3451
2	C	0.22	0/59	0.77	0/77
2	D	0.24	0/59	0.77	0/77
All	All	0.29	0/5102	0.73	0/6960

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2400	0	2465	29	0
1	B	2463	0	2512	27	0
2	C	58	0	63	1	0
2	D	58	0	63	2	0
3	A	40	0	16	0	0
3	B	15	0	6	1	0
All	All	5034	0	5125	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:ARG:HH21	1:A:379:VAL:HG12	1.50	0.77
1:A:127:ASN:HD21	1:A:136:ALA:H	1.39	0.70
1:A:239:GLN:NE2	1:A:333:TYR:OH	2.25	0.69
1:B:56:ASP:O	2:D:8:ARG:NH2	2.26	0.69
1:B:239:GLN:NE2	1:B:333:TYR:OH	2.26	0.68
1:A:219:HIS:HD2	1:A:221:GLY:H	1.41	0.67
1:B:127:ASN:HD21	1:B:136:ALA:H	1.43	0.65
1:B:219:HIS:HD2	1:B:221:GLY:H	1.45	0.62
1:B:139:ASP:OD2	1:B:143:ARG:NH1	2.33	0.60
1:A:264:LEU:HD22	1:A:307:VAL:HG21	1.85	0.59
1:A:98:LEU:O	1:A:101:THR:OG1	2.20	0.57
1:A:323:PRO:HB2	1:A:354:THR:HG22	1.88	0.56
1:A:98:LEU:HD21	1:A:177:ALA:HB1	1.90	0.54
1:A:167:LEU:HD23	1:A:260:ILE:HD13	1.91	0.53
1:A:192:ALA:HA	1:B:240:VAL:HG22	1.91	0.53
1:A:100:SER:HB3	1:A:183:ARG:HH21	1.75	0.52
1:A:219:HIS:CD2	1:A:221:GLY:H	2.26	0.52
1:B:78:GLY:O	1:B:82:ASN:ND2	2.40	0.51
1:B:294:THR:OG1	1:B:295:ILE:N	2.37	0.51
1:B:174:PRO:HB3	1:B:295:ILE:HD13	1.93	0.50
1:A:54:ASP:O	2:C:8:ARG:NH1	2.45	0.50
1:B:219:HIS:CD2	1:B:221:GLY:H	2.29	0.49
1:A:67:VAL:HG21	1:A:352:MET:HB3	1.94	0.49
1:A:171:ILE:HG22	1:A:303:LEU:HG	1.95	0.49
1:A:375:ASN:O	1:A:379:VAL:HG23	2.14	0.48
1:A:241:ASN:O	1:A:245:SER:OG	2.29	0.48
1:B:211:GLN:HG3	1:B:228:ILE:HG22	1.95	0.48
1:A:139:ASP:OD2	1:A:143:ARG:NH1	2.47	0.48
1:B:209:GLY:HA3	1:B:228:ILE:HD13	1.97	0.47
1:B:145:TYR:CZ	1:B:149:ARG:HD2	2.50	0.46
1:B:57:VAL:HB	1:B:132:HIS:CD2	2.50	0.46
1:A:309:VAL:CG2	1:A:372:VAL:HG11	2.46	0.45
1:B:99:GLN:NE2	3:B:1388:GLY:O	2.50	0.45
1:B:219:HIS:HA	1:B:220:PRO:HD3	1.85	0.44
1:B:143:ARG:NE	1:B:205:LEU:O	2.43	0.44
1:B:181:MET:HB3	1:B:185:ARG:HB2	2.00	0.44
1:B:146:TYR:CZ	1:B:208:MET:HE2	2.53	0.44
1:B:101:THR:OG1	1:B:179:THR:HG21	2.18	0.44
1:A:124:GLU:OE1	1:A:359:TYR:OH	2.37	0.43
1:A:91:ARG:NH2	1:A:379:VAL:HG12	2.26	0.43
1:A:102:VAL:HA	1:A:105:TYR:HD2	1.84	0.42
1:A:127:ASN:HD21	1:A:136:ALA:N	2.10	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:323:PRO:HB2	1:B:354:THR:HG22	2.01	0.42
1:B:339:TRP:CD1	2:D:9:ARG:HG3	2.55	0.42
1:B:67:VAL:HG21	1:B:352:MET:HB3	2.01	0.42
1:B:292:ASN:O	1:B:292:ASN:ND2	2.48	0.42
1:B:291:PHE:HB2	1:B:293:MET:HE2	2.03	0.41
1:A:164:SER:HA	1:A:310:LEU:HD21	2.03	0.41
1:A:57:VAL:HB	1:A:132:HIS:CD2	2.56	0.41
1:B:167:LEU:HD21	1:B:307:VAL:HA	2.01	0.41
1:B:164:SER:HA	1:B:310:LEU:HD21	2.02	0.41
1:A:127:ASN:ND2	1:A:136:ALA:H	2.12	0.40
1:A:219:HIS:HA	1:A:220:PRO:HD3	1.93	0.40
1:A:75:PHE:HB2	1:A:121:MET:SD	2.61	0.40
1:A:324:TYR:CZ	1:A:328:ARG:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/335 (89%)	285 (96%)	13 (4%)	0	100	100
1	B	310/335 (92%)	296 (96%)	14 (4%)	0	100	100
2	C	4/10 (40%)	4 (100%)	0	0	100	100
2	D	4/10 (40%)	4 (100%)	0	0	100	100
All	All	616/690 (89%)	589 (96%)	27 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/290 (91%)	258 (98%)	6 (2%)	44	66
1	B	270/290 (93%)	269 (100%)	1 (0%)	84	91
2	C	6/7 (86%)	6 (100%)	0	100	100
2	D	6/7 (86%)	6 (100%)	0	100	100
All	All	546/594 (92%)	539 (99%)	7 (1%)	61	76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	102	VAL
1	A	129	ILE
1	A	131	VAL
1	A	237	VAL
1	A	383	THR
1	A	384	LEU
1	B	292	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	127	ASN
1	A	211	GLN
1	A	219	HIS
1	A	239	GLN
1	A	301	GLN
1	A	338	GLN
1	A	365	ASN
1	A	370	ASN
1	B	127	ASN
1	B	211	GLN
1	B	219	HIS
1	B	239	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	270	GLN
1	B	370	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLY	A	1392	-	4,4,4	1.15	1 (25%)	3,4,4	1.66	1 (33%)
3	GLY	A	1388	-	4,4,4	1.15	1 (25%)	3,4,4	1.64	1 (33%)
3	GLY	A	1390	-	4,4,4	1.13	1 (25%)	3,4,4	1.65	1 (33%)
3	GLY	B	1389	-	4,4,4	1.16	1 (25%)	3,4,4	1.62	1 (33%)
3	GLY	A	1389	-	4,4,4	1.15	1 (25%)	3,4,4	1.63	1 (33%)
3	GLY	A	1386	-	4,4,4	1.15	1 (25%)	3,4,4	1.64	1 (33%)
3	GLY	A	1385	-	4,4,4	1.15	1 (25%)	3,4,4	1.62	1 (33%)
3	GLY	A	1391	-	4,4,4	1.13	1 (25%)	3,4,4	1.61	0
3	GLY	B	1387	-	4,4,4	1.14	1 (25%)	3,4,4	1.65	1 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLY	B	1388	-	4,4,4	1.17	1 (25%)	3,4,4	1.67	1 (33%)
3	GLY	A	1387	-	4,4,4	1.14	1 (25%)	3,4,4	1.65	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLY	A	1392	-	-	0/2/2/2	-
3	GLY	A	1388	-	-	0/2/2/2	-
3	GLY	A	1390	-	-	0/2/2/2	-
3	GLY	B	1389	-	-	2/2/2/2	-
3	GLY	A	1389	-	-	0/2/2/2	-
3	GLY	A	1386	-	-	0/2/2/2	-
3	GLY	A	1385	-	-	0/2/2/2	-
3	GLY	A	1391	-	-	0/2/2/2	-
3	GLY	B	1387	-	-	0/2/2/2	-
3	GLY	B	1388	-	-	0/2/2/2	-
3	GLY	A	1387	-	-	0/2/2/2	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1388	GLY	OXT-C	-2.22	1.23	1.30
3	B	1389	GLY	OXT-C	-2.20	1.23	1.30
3	A	1385	GLY	OXT-C	-2.19	1.23	1.30
3	A	1386	GLY	OXT-C	-2.19	1.23	1.30
3	A	1389	GLY	OXT-C	-2.18	1.23	1.30
3	A	1392	GLY	OXT-C	-2.18	1.23	1.30
3	A	1388	GLY	OXT-C	-2.18	1.23	1.30
3	A	1387	GLY	OXT-C	-2.17	1.23	1.30
3	B	1387	GLY	OXT-C	-2.15	1.23	1.30
3	A	1390	GLY	OXT-C	-2.15	1.23	1.30
3	A	1391	GLY	OXT-C	-2.13	1.23	1.30

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1388	GLY	OXT-C-O	-2.12	117.87	123.33
3	A	1392	GLY	OXT-C-O	-2.10	117.93	123.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1390	GLY	OXT-C-O	-2.10	117.94	123.33
3	B	1387	GLY	OXT-C-O	-2.09	117.94	123.33
3	A	1387	GLY	OXT-C-O	-2.08	117.98	123.33
3	A	1386	GLY	OXT-C-O	-2.07	118.02	123.33
3	A	1388	GLY	OXT-C-O	-2.06	118.04	123.33
3	A	1389	GLY	OXT-C-O	-2.05	118.06	123.33
3	A	1385	GLY	OXT-C-O	-2.04	118.09	123.33
3	B	1389	GLY	OXT-C-O	-2.03	118.11	123.33

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1389	GLY	O-C-CA-N
3	B	1389	GLY	OXT-C-CA-N

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1388	GLY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/335 (90%)	-0.07	7 (2%) 61 59	25, 63, 151, 300	0
1	B	314/335 (93%)	0.06	8 (2%) 58 56	32, 78, 163, 298	0
2	C	6/10 (60%)	0.27	0 100 100	45, 52, 77, 84	0
2	D	6/10 (60%)	0.09	0 100 100	57, 63, 87, 90	0
All	All	630/690 (91%)	0.00	15 (2%) 59 57	25, 69, 158, 300	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	ASN	4.7
1	B	295	ILE	4.3
1	B	156	THR	4.1
1	B	372	VAL	3.4
1	A	344	PHE	2.9
1	B	100	SER	2.8
1	A	98	LEU	2.8
1	B	180	LEU	2.8
1	B	157	ALA	2.7
1	A	52	ASN	2.6
1	A	152	CYS	2.3
1	A	124	GLU	2.2
1	B	53	SER	2.2
1	A	384	LEU	2.2
1	A	376	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GLY	A	1387	5/5	0.55	0.24	92,92,92,125	0
3	GLY	A	1391	5/5	0.56	0.25	89,89,89,101	0
3	GLY	A	1386	5/5	0.61	0.23	90,102,102,102	0
3	GLY	A	1392	5/5	0.68	0.14	68,77,77,77	0
3	GLY	B	1388	5/5	0.69	0.09	96,104,104,104	0
3	GLY	B	1387	5/5	0.83	0.14	79,94,94,94	0
3	GLY	A	1388	5/5	0.84	0.17	76,98,98,98	0
3	GLY	A	1390	5/5	0.84	0.28	72,72,72,144	0
3	GLY	B	1389	5/5	0.85	0.39	108,108,108,143	0
3	GLY	A	1389	5/5	0.87	0.14	89,89,89,107	0
3	GLY	A	1385	5/5	0.89	0.13	67,67,67,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.