



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 02:41 PM EDT

PDB ID : 3BVE / pdb_00003bve
Title : Structural basis for the iron uptake mechanism of Helicobacter pylori ferritin
Authors : Kim, K.H.; Cho, K.J.; Lee, J.H.; Shin, H.J.; Yang, I.S.
Deposited on : 2008-01-07
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	
1	B	181	
1	C	181	
1	D	181	
1	E	181	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	181	2% 85% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9460 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	173	Total	C	N	O	S	0	3	0
			1433	912	241	275	5			
1	B	173	Total	C	N	O	S	0	6	0
			1450	925	244	276	5			
1	C	173	Total	C	N	O	S	0	4	0
			1441	916	242	277	6			
1	D	173	Total	C	N	O	S	0	5	0
			1440	917	243	275	5			
1	E	172	Total	C	N	O	S	0	6	0
			1434	914	239	276	5			
1	F	173	Total	C	N	O	S	0	4	0
			1434	912	240	277	5			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1013	MET	-	expression tag	UNP Q9ZLI1
A	1012	GLY	-	expression tag	UNP Q9ZLI1
A	1011	SER	-	expression tag	UNP Q9ZLI1
A	1010	SER	-	expression tag	UNP Q9ZLI1
A	1009	HIS	-	expression tag	UNP Q9ZLI1
A	1008	HIS	-	expression tag	UNP Q9ZLI1
A	1007	HIS	-	expression tag	UNP Q9ZLI1
A	1006	HIS	-	expression tag	UNP Q9ZLI1
A	1005	HIS	-	expression tag	UNP Q9ZLI1
A	1004	HIS	-	expression tag	UNP Q9ZLI1
A	1003	SER	-	expression tag	UNP Q9ZLI1
A	1002	GLN	-	expression tag	UNP Q9ZLI1
A	1001	ASP	-	expression tag	UNP Q9ZLI1
A	0	PRO	-	expression tag	UNP Q9ZLI1
B	1013	MET	-	expression tag	UNP Q9ZLI1
B	1012	GLY	-	expression tag	UNP Q9ZLI1
B	1011	SER	-	expression tag	UNP Q9ZLI1

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	1010	SER	-	expression tag	UNP Q9ZLI1
B	1009	HIS	-	expression tag	UNP Q9ZLI1
B	1008	HIS	-	expression tag	UNP Q9ZLI1
B	1007	HIS	-	expression tag	UNP Q9ZLI1
B	1006	HIS	-	expression tag	UNP Q9ZLI1
B	1005	HIS	-	expression tag	UNP Q9ZLI1
B	1004	HIS	-	expression tag	UNP Q9ZLI1
B	1003	SER	-	expression tag	UNP Q9ZLI1
B	1002	GLN	-	expression tag	UNP Q9ZLI1
B	1001	ASP	-	expression tag	UNP Q9ZLI1
B	0	PRO	-	expression tag	UNP Q9ZLI1
C	1013	MET	-	expression tag	UNP Q9ZLI1
C	1012	GLY	-	expression tag	UNP Q9ZLI1
C	1011	SER	-	expression tag	UNP Q9ZLI1
C	1010	SER	-	expression tag	UNP Q9ZLI1
C	1009	HIS	-	expression tag	UNP Q9ZLI1
C	1008	HIS	-	expression tag	UNP Q9ZLI1
C	1007	HIS	-	expression tag	UNP Q9ZLI1
C	1006	HIS	-	expression tag	UNP Q9ZLI1
C	1005	HIS	-	expression tag	UNP Q9ZLI1
C	1004	HIS	-	expression tag	UNP Q9ZLI1
C	1003	SER	-	expression tag	UNP Q9ZLI1
C	1002	GLN	-	expression tag	UNP Q9ZLI1
C	1001	ASP	-	expression tag	UNP Q9ZLI1
C	0	PRO	-	expression tag	UNP Q9ZLI1
D	1013	MET	-	expression tag	UNP Q9ZLI1
D	1012	GLY	-	expression tag	UNP Q9ZLI1
D	1011	SER	-	expression tag	UNP Q9ZLI1
D	1010	SER	-	expression tag	UNP Q9ZLI1
D	1009	HIS	-	expression tag	UNP Q9ZLI1
D	1008	HIS	-	expression tag	UNP Q9ZLI1
D	1007	HIS	-	expression tag	UNP Q9ZLI1
D	1006	HIS	-	expression tag	UNP Q9ZLI1
D	1005	HIS	-	expression tag	UNP Q9ZLI1
D	1004	HIS	-	expression tag	UNP Q9ZLI1
D	1003	SER	-	expression tag	UNP Q9ZLI1
D	1002	GLN	-	expression tag	UNP Q9ZLI1
D	1001	ASP	-	expression tag	UNP Q9ZLI1
D	0	PRO	-	expression tag	UNP Q9ZLI1
E	1013	MET	-	expression tag	UNP Q9ZLI1
E	1012	GLY	-	expression tag	UNP Q9ZLI1
E	1011	SER	-	expression tag	UNP Q9ZLI1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1010	SER	-	expression tag	UNP Q9ZLI1
E	1009	HIS	-	expression tag	UNP Q9ZLI1
E	1008	HIS	-	expression tag	UNP Q9ZLI1
E	1007	HIS	-	expression tag	UNP Q9ZLI1
E	1006	HIS	-	expression tag	UNP Q9ZLI1
E	1005	HIS	-	expression tag	UNP Q9ZLI1
E	1004	HIS	-	expression tag	UNP Q9ZLI1
E	1003	SER	-	expression tag	UNP Q9ZLI1
E	1002	GLN	-	expression tag	UNP Q9ZLI1
E	1001	ASP	-	expression tag	UNP Q9ZLI1
E	0	PRO	-	expression tag	UNP Q9ZLI1
F	1013	MET	-	expression tag	UNP Q9ZLI1
F	1012	GLY	-	expression tag	UNP Q9ZLI1
F	1011	SER	-	expression tag	UNP Q9ZLI1
F	1010	SER	-	expression tag	UNP Q9ZLI1
F	1009	HIS	-	expression tag	UNP Q9ZLI1
F	1008	HIS	-	expression tag	UNP Q9ZLI1
F	1007	HIS	-	expression tag	UNP Q9ZLI1
F	1006	HIS	-	expression tag	UNP Q9ZLI1
F	1005	HIS	-	expression tag	UNP Q9ZLI1
F	1004	HIS	-	expression tag	UNP Q9ZLI1
F	1003	SER	-	expression tag	UNP Q9ZLI1
F	1002	GLN	-	expression tag	UNP Q9ZLI1
F	1001	ASP	-	expression tag	UNP Q9ZLI1
F	0	PRO	-	expression tag	UNP Q9ZLI1

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total	O	0	0
			126	126		
3	B	140	Total	O	0	0
			140	140		
3	C	132	Total	O	0	0
			132	132		
3	D	131	Total	O	0	0
			131	131		
3	E	143	Total	O	0	0
			143	143		
3	F	138	Total	O	0	0
			138	138		

- Molecule 1: Ferritin





4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	128.71Å 128.71Å 165.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.90 – 1.80 34.90 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.2 (34.90-1.80) 96.2 (34.90-1.80)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.222 0.186 , 0.223	Depositor DCC
R_{free} test set	5988 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9460	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1476	0.76	0/1992
1	B	0.48	0/1503	0.77	0/2027
1	C	0.48	0/1481	0.76	0/1998
1	D	0.46	0/1490	0.77	0/2011
1	E	0.46	0/1486	0.75	0/2005
1	F	0.49	0/1479	0.76	0/1996
All	All	0.47	0/8915	0.76	0/12029

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1433	0	1379	10	0
1	B	1450	0	1406	15	0
1	C	1441	0	1384	18	0
1	D	1440	0	1390	19	0
1	E	1434	0	1388	15	0
1	F	1434	0	1383	14	0
2	B	6	0	8	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	6	0	8	0	0
2	F	6	0	8	0	0
3	A	126	0	0	1	0
3	B	140	0	0	1	0
3	C	132	0	0	5	0
3	D	131	0	0	5	0
3	E	143	0	0	3	0
3	F	138	0	0	2	0
All	All	9460	0	8354	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 83 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:88[A]:GLN:NE2	3:E:1146:HOH:O	2.08	0.87
1:D:88[A]:GLN:NE2	3:D:1132:HOH:O	2.10	0.83
1:B:19:ASN:HD21	1:B:74:SER:H	1.22	0.82
1:F:88:GLN:HG3	3:F:1056:HOH:O	1.86	0.76
1:E:30:TRP:O	1:E:34:HIS:HD2	1.74	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/181 (96%)	171 (98%)	2 (1%)	1 (1%)	21	11
1	B	177/181 (98%)	175 (99%)	1 (1%)	1 (1%)	21	11
1	C	175/181 (97%)	172 (98%)	2 (1%)	1 (1%)	21	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	176/181 (97%)	174 (99%)	2 (1%)	0	100	100
1	E	176/181 (97%)	172 (98%)	3 (2%)	1 (1%)	21	11
1	F	175/181 (97%)	171 (98%)	3 (2%)	1 (1%)	21	11
All	All	1053/1086 (97%)	1035 (98%)	13 (1%)	5 (0%)	24	14

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	148	ASN
1	F	148	ASN
1	A	148	ASN
1	E	148	ASN
1	C	148	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/163 (98%)	156 (98%)	3 (2%)	50	41
1	B	162/163 (99%)	160 (99%)	2 (1%)	63	57
1	C	160/163 (98%)	158 (99%)	2 (1%)	61	54
1	D	161/163 (99%)	156 (97%)	5 (3%)	35	23
1	E	161/163 (99%)	157 (98%)	4 (2%)	42	30
1	F	160/163 (98%)	159 (99%)	1 (1%)	78	77
All	All	963/978 (98%)	946 (98%)	17 (2%)	57	43

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	140	LYS
1	F	151	LEU
1	D	1006	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	85	GLN
1	D	112	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	88	GLN
1	D	85	GLN
1	F	156	GLN
1	F	34	HIS
1	C	96	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	1007	-	5,5,5	0.38	0	5,5,5	0.41	0
2	GOL	F	1006	-	5,5,5	0.39	0	5,5,5	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	1007	-	5,5,5	0.39	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	1007	-	-	2/4/4/4	-
2	GOL	F	1006	-	-	4/4/4/4	-
2	GOL	B	1007	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1007	GOL	O1-C1-C2-C3
2	C	1007	GOL	O1-C1-C2-C3
2	F	1006	GOL	O1-C1-C2-C3
2	B	1007	GOL	O1-C1-C2-O2
2	F	1006	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1007	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	173/181 (95%)	-0.29	3 (1%) 69 69	9, 15, 26, 35	3 (1%)
1	B	173/181 (95%)	-0.22	3 (1%) 69 69	9, 16, 26, 36	6 (3%)
1	C	173/181 (95%)	-0.26	5 (2%) 53 53	7, 15, 28, 36	4 (2%)
1	D	173/181 (95%)	-0.11	4 (2%) 61 61	10, 17, 29, 37	5 (2%)
1	E	172/181 (95%)	-0.27	3 (1%) 69 69	8, 15, 26, 35	6 (3%)
1	F	173/181 (95%)	-0.33	4 (2%) 61 61	9, 14, 24, 40	4 (2%)
All	All	1037/1086 (95%)	-0.25	22 (2%) 63 64	7, 16, 27, 40	28 (2%)

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	148	ASN	5.3
1	E	146	ASN	4.4
1	D	73	ILE	4.1
1	A	148	ASN	3.8
1	E	148	ASN	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	F	1006	6/6	0.80	0.17	39,42,45,47	0
2	GOL	C	1007	6/6	0.81	0.24	45,52,56,59	0
2	GOL	B	1007	6/6	0.88	0.14	33,40,42,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.