



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 23, 2026 – 12:59 AM UTC

PDB ID : 3BVV / pdb_00003bvv
Title : Golgi mannosidase II D204A catalytic nucleophile mutant complex with METHYL ALPHA-D-MANNOPYRANOSYL-(1->3)-[6-THIO-ALPHA-D-MANNOPYRANOSYL-(1->6)]-BETA-D-MANNOPYRANOSIDE
Authors : Kuntz, D.A.; Rose, D.R.
Deposited on : 2008-01-07
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1045	4% 90% 7%
2	B	3	33% 67%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10336 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-mannosidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1016	Total	C	N	O	S	0	55	0
			8581	5465	1505	1566	45			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ARG	-	expression tag	UNP Q24451
A	2	SER	-	expression tag	UNP Q24451
A	3	SER	-	expression tag	UNP Q24451
A	4	HIS	-	expression tag	UNP Q24451
A	5	HIS	-	expression tag	UNP Q24451
A	6	HIS	-	expression tag	UNP Q24451
A	7	HIS	-	expression tag	UNP Q24451
A	8	HIS	-	expression tag	UNP Q24451
A	9	HIS	-	expression tag	UNP Q24451
A	10	GLY	-	expression tag	UNP Q24451
A	11	GLU	-	expression tag	UNP Q24451
A	12	PHE	-	expression tag	UNP Q24451
A	204	ALA	ASP	engineered mutation	UNP Q24451
A	907	LYS	GLU	SEE REMARK 999	UNP Q24451

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[6-thio-alpha-D-mannopyranose-(1-6)]methyl beta-D-mannopyranoside.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	O	S	0	0	0
			35	19	15	1			

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1657	Total	O	0	36
			1690	1690		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-mannosidase 2



- Molecule 2: alpha-D-mannopyranose-(1-3)-[6-thio-alpha-D-mannopyranose-(1-6)]methyl beta-D-mannopyranoside



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.10Å 109.81Å 139.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.61 – 1.30 19.61 – 1.30	Depositor EDS
% Data completeness (in resolution range)	97.8 (19.61-1.30) 98.0 (19.61-1.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 1.30Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.152 , 0.173 0.150 , 0.148	Depositor DCC
R_{free} test set	3816 reflections (1.47%)	wwPDB-VP
Wilson B-factor (Å ²)	12.1	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10336	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MAN, MRD, MPD, ZN, YZ0, Z4Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/8909	0.77	0/12082

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8581	0	8429	81	0
2	B	35	0	8	3	0
3	A	5	0	0	0	0
4	A	1	0	0	0	0
5	A	8	0	14	0	0
6	A	16	0	28	2	0
7	A	1690	0	0	38	0
All	All	10336	0	8479	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:663:GLN:HG3	7:A:1741:HOH:O	1.60	0.99
1:A:96[A]:ILE:HG22	7:A:2227:HOH:O	1.71	0.91
1:A:846:LEU:HD23	1:A:864[B]:ILE:CD1	2.00	0.91
1:A:75[B]:TYR:HD2	7:A:1385:HOH:O	1.53	0.90
1:A:82[A]:LYS:HD2	1:A:375[A]:GLN:HE22	1.38	0.88
1:A:96[A]:ILE:CG2	7:A:2227:HOH:O	2.23	0.83
6:A:1104:MPD:H53	7:A:1597[B]:HOH:O	1.84	0.77
1:A:846:LEU:CD2	1:A:864[B]:ILE:HD12	2.16	0.76
1:A:846:LEU:HD23	1:A:864[B]:ILE:HD12	1.67	0.74
1:A:846:LEU:HD23	1:A:864[B]:ILE:HD11	1.67	0.74
1:A:380:GLN:HG2	7:A:2289:HOH:O	1.89	0.71
1:A:346[A]:GLN:NE2	7:A:1201:HOH:O	2.23	0.71
1:A:567:GLN:HB2	7:A:2160:HOH:O	1.90	0.71
7:A:2336[B]:HOH:O	2:B:1:YZ0:C1M	2.43	0.67
1:A:250:ILE:HD11	7:A:2239:HOH:O	1.94	0.66
1:A:435[B]:TYR:CE2	7:A:1597[B]:HOH:O	2.48	0.66
1:A:954[A]:ARG:HH21	1:A:981[A]:ARG:NE	1.93	0.66
1:A:971[A]:LYS:HE2	7:A:2483:HOH:O	1.96	0.64
1:A:136[B]:ARG:HD2	7:A:1365:HOH:O	1.97	0.64
1:A:678:LEU:HD12	1:A:769[B]:MET:HE1	1.80	0.64
1:A:957:LEU:HD11	1:A:979[B]:LEU:HG	1.81	0.62
1:A:295:LEU:HD22	2:B:3:Z4Y:S6	2.39	0.62
1:A:138[A]:TYR:CE1	1:A:193[A]:MET:CE	2.83	0.61
1:A:742[A]:GLN:HG3	7:A:2024:HOH:O	2.01	0.61
1:A:541:THR:HG23	7:A:1485:HOH:O	2.00	0.61
1:A:82[A]:LYS:CD	1:A:375[A]:GLN:HE22	2.12	0.60
1:A:709[A]:HIS:ND1	7:A:1205:HOH:O	2.31	0.60
1:A:704:PRO:HA	7:A:2202[B]:HOH:O	2.02	0.59
6:A:1105:MPD:H12	7:A:2581:HOH:O	2.03	0.58
1:A:82[A]:LYS:HD2	1:A:375[A]:GLN:NE2	2.14	0.58
1:A:904:ARG:HG2	1:A:985:MET:SD	2.44	0.57
1:A:895:VAL:HG12	1:A:897:GLU:HG3	1.87	0.57
1:A:228[A]:ARG:NE	7:A:1215[A]:HOH:O	2.38	0.56
1:A:51:ARG:NH2	7:A:1214:HOH:O	2.38	0.56
1:A:992[A]:GLU:HG2	1:A:993:HIS:H	1.70	0.55
1:A:82[A]:LYS:CE	1:A:375[A]:GLN:NE2	2.69	0.55
1:A:678:LEU:CD1	1:A:769[B]:MET:HE1	2.38	0.54
1:A:846:LEU:HD21	1:A:864[B]:ILE:HD12	1.90	0.54
1:A:491:GLN:NE2	7:A:1206:HOH:O	2.32	0.53
1:A:75[B]:TYR:CD2	7:A:1385:HOH:O	2.41	0.53
1:A:848[B]:THR:CG2	1:A:850:GLN:NE2	2.72	0.53
1:A:32:GLN:NE2	7:A:1221:HOH:O	2.41	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:LYS:NZ	7:A:1227:HOH:O	2.44	0.51
1:A:241[B]:ARG:CZ	7:A:1503:HOH:O	2.58	0.51
1:A:704:PRO:HG3	7:A:2709:HOH:O	2.10	0.50
1:A:174:HIS:CE1	1:A:176:ARG:HD3	2.46	0.50
1:A:256:ASP:HB2	7:A:1221:HOH:O	2.11	0.49
1:A:990:PRO:O	1:A:991:GLU:C	2.55	0.49
1:A:651:LEU:CD1	1:A:653:ARG:HG2	2.43	0.48
1:A:62:TRP:CD2	1:A:65:GLY:HA3	2.49	0.48
1:A:138[A]:TYR:CE1	1:A:193[A]:MET:HE1	2.48	0.48
1:A:993:HIS:O	1:A:993:HIS:ND1	2.48	0.47
1:A:954[A]:ARG:HH21	1:A:981[A]:ARG:CD	2.27	0.47
1:A:979[A]:LEU:HD21	1:A:999:VAL:HG11	1.97	0.47
1:A:990:PRO:O	1:A:991:GLU:O	2.33	0.47
1:A:251:TRP:CD1	1:A:251:TRP:C	2.93	0.46
1:A:82[B]:LYS:HD2	7:A:2337:HOH:O	2.16	0.46
1:A:707:PRO:HG2	1:A:797[A]:HIS:CD2	2.51	0.46
1:A:580:VAL:HG22	1:A:634:LEU:HD22	1.97	0.45
1:A:138[B]:TYR:CE1	1:A:146[B]:LYS:HE3	2.52	0.45
1:A:341:ASP:OD2	2:B:1:YZ0:O4	2.34	0.45
1:A:75[B]:TYR:HE2	7:A:2245:HOH:O	1.99	0.44
1:A:82[B]:LYS:HD2	7:A:2280:HOH:O	2.16	0.44
1:A:848[B]:THR:HG22	1:A:850:GLN:H	1.83	0.44
1:A:797[B]:HIS:HA	7:A:2283:HOH:O	2.19	0.43
1:A:686:LEU:HD22	1:A:697:ILE:HG12	2.01	0.43
1:A:32:GLN:HG3	7:A:1484:HOH:O	2.18	0.43
1:A:96[A]:ILE:HG23	1:A:479:LYS:HE2	2.01	0.42
1:A:228[A]:ARG:NH2	7:A:1215[A]:HOH:O	2.52	0.42
1:A:848[B]:THR:HG22	1:A:850:GLN:NE2	2.34	0.42
1:A:582:ASP:HB2	7:A:1477[A]:HOH:O	2.19	0.42
1:A:82[A]:LYS:CD	1:A:375[A]:GLN:NE2	2.79	0.42
1:A:280:LYS:NZ	1:A:301:VAL:HG21	2.35	0.42
1:A:848[B]:THR:HG23	1:A:850:GLN:NE2	2.35	0.41
1:A:992[A]:GLU:HB3	1:A:993:HIS:CD2	2.56	0.41
1:A:995[B]:GLN:H	1:A:995[B]:GLN:NE2	2.19	0.41
1:A:797[A]:HIS:HB2	7:A:2554:HOH:O	2.21	0.40
1:A:898:LYS:HA	1:A:898:LYS:HD2	1.86	0.40
1:A:325:LYS:NZ	7:A:2177[B]:HOH:O	2.54	0.40
1:A:488:GLN:HG3	7:A:2005:HOH:O	2.20	0.40
1:A:906:SER:HB3	7:A:2412:HOH:O	2.21	0.40
1:A:934:PHE:CE2	1:A:936[A]:GLU:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1069/1045 (102%)	1040 (97%)	25 (2%)	4 (0%)	30	9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	991	GLU
1	A	95	TRP
1	A	993	HIS
1	A	204	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	957/928 (103%)	948 (99%)	9 (1%)	70	42

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	ASN
1	A	275	CYS
1	A	394	GLN
1	A	653	ARG
1	A	655	ASN
1	A	663	GLN
1	A	691	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	898	LYS
1	A	991	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	32	GLN
1	A	148	GLN
1	A	347	ASN
1	A	380	GLN
1	A	663	GLN
1	A	923	GLN
1	A	1005	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	YZ0	B	1	2	13,13,13	0.74	0	18,18,18	1.57	3 (16%)
2	MAN	B	2	2,4	11,11,12	1.26	1 (9%)	15,15,17	1.00	1 (6%)
2	Z4Y	B	3	2	11,11,12	1.40	1 (9%)	14,15,17	1.67	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YZ0	B	1	2	-	4/4/24/24	0/1/1/1
2	MAN	B	2	2,4	-	0/2/19/22	0/1/1/1
2	Z4Y	B	3	2	-	0/2/19/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	MAN	C2-C3	2.67	1.56	1.52
2	B	3	Z4Y	C2-C3	2.46	1.56	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	YZ0	C4-C3-C2	-4.12	103.59	110.83
2	B	3	Z4Y	O5-C5-C6	3.33	112.06	106.50
2	B	1	YZ0	C1M-O1-C1	-3.19	108.41	113.26
2	B	3	Z4Y	O4-C4-C3	3.06	117.58	110.38
2	B	2	MAN	C1-C2-C3	2.95	113.94	109.64
2	B	3	Z4Y	C6-C5-C4	2.62	119.41	112.86
2	B	1	YZ0	O6-C6-C5	-2.32	103.44	111.33
2	B	3	Z4Y	C3-C4-C5	-2.20	106.24	110.23

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	YZ0	O5-C1-O1-C1M
2	B	1	YZ0	C2-C1-O1-C1M
2	B	1	YZ0	O5-C5-C6-O6
2	B	1	YZ0	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	YZ0	2	0
2	B	3	Z4Y	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	MPD	A	1104	-	7,7,7	0.38	0	9,10,10	0.51	0
6	MPD	A	1105	-	7,7,7	0.42	0	9,10,10	0.47	0
3	PO4	A	1101	-	4,4,4	0.95	0	6,6,6	0.99	0
5	MRD	A	1103	-	7,7,7	0.24	0	9,10,10	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MPD	A	1104	-	-	3/5/5/5	-
6	MPD	A	1105	-	-	1/5/5/5	-
5	MRD	A	1103	-	-	3/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1103	MRD	C2-C3-C4-O4
5	A	1103	MRD	C2-C3-C4-C5
6	A	1104	MPD	C1-C2-C3-C4
6	A	1105	MPD	O2-C2-C3-C4
6	A	1104	MPD	CM-C2-C3-C4
5	A	1103	MRD	O2-C2-C3-C4
6	A	1104	MPD	C2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1104	MPD	1	0
6	A	1105	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1016/1045 (97%)	0.02	42 (4%)	41 47	4, 12, 24, 33	55 (5%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	990	PRO	7.6
1	A	534	SER	6.3
1	A	993	HIS	6.0
1	A	602	THR	5.4
1	A	538	ASP	4.8
1	A	991	GLU	4.8
1	A	702	ASP	4.5
1	A	1045	SER	4.4
1	A	539	SER	4.3
1	A	600	HIS	4.0
1	A	1024	ASP	3.9
1	A	720	HIS	3.9
1	A	701	GLN	3.7
1	A	989	THR	3.7
1	A	987	CYS	3.4
1	A	537	GLU	3.4
1	A	603	LEU	3.1
1	A	78	HIS	3.1
1	A	396	GLU	3.1
1	A	682	ASN	3.0
1	A	30	GLN	2.9
1	A	1028	ALA	2.8
1	A	638	ASP	2.7
1	A	703	SER	2.6
1	A	604	THR	2.5
1	A	662	GLY	2.5
1	A	366	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	ASN	2.5
1	A	601	ASP	2.4
1	A	139	HIS	2.4
1	A	613	THR	2.3
1	A	784	SER	2.2
1	A	1022	HIS	2.2
1	A	57	ILE	2.2
1	A	700	THR	2.2
1	A	994	THR	2.2
1	A	655	ASN	2.2
1	A	1029	PRO	2.2
1	A	988	GLY	2.1
1	A	1044	SER	2.1
1	A	583	LEU	2.0
1	A	547	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	Z4Y	B	3	11/12	0.79	0.17	10,18,24,29	11
2	YZ0	B	1	13/13	0.92	0.08	14,18,21,25	12
2	MAN	B	2	11/12	0.95	0.08	7,12,13,13	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MPD	A	1105	8/8	0.75	0.14	26,29,31,32	0
6	MPD	A	1104	8/8	0.76	0.14	33,34,34,35	0
5	MRD	A	1103	8/8	0.94	0.09	13,18,19,21	0
3	PO4	A	1101	5/5	0.95	0.09	12,15,17,18	5
4	ZN	A	1102	1/1	1.00	0.01	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.