



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:46 PM UTC

PDB ID : 5BV5 / pdb_00005bv5
Title : Structure of CYP119 with T213A and C317H mutations
Authors : Buller, A.R.; Heel, T.; McIntosh, J.A.; Arnold, F.H.
Deposited on : 2015-06-04
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	5% 82% 14%
1	B	368	6% 73% 5% 22%
1	C	368	5% 80% 15%
1	D	368	5% 82% 7% 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 119.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2411	1545	412	450	4			
1	B	288	Total	C	N	O	S	0	0	0
			2236	1438	382	413	3			
1	D	327	Total	C	N	O	S	0	0	0
			2553	1648	427	474	4			
1	C	311	Total	C	N	O	S	0	0	0
			2401	1537	409	451	4			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

-
- Chemical structure of HEM (heme) showing a central iron atom coordinated by four nitrogen atoms in a porphyrin-like ring, with various side chains and a central heme label.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C N 11 9 2	0	0
4	D	1	Total C N 11 9 2	0	0
4	C	1	Total C N 11 9 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total O 2 2	0	0
5	B	3	Total O 3 3	0	0
5	D	3	Total O 3 3	0	0
5	C	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome P450 119




• Molecule 1: Cytochrome P450 119



• Molecule 1: Cytochrome P450 119



● Molecule 1: Cytochrome P450 119



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.49Å 137.87Å 91.78Å 90.00° 101.18° 90.00°	Depositor
Resolution (Å)	50.00 – 2.70 50.00 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.8 (50.00-2.70) 97.8 (50.00-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.238 , 0.277 0.237 , 0.275	Depositor DCC
R_{free} test set	2143 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	63.4	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9827	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, HEM, PIM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2458	0.67	0/3341
1	B	0.32	0/2280	0.67	0/3098
1	C	0.33	0/2450	0.67	0/3333
1	D	0.32	0/2607	0.67	0/3545
All	All	0.32	0/9795	0.67	0/13317

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2411	0	2267	7	0
1	B	2236	0	2123	11	0
1	C	2401	0	2283	6	0
1	D	2553	0	2453	13	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	43	0	30	4	0
3	B	43	0	30	5	0
3	C	43	0	30	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	43	0	30	5	0
4	B	11	0	8	0	0
4	C	11	0	8	0	0
4	D	11	0	8	1	0
5	A	2	0	0	0	0
5	B	3	0	0	0	0
5	C	3	0	0	0	0
5	D	3	0	0	0	0
All	All	9827	0	9270	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:29:THR:HG23	1:B:280:VAL:HG13	1.65	0.77
1:A:29:THR:HG23	1:A:280:VAL:CG1	2.21	0.71
3:C:500:HEM:HMC1	3:C:500:HEM:HBC2	1.74	0.70
3:C:500:HEM:HMB1	3:C:500:HEM:HBB2	1.74	0.69
3:D:500:HEM:HBB2	3:D:500:HEM:HMB1	1.75	0.69
3:A:602:HEM:HBC2	3:A:602:HEM:HMC1	1.75	0.67
3:B:602:HEM:HBC2	3:B:602:HEM:HMC2	1.75	0.67
3:D:500:HEM:HMC2	3:D:500:HEM:HBC2	1.77	0.66
3:B:602:HEM:HMB1	3:B:602:HEM:HBB2	1.77	0.66
3:A:602:HEM:HBB2	3:A:602:HEM:HMB2	1.80	0.63
1:C:29:THR:HG23	1:C:280:VAL:CG1	2.29	0.62
1:B:29:THR:HG23	1:B:280:VAL:CG1	2.34	0.58
1:A:223:ILE:HG12	1:A:360:LEU:HD11	1.86	0.56
1:D:26:TYR:CD2	1:D:291:VAL:HG21	2.41	0.55
1:D:125:LEU:HB3	1:D:126:PRO:HD3	1.92	0.52
1:A:223:ILE:HD13	1:A:343:ILE:HD13	1.91	0.52
1:D:136:LEU:HD22	1:D:171:LEU:HD22	1.92	0.51
1:A:125:LEU:HB3	1:A:126:PRO:HD3	1.92	0.51
1:B:252:PRO:HG2	1:B:255:MET:HE2	1.93	0.50
1:A:29:THR:HG23	1:A:280:VAL:HG13	1.95	0.49
1:B:210:GLY:HA3	3:B:602:HEM:C3C	2.47	0.49
1:B:125:LEU:HB3	1:B:126:PRO:HD3	1.93	0.49
1:C:125:LEU:HB3	1:C:126:PRO:HD3	1.96	0.48
1:A:123:VAL:HB	1:A:124:PRO:HD3	1.96	0.47
1:D:26:TYR:CG	1:D:291:VAL:HG21	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:500:HEM:C1A	4:D:501:PIM:H4	2.49	0.47
1:D:144:PHE:O	1:D:148:SER:N	2.48	0.47
3:D:500:HEM:HBC2	3:D:500:HEM:CMC	2.44	0.47
1:D:147:TRP:CZ3	1:D:171:LEU:HD11	2.50	0.46
3:A:602:HEM:HBC2	3:A:602:HEM:CMC	2.46	0.45
3:C:500:HEM:HBC2	3:C:500:HEM:CMC	2.46	0.45
1:B:117:ILE:HD11	1:B:362:VAL:HG21	1.97	0.45
1:D:246:GLU:HG3	1:D:299:ILE:O	2.18	0.44
1:B:226:THR:HB	1:B:343:ILE:HD11	1.99	0.43
3:B:602:HEM:HBC2	3:B:602:HEM:CMC	2.44	0.43
3:D:500:HEM:HBB2	3:D:500:HEM:CMB	2.47	0.43
3:B:602:HEM:HBB2	3:B:602:HEM:CMB	2.46	0.43
1:D:108:ASP:OD1	1:D:337:ARG:NH2	2.49	0.43
1:C:217:LEU:HD22	1:C:248:LEU:HD21	1.99	0.43
1:B:244:ILE:CD1	1:B:328:ARG:HA	2.49	0.42
3:C:500:HEM:HBB2	3:C:500:HEM:CMB	2.45	0.42
1:C:282:ILE:HG21	1:C:309:SER:HA	2.00	0.42
1:A:118:VAL:HA	1:A:122:ALA:HB3	2.02	0.42
3:A:602:HEM:HBB2	3:A:602:HEM:CMB	2.46	0.42
1:D:185:VAL:O	1:D:189:VAL:HG23	2.19	0.42
1:D:161:ILE:HA	1:D:164:LEU:HB3	2.02	0.42
1:C:123:VAL:HB	1:C:124:PRO:HD3	2.02	0.41
1:B:223:ILE:HD13	1:B:343:ILE:HD13	2.03	0.41
1:C:344:LEU:HD11	1:C:363:ARG:HB2	2.02	0.41
1:B:320:ALA:N	1:B:321:PRO:CD	2.84	0.41
1:B:123:VAL:HB	1:B:124:PRO:HD3	2.02	0.40
1:D:123:VAL:HB	1:D:124:PRO:HD3	2.02	0.40
1:D:282:ILE:HG21	1:D:309:SER:HA	2.04	0.40
1:D:28:TYR:O	1:D:32:VAL:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/368 (82%)	299 (99%)	4 (1%)	0	100	100
1	B	276/368 (75%)	272 (99%)	4 (1%)	0	100	100
1	C	303/368 (82%)	294 (97%)	8 (3%)	1 (0%)	36	60
1	D	317/368 (86%)	307 (97%)	10 (3%)	0	100	100
All	All	1199/1472 (82%)	1172 (98%)	26 (2%)	1 (0%)	48	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	312	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	242/337 (72%)	241 (100%)	1 (0%)	84	93
1	B	226/337 (67%)	225 (100%)	1 (0%)	84	93
1	C	247/337 (73%)	245 (99%)	2 (1%)	73	88
1	D	266/337 (79%)	263 (99%)	3 (1%)	65	85
All	All	981/1348 (73%)	974 (99%)	7 (1%)	76	90

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	ARG
1	B	359	ARG
1	D	143	LYS
1	D	339	ARG
1	D	360	LEU
1	C	235	ARG
1	C	242	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	211	ASN
1	A	293	HIS
1	B	34	ASN
1	B	293	HIS
1	D	22	GLN
1	D	351	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	B	602	4,1	50,50,50	1.58	8 (16%)	67,82,82	1.49	8 (11%)
4	PIM	B	603	3	11,12,12	1.42	1 (9%)	14,15,15	1.77	4 (28%)
3	HEM	C	500	4,1	50,50,50	1.58	9 (18%)	67,82,82	1.48	8 (11%)
4	PIM	D	501	3	11,12,12	1.46	1 (9%)	14,15,15	1.72	4 (28%)
3	HEM	A	602	5,1	50,50,50	1.56	7 (14%)	67,82,82	1.56	11 (16%)
2	PO4	B	601	-	4,4,4	0.91	0	6,6,6	0.50	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	HEM	D	500	4,1	50,50,50	1.57	6 (12%)	67,82,82	1.61	10 (14%)
4	PIM	C	501	3	11,12,12	1.39	1 (9%)	14,15,15	1.82	3 (21%)
2	PO4	A	601	-	4,4,4	0.93	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	B	602	4,1	-	5/14/54/54	-
4	PIM	B	603	3	-	2/4/4/4	0/2/2/2
3	HEM	C	500	4,1	-	7/14/54/54	-
4	PIM	D	501	3	-	0/4/4/4	0/2/2/2
3	HEM	A	602	5,1	-	2/14/54/54	-
4	PIM	C	501	3	-	0/4/4/4	0/2/2/2
3	HEM	D	500	4,1	-	2/14/54/54	-

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	HEM	FE-NB	5.47	2.11	1.94
3	C	500	HEM	FE-NB	5.30	2.11	1.94
3	A	602	HEM	FE-NB	5.05	2.10	1.94
3	D	500	HEM	FE-NB	4.99	2.10	1.94
3	D	500	HEM	FE-NC	4.68	2.10	1.95
3	B	602	HEM	FE-NC	4.57	2.10	1.95
3	C	500	HEM	FE-NC	4.56	2.10	1.95
3	A	602	HEM	FE-NC	4.51	2.10	1.95
4	D	501	PIM	C5-N1	-4.20	1.33	1.38
4	B	603	PIM	C5-N1	-4.02	1.34	1.38
4	C	501	PIM	C5-N1	-3.86	1.34	1.38
3	A	602	HEM	C1B-NB	-3.64	1.33	1.40
3	D	500	HEM	C1B-NB	-3.59	1.34	1.40
3	C	500	HEM	C1B-NB	-3.26	1.34	1.40
3	B	602	HEM	C1B-NB	-3.22	1.34	1.40
3	C	500	HEM	C4D-ND	-3.17	1.34	1.40
3	B	602	HEM	C4D-ND	-3.01	1.35	1.40
3	D	500	HEM	C4D-ND	-2.94	1.35	1.40
3	A	602	HEM	C4D-ND	-2.73	1.35	1.40
3	A	602	HEM	C4B-NB	-2.40	1.34	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	500	HEM	C4B-NB	-2.36	1.34	1.38
3	C	500	HEM	FE-ND	-2.30	1.87	1.94
3	B	602	HEM	FE-ND	-2.25	1.88	1.94
3	A	602	HEM	C1C-C2C	-2.21	1.41	1.45
3	B	602	HEM	C4B-NB	-2.19	1.34	1.38
3	B	602	HEM	C1C-C2C	-2.16	1.41	1.45
3	C	500	HEM	C4B-NB	-2.16	1.34	1.38
3	D	500	HEM	C1C-C2C	-2.15	1.41	1.45
3	B	602	HEM	C1D-ND	-2.14	1.34	1.38
3	C	500	HEM	C1D-ND	-2.11	1.34	1.38
3	C	500	HEM	C1C-C2C	-2.10	1.41	1.45
3	C	500	HEM	C3B-C4B	2.07	1.48	1.44
3	A	602	HEM	C3B-C4B	2.05	1.48	1.44

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	500	HEM	CHC-C4B-NB	5.00	129.81	124.42
3	B	602	HEM	CHC-C4B-NB	4.98	129.78	124.42
3	A	602	HEM	CHD-C1D-ND	4.85	129.64	124.42
3	B	602	HEM	CHD-C1D-ND	4.58	129.35	124.42
3	D	500	HEM	CHD-C1D-ND	4.54	129.31	124.42
3	D	500	HEM	CHC-C4B-NB	4.54	129.31	124.42
3	D	500	HEM	C1B-NB-C4B	4.39	110.40	105.21
3	A	602	HEM	CHC-C4B-NB	4.37	129.12	124.42
4	C	501	PIM	C6-C5-N1	-4.33	118.68	123.72
4	D	501	PIM	C6-C5-N1	-4.23	118.80	123.72
4	B	603	PIM	C6-C5-N1	-4.14	118.90	123.72
3	C	500	HEM	CHD-C1D-ND	4.08	128.82	124.42
3	D	500	HEM	CHA-C4D-ND	3.89	129.17	124.37
3	A	602	HEM	C1B-NB-C4B	3.83	109.75	105.21
3	A	602	HEM	CHD-C1D-C2D	-3.75	119.11	125.03
3	D	500	HEM	CHD-C1D-C2D	-3.70	119.19	125.03
3	A	602	HEM	CHA-C4D-ND	3.53	128.73	124.37
3	C	500	HEM	CHA-C4D-ND	3.52	128.72	124.37
3	D	500	HEM	CHA-C4D-C3D	-3.29	119.16	125.23
3	B	602	HEM	CHA-C4D-ND	3.15	128.26	124.37
3	B	602	HEM	CHD-C1D-C2D	-3.09	120.16	125.03
4	C	501	PIM	C5-C4-N3	-3.06	106.36	111.04
3	C	500	HEM	C1B-NB-C4B	3.02	108.78	105.21
4	C	501	PIM	C4-C5-N1	2.95	109.83	104.68
4	B	603	PIM	C4-C5-N1	2.93	109.79	104.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	500	HEM	CHB-C1B-NB	2.86	127.91	124.37
4	B	603	PIM	C5-C4-N3	-2.82	106.72	111.04
4	D	501	PIM	C4-C5-N1	2.80	109.56	104.68
3	A	602	HEM	CHA-C4D-C3D	-2.79	120.08	125.23
3	B	602	HEM	C1B-NB-C4B	2.77	108.49	105.21
3	D	500	HEM	C3B-C4B-NB	-2.70	107.53	109.47
3	C	500	HEM	CHD-C1D-C2D	-2.70	120.77	125.03
4	D	501	PIM	C5-C4-N3	-2.53	107.17	111.04
3	B	602	HEM	CHB-C1B-NB	2.47	127.42	124.37
3	C	500	HEM	CHA-C4D-C3D	-2.45	120.72	125.23
3	A	602	HEM	CHB-C1B-NB	2.41	127.35	124.37
3	D	500	HEM	CHB-C1B-NB	2.35	127.28	124.37
3	D	500	HEM	C1A-CHA-C4D	-2.30	120.84	126.25
3	A	602	HEM	C1A-CHA-C4D	-2.26	120.92	126.25
3	C	500	HEM	C4B-C3B-C2B	-2.23	105.23	107.28
4	D	501	PIM	C2-N1-C5	-2.20	106.40	107.53
3	A	602	HEM	C4B-C3B-C2B	-2.18	105.27	107.28
4	B	603	PIM	C2-N1-C5	-2.14	106.43	107.53
3	B	602	HEM	C4B-C3B-C2B	-2.13	105.32	107.28
3	A	602	HEM	CAD-C3D-C4D	2.12	128.40	124.70
3	A	602	HEM	C3B-C4B-NB	-2.12	107.94	109.47
3	B	602	HEM	CHA-C4D-C3D	-2.10	121.36	125.23
3	D	500	HEM	O2D-CGD-CBD	2.01	120.34	114.00

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	500	HEM	C3D-CAD-CBD-CGD
3	B	602	HEM	C2A-CAA-CBA-CGA
4	B	603	PIM	N1-C5-C6-C7
4	B	603	PIM	N1-C5-C6-C11
3	B	602	HEM	CAA-CBA-CGA-O2A
3	B	602	HEM	CAA-CBA-CGA-O1A
3	C	500	HEM	C4D-C3D-CAD-CBD
3	B	602	HEM	CAD-CBD-CGD-O1D
3	B	602	HEM	CAD-CBD-CGD-O2D
3	C	500	HEM	CAD-CBD-CGD-O1D
3	C	500	HEM	CAD-CBD-CGD-O2D
3	A	602	HEM	CAD-CBD-CGD-O2D
3	D	500	HEM	CAD-CBD-CGD-O1D
3	C	500	HEM	CAA-CBA-CGA-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	500	HEM	CAD-CBD-CGD-O2D
3	C	500	HEM	C2D-C3D-CAD-CBD
3	A	602	HEM	CAD-CBD-CGD-O1D
3	C	500	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.


5 monomers are involved in 18 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	HEM	5	0
3	C	500	HEM	4	0
4	D	501	PIM	1	0
3	A	602	HEM	4	0
3	D	500	HEM	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	315/368 (85%)	0.30	18 (5%)	29 26	43, 61, 98, 120	0
1	B	288/368 (78%)	0.49	21 (7%)	21 18	44, 66, 102, 122	0
1	C	311/368 (84%)	0.41	18 (5%)	29 25	51, 71, 89, 115	0
1	D	327/368 (88%)	0.54	19 (5%)	29 25	47, 66, 106, 119	0
All	All	1241/1472 (84%)	0.44	76 (6%)	27 24	43, 66, 101, 122	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	190	VAL	4.1
1	D	152	ALA	3.9
1	B	278	VAL	3.6
1	D	19	ASN	3.6
1	D	272	ILE	3.6
1	C	159	GLY	3.6
1	C	74	PRO	3.6
1	A	139	GLU	3.5
1	C	182	GLY	3.5
1	A	147	TRP	3.5
1	C	334	PHE	3.4
1	A	272	ILE	3.4
1	B	316	LEU	3.1
1	C	181	SER	3.1
1	C	193	ASN	3.0
1	D	138	ILE	3.0
1	A	351	ASN	2.9
1	B	135	GLY	2.9
1	D	81	SER	2.9
1	D	269	ASP	2.9
1	D	73	PRO	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	161	ILE	2.8
1	B	277	TYR	2.7
1	D	181	SER	2.7
1	D	23	VAL	2.7
1	B	138	ILE	2.7
1	A	138	ILE	2.6
1	C	135	GLY	2.5
1	C	162	PHE	2.5
1	A	365	LYS	2.5
1	B	150	LEU	2.5
1	A	185	VAL	2.5
1	C	364	LEU	2.4
1	A	12	ASP	2.4
1	B	269	ASP	2.4
1	B	141	LYS	2.4
1	A	188	ARG	2.3
1	B	258	VAL	2.3
1	B	318	LEU	2.3
1	C	314	ILE	2.3
1	B	262	LYS	2.3
1	C	137	PRO	2.3
1	B	223	ILE	2.3
1	D	140	ASP	2.3
1	B	164	LEU	2.3
1	B	272	ILE	2.2
1	A	350	PRO	2.2
1	D	258	VAL	2.2
1	B	174	TYR	2.2
1	D	22	GLN	2.2
1	B	313	GLY	2.2
1	A	263	GLU	2.2
1	C	202	TYR	2.2
1	D	40	SER	2.2
1	A	338	PHE	2.2
1	B	39	PHE	2.2
1	A	145	LYS	2.1
1	B	199	LYS	2.1
1	C	267	LEU	2.1
1	C	90	GLN	2.1
1	D	267	LEU	2.1
1	A	177	ASP	2.1
1	D	273	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	227	ARG	2.1
1	A	179	LEU	2.1
1	C	315	HIS	2.1
1	A	296	GLU	2.1
1	B	89	PRO	2.1
1	A	341	ILE	2.1
1	D	151	VAL	2.1
1	C	269	ASP	2.0
1	B	137	PRO	2.0
1	C	190	VAL	2.0
1	A	161	ILE	2.0
1	D	264	ARG	2.0
1	C	188	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	B	601	5/5	0.89	0.11	72,72,73,75	0
4	PIM	B	603	11/11	0.90	0.12	61,65,70,70	0
2	PO4	A	601	5/5	0.92	0.14	54,57,58,60	0
4	PIM	D	501	11/11	0.93	0.10	48,49,49,49	0
4	PIM	C	501	11/11	0.93	0.11	51,51,51,52	0
3	HEM	B	602	43/43	0.95	0.10	52,56,67,72	0
3	HEM	D	500	43/43	0.96	0.09	47,53,68,76	0
3	HEM	A	602	43/43	0.96	0.09	42,47,62,70	0
3	HEM	C	500	43/43	0.97	0.08	45,50,60,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM D 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around HEM C 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.