



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 03:00 PM UTC

PDB ID : 7BVR / pdb_00007bvr
Title : DgpB-DgpC complex apo
Authors : Mori, T.; He, H.; Abe, I.
Deposited on : 2020-04-11
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	
1	C	337	
1	E	337	
1	G	337	
2	B	142	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	142	 89% 8% •
2	F	142	 92% 6% ••
2	H	142	 86% 12% •

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 15165 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AP_endonuc_2 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2550	1617	421	485	27	0	0	0
1	C	323	2552	1619	419	487	27	0	0	0
1	E	323	2557	1621	422	487	27	0	0	0
1	G	323	2554	1619	421	487	27	0	0	0

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	325	LYS	-	expression tag	UNP A0A3Q9WXL1
A	326	LEU	-	expression tag	UNP A0A3Q9WXL1
A	327	ALA	-	expression tag	UNP A0A3Q9WXL1
A	328	ALA	-	expression tag	UNP A0A3Q9WXL1
A	329	ALA	-	expression tag	UNP A0A3Q9WXL1
A	330	LEU	-	expression tag	UNP A0A3Q9WXL1
A	331	GLU	-	expression tag	UNP A0A3Q9WXL1
A	332	HIS	-	expression tag	UNP A0A3Q9WXL1
A	333	HIS	-	expression tag	UNP A0A3Q9WXL1
A	334	HIS	-	expression tag	UNP A0A3Q9WXL1
A	335	HIS	-	expression tag	UNP A0A3Q9WXL1
A	336	HIS	-	expression tag	UNP A0A3Q9WXL1
A	337	HIS	-	expression tag	UNP A0A3Q9WXL1
C	325	LYS	-	expression tag	UNP A0A3Q9WXL1
C	326	LEU	-	expression tag	UNP A0A3Q9WXL1
C	327	ALA	-	expression tag	UNP A0A3Q9WXL1
C	328	ALA	-	expression tag	UNP A0A3Q9WXL1
C	329	ALA	-	expression tag	UNP A0A3Q9WXL1
C	330	LEU	-	expression tag	UNP A0A3Q9WXL1
C	331	GLU	-	expression tag	UNP A0A3Q9WXL1
C	332	HIS	-	expression tag	UNP A0A3Q9WXL1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	333	HIS	-	expression tag	UNP A0A3Q9WXL1
C	334	HIS	-	expression tag	UNP A0A3Q9WXL1
C	335	HIS	-	expression tag	UNP A0A3Q9WXL1
C	336	HIS	-	expression tag	UNP A0A3Q9WXL1
C	337	HIS	-	expression tag	UNP A0A3Q9WXL1
E	325	LYS	-	expression tag	UNP A0A3Q9WXL1
E	326	LEU	-	expression tag	UNP A0A3Q9WXL1
E	327	ALA	-	expression tag	UNP A0A3Q9WXL1
E	328	ALA	-	expression tag	UNP A0A3Q9WXL1
E	329	ALA	-	expression tag	UNP A0A3Q9WXL1
E	330	LEU	-	expression tag	UNP A0A3Q9WXL1
E	331	GLU	-	expression tag	UNP A0A3Q9WXL1
E	332	HIS	-	expression tag	UNP A0A3Q9WXL1
E	333	HIS	-	expression tag	UNP A0A3Q9WXL1
E	334	HIS	-	expression tag	UNP A0A3Q9WXL1
E	335	HIS	-	expression tag	UNP A0A3Q9WXL1
E	336	HIS	-	expression tag	UNP A0A3Q9WXL1
E	337	HIS	-	expression tag	UNP A0A3Q9WXL1
G	325	LYS	-	expression tag	UNP A0A3Q9WXL1
G	326	LEU	-	expression tag	UNP A0A3Q9WXL1
G	327	ALA	-	expression tag	UNP A0A3Q9WXL1
G	328	ALA	-	expression tag	UNP A0A3Q9WXL1
G	329	ALA	-	expression tag	UNP A0A3Q9WXL1
G	330	LEU	-	expression tag	UNP A0A3Q9WXL1
G	331	GLU	-	expression tag	UNP A0A3Q9WXL1
G	332	HIS	-	expression tag	UNP A0A3Q9WXL1
G	333	HIS	-	expression tag	UNP A0A3Q9WXL1
G	334	HIS	-	expression tag	UNP A0A3Q9WXL1
G	335	HIS	-	expression tag	UNP A0A3Q9WXL1
G	336	HIS	-	expression tag	UNP A0A3Q9WXL1
G	337	HIS	-	expression tag	UNP A0A3Q9WXL1

- Molecule 2 is a protein called DgpB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	B	139	Total	C	N	O	S	0	0	0
			1111	714	179	214	4			
2	D	138	Total	C	N	O	S	0	0	0
			1103	708	178	213	4			
2	F	140	Total	C	N	O	S	0	0	0
			1111	714	181	212	4			
2	H	139	Total	C	N	O	S	0	0	0
			1111	714	179	214	4			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	E	1	Total C O 6 3 3	0	0
3	G	1	Total C O 6 3 3	0	0

- Molecule 4 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mn 1 1	0	0
4	C	1	Total Mn 1 1	0	0
4	E	1	Total Mn 1 1	0	0
4	G	1	Total Mn 1 1	0	0

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
5	B	1	Total 15	8	2	4	1	0	0
5	B	1	Total 15	8	2	4	1	0	0
5	D	1	Total 15	8	2	4	1	0	0
5	F	1	Total 15	8	2	4	1	0	0
5	H	1	Total 15	8	2	4	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	44	Total 44	44	0	0
6	B	57	Total 57	57	0	0
6	C	68	Total 68	68	0	0
6	D	35	Total 35	35	0	0
6	E	79	Total 79	79	0	0
6	F	49	Total 49	49	0	0
6	G	39	Total 39	39	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	42	Total	O	0	0
			42	42		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: AP_endonuc_2 domain-containing protein




- Molecule 1: AP_endonuc_2 domain-containing protein




- Molecule 1: AP_endonuc_2 domain-containing protein




- Molecule 1: AP_endonuc_2 domain-containing protein







• Molecule 2: DgpB




• Molecule 2: DgpB



• Molecule 2: DgpB



• Molecule 2: DgpB



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.52Å 156.72Å 158.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.89 – 2.60 46.89 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.89-2.60) 99.9 (46.89-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.191 , 0.249 0.191 , 0.250	Depositor DCC
R_{free} test set	2008 reflections (2.96%)	wwPDB-VP
Wilson B-factor (Å ²)	38.1	Xtrriage
Anisotropy	0.176	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15165	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.5972e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EPE, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2606	0.35	0/3517
1	C	0.13	0/2608	0.35	0/3519
1	E	0.13	0/2613	0.36	0/3525
1	G	0.12	0/2610	0.33	0/3522
2	B	0.18	0/1137	0.35	0/1543
2	D	0.13	0/1129	0.35	0/1532
2	F	0.26	0/1137	0.42	0/1544
2	H	0.16	0/1137	0.32	0/1543
All	All	0.15	0/14977	0.35	0/20245

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2550	0	2471	45	0
1	C	2552	0	2475	29	0
1	E	2557	0	2484	25	0
1	G	2554	0	2475	31	0
2	B	1111	0	1072	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1103	0	1061	9	0
2	F	1111	0	1070	6	0
2	H	1111	0	1072	10	0
3	A	6	0	8	0	0
3	C	6	0	8	2	0
3	E	6	0	8	0	0
3	G	6	0	8	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	G	1	0	0	0	0
5	B	30	0	34	2	0
5	D	15	0	17	0	0
5	F	15	0	17	0	0
5	H	15	0	17	0	0
6	A	44	0	0	0	0
6	B	57	0	0	0	0
6	C	68	0	0	1	0
6	D	35	0	0	0	0
6	E	79	0	0	1	0
6	F	49	0	0	0	0
6	G	39	0	0	0	0
6	H	42	0	0	0	0
All	All	15165	0	14297	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:303:TYR:CE2	2:F:7:LEU:HD21	2.12	0.84
1:A:269:MET:H	1:A:311:MET:HE3	1.49	0.78
1:C:172:PRO:HG2	1:C:262:MET:HG2	1.65	0.77
2:B:49:GLU:HA	5:B:201:EPE:H101	1.68	0.76
2:D:83:ASN:HD21	2:H:83:ASN:HB3	1.47	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/337 (95%)	313 (98%)	8 (2%)	0	100	100
1	C	321/337 (95%)	312 (97%)	9 (3%)	0	100	100
1	E	321/337 (95%)	311 (97%)	10 (3%)	0	100	100
1	G	321/337 (95%)	309 (96%)	12 (4%)	0	100	100
2	B	137/142 (96%)	133 (97%)	4 (3%)	0	100	100
2	D	136/142 (96%)	131 (96%)	5 (4%)	0	100	100
2	F	138/142 (97%)	133 (96%)	5 (4%)	0	100	100
2	H	137/142 (96%)	135 (98%)	2 (2%)	0	100	100
All	All	1832/1916 (96%)	1777 (97%)	55 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/287 (94%)	271 (100%)	0	100	100
1	C	272/287 (95%)	272 (100%)	0	100	100
1	E	273/287 (95%)	273 (100%)	0	100	100
1	G	272/287 (95%)	272 (100%)	0	100	100
2	B	123/125 (98%)	123 (100%)	0	100	100
2	D	122/125 (98%)	122 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	122/125 (98%)	121 (99%)	1 (1%)	73	88
2	H	123/125 (98%)	123 (100%)	0	100	100
All	All	1578/1648 (96%)	1577 (100%)	1 (0%)	88	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	7	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
2	F	31	GLN
1	G	19	GLN
2	H	141	GLN
2	H	8	ASN
2	H	31	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EPE	B	201	-	15,15,15	0.81	1 (6%)	19,20,20	1.86	5 (26%)
3	GOL	A	401	-	5,5,5	0.94	0	5,5,5	1.03	0
3	GOL	E	401	-	5,5,5	0.81	0	5,5,5	1.07	0
5	EPE	H	201	-	15,15,15	0.80	1 (6%)	19,20,20	1.96	4 (21%)
3	GOL	C	401	-	5,5,5	0.90	0	5,5,5	1.05	0
5	EPE	B	202	-	15,15,15	0.86	1 (6%)	19,20,20	1.74	4 (21%)
5	EPE	F	201	-	15,15,15	0.83	1 (6%)	19,20,20	1.86	6 (31%)
3	GOL	G	401	-	5,5,5	0.83	0	5,5,5	1.12	0
5	EPE	D	201	-	15,15,15	0.81	1 (6%)	19,20,20	1.90	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	B	201	-	-	7/9/19/19	0/1/1/1
3	GOL	A	401	-	-	1/4/4/4	-
3	GOL	E	401	-	-	2/4/4/4	-
5	EPE	H	201	-	-	4/9/19/19	0/1/1/1
3	GOL	C	401	-	-	0/4/4/4	-
5	EPE	B	202	-	-	3/9/19/19	0/1/1/1
5	EPE	F	201	-	-	4/9/19/19	0/1/1/1
3	GOL	G	401	-	-	4/4/4/4	-
5	EPE	D	201	-	-	2/9/19/19	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	202	EPE	C10-S	2.92	1.81	1.77
5	F	201	EPE	C10-S	2.73	1.81	1.77
5	D	201	EPE	C10-S	2.69	1.81	1.77
5	H	201	EPE	C10-S	2.69	1.81	1.77
5	B	201	EPE	C10-S	2.63	1.81	1.77

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	201	EPE	C5-N4-C3	5.70	121.11	108.84
5	B	201	EPE	C5-N4-C3	4.64	118.84	108.84
5	B	202	EPE	C5-N4-C3	4.54	118.62	108.84
5	D	201	EPE	C5-N4-C3	4.40	118.31	108.84
5	F	201	EPE	C5-N4-C3	4.24	117.98	108.84

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	401	GOL	C1-C2-C3-O3
5	B	201	EPE	C8-C7-N4-C5
5	B	201	EPE	C9-C10-S-O1S
5	B	201	EPE	C9-C10-S-O3S
5	B	202	EPE	C9-C10-S-O2S

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	201	EPE	2	0
3	C	401	GOL	2	0
3	G	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/337 (95%)	-0.17	2 (0%) 85 83	25, 43, 60, 76	0
1	C	323/337 (95%)	-0.12	4 (1%) 76 73	24, 40, 76, 89	0
1	E	323/337 (95%)	-0.36	1 (0%) 90 88	22, 33, 55, 66	0
1	G	323/337 (95%)	-0.10	3 (0%) 81 78	26, 44, 70, 86	0
2	B	139/142 (97%)	-0.47	0 100 100	25, 33, 43, 58	0
2	D	138/142 (97%)	-0.31	0 100 100	25, 38, 52, 58	0
2	F	140/142 (98%)	-0.34	2 (1%) 73 69	23, 35, 49, 60	0
2	H	139/142 (97%)	-0.46	1 (0%) 84 82	26, 34, 50, 61	0
All	All	1848/1916 (96%)	-0.25	13 (0%) 84 82	22, 37, 62, 89	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	166	LYS	2.8
1	A	324	ASP	2.7
1	C	226	LYS	2.6
1	G	143	HIS	2.5
2	F	142	ASN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	G	401	6/6	0.82	0.14	39,41,42,49	0
3	GOL	A	401	6/6	0.85	0.15	37,41,41,47	0
5	EPE	B	201	15/15	0.85	0.17	40,48,68,73	0
3	GOL	E	401	6/6	0.86	0.14	34,35,37,37	0
5	EPE	D	201	15/15	0.91	0.15	42,57,73,74	0
5	EPE	H	201	15/15	0.91	0.16	35,60,82,87	0
5	EPE	F	201	15/15	0.92	0.14	38,50,70,71	0
3	GOL	C	401	6/6	0.92	0.10	38,39,43,46	0
5	EPE	B	202	15/15	0.94	0.13	37,59,77,77	0
4	MN	G	402	1/1	0.98	0.12	32,32,32,32	1
4	MN	C	402	1/1	0.98	0.10	29,29,29,29	1
4	MN	E	402	1/1	0.98	0.10	19,19,19,19	1
4	MN	A	402	1/1	1.00	0.11	30,30,30,30	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MN G 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN C 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN E 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.