



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 08:47 PM UTC

PDB ID : 8BV1 / pdb_00008bv1
Title : Peptide inhibitor P4 in complex with ASF1 histone chaperone
Authors : Perrin, M.E.; Li, B.; Mbianda, J.; Ropars, V.; Legrand, P.; Douat, C.; Ochsenbein, F.; Guichard, G.
Deposited on : 2022-12-01
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	2% 91% 8% .
1	B	158	% 90% 8% ..
1	C	158	2% 89% 7% ..
1	D	158	2% 95% . .
1	E	158	3% 92% 6% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	158	3% 89% 9%
2	G	13	85% 15%
2	H	13	85% 15%
2	I	13	92% 8%
2	K	13	62% 38%
2	L	13	77% 23%
2	N	13	69% 31%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone chaperone ASF1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	155	Total	C	N	O	S	0	0	0
			1248	800	203	241	4			
1	B	155	Total	C	N	O	S	0	0	0
			1248	799	204	242	3			
1	C	152	Total	C	N	O	S	0	0	0
			1223	786	200	234	3			
1	D	154	Total	C	N	O	S	0	0	0
			1240	795	202	240	3			
1	E	154	Total	C	N	O	S	0	0	0
			1240	795	202	240	3			
1	F	154	Total	C	N	O	S	0	0	0
			1240	795	202	240	3			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9Y294
A	0	ALA	-	expression tag	UNP Q9Y294
B	-1	GLY	-	expression tag	UNP Q9Y294
B	0	ALA	-	expression tag	UNP Q9Y294
C	-1	GLY	-	expression tag	UNP Q9Y294
C	0	ALA	-	expression tag	UNP Q9Y294
D	-1	GLY	-	expression tag	UNP Q9Y294
D	0	ALA	-	expression tag	UNP Q9Y294
E	-1	GLY	-	expression tag	UNP Q9Y294
E	0	ALA	-	expression tag	UNP Q9Y294
F	-1	GLY	-	expression tag	UNP Q9Y294
F	0	ALA	-	expression tag	UNP Q9Y294

- Molecule 2 is a protein called P4 peptide inhibitor of histone chaperone ASF1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	N	13	Total	C	N	O	0	0	1
			101	65	22	14			
2	G	13	Total	C	N	O	0	0	1
			101	65	22	14			
2	K	13	Total	C	N	O	0	0	1
			101	65	22	14			
2	H	13	Total	C	N	O	0	0	1
			101	65	22	14			
2	L	13	Total	C	N	O	0	0	1
			101	65	22	14			
2	I	13	Total	C	N	O	0	0	1
			101	65	22	14			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	29	Total	O	0	0
			29	29		
4	C	19	Total	O	0	0
			19	19		
4	D	21	Total	O	0	0
			21	21		
4	E	18	Total	O	0	0
			18	18		
4	F	12	Total	O	0	0
			12	12		
4	N	4	Total	O	0	0
			4	4		
4	G	4	Total	O	0	0
			4	4		
4	H	1	Total	O	0	0
			1	1		
4	L	4	Total	O	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone chaperone ASF1A



- Molecule 1: Histone chaperone ASF1A




- Molecule 1: Histone chaperone ASF1A





- Molecule 1: Histone chaperone ASF1A




- Molecule 1: Histone chaperone ASF1A





- Molecule 1: Histone chaperone ASF1A



- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1





- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1





- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1





- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1



- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1



- Molecule 2: P4 peptide inhibitor of histone chaperone ASF1



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	123.89Å 123.89Å 179.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.79 – 2.83 29.79 – 2.83	Depositor EDS
% Data completeness (in resolution range)	88.8 (29.79-2.83) 88.7 (29.79-2.83)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.85Å)	Xtriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.204 , 0.221 0.193 , 0.212	Depositor DCC
R_{free} test set	2830 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	91.2	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 60.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8282	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, GOL, NH2, ALN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/1281	0.99	1/1752 (0.1%)
1	B	0.76	0/1281	1.00	0/1753
1	C	0.69	0/1256	0.95	2/1719 (0.1%)
1	D	0.66	0/1273	0.93	2/1742 (0.1%)
1	E	0.71	0/1273	0.97	2/1742 (0.1%)
1	F	0.67	0/1273	0.94	1/1742 (0.1%)
2	G	0.85	0/81	1.02	1/104 (1.0%)
2	H	0.63	0/81	0.80	0/104
2	I	0.68	0/81	0.94	0/104
2	K	0.73	0/81	1.06	0/104
2	L	0.85	0/81	0.87	0/104
2	N	0.74	0/81	1.00	0/104
All	All	0.70	0/8123	0.96	9/11074 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	13	ASP	CA-CB-CG	5.54	118.14	112.60
1	E	155	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	13	ASP	CA-CB-CG	5.29	117.89	112.60
1	F	86	ASP	CA-CB-CG	5.27	117.87	112.60
1	C	13	ASP	CA-CB-CG	5.18	117.78	112.60
1	D	13	ASP	CA-CB-CG	5.13	117.73	112.60
1	C	116	GLU	CB-CG-CD	5.08	121.24	112.60
2	G	11	ILE	N-CA-C	-5.07	106.41	113.00
1	D	86	ASP	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1248	0	1198	8	0
1	B	1248	0	1192	10	0
1	C	1223	0	1176	6	0
1	D	1240	0	1186	4	0
1	E	1240	0	1186	3	0
1	F	1240	0	1186	8	0
2	G	101	0	108	2	0
2	H	101	0	108	1	0
2	I	101	0	108	1	0
2	K	101	0	108	6	0
2	L	101	0	108	2	0
2	N	101	0	108	2	0
3	A	36	0	48	3	0
3	B	18	0	24	1	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
3	E	18	0	24	0	0
3	F	6	0	8	0	0
4	A	30	0	0	0	0
4	B	29	0	0	0	0
4	C	19	0	0	0	0
4	D	21	0	0	0	0
4	E	18	0	0	0	0
4	F	12	0	0	0	0
4	G	4	0	0	0	0
4	H	1	0	0	0	0
4	I	5	0	0	0	0
4	L	4	0	0	0	0
4	N	4	0	0	0	0
All	All	8282	0	7892	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:ASN:HD21	1:B:139:ILE:H	1.32	0.77
1:F:113:VAL:HG22	1:F:139:ILE:HG12	1.80	0.64
2:N:1:ACE:H2	2:N:5:ALA:H	1.62	0.63
1:B:145:ARG:HD3	2:K:11:ILE:HG21	1.81	0.63
1:E:36:GLU:HG3	1:E:102:ARG:HD3	1.83	0.60
1:E:10:VAL:HB	1:E:25:GLU:HB3	1.86	0.56
2:K:2:GLU:O	2:K:6:ARG:HG3	2.08	0.53
1:B:90:VAL:HG22	1:B:135:LEU:HD21	1.91	0.51
1:A:4:VAL:HG21	1:A:38:LEU:HD21	1.92	0.51
1:A:69:ARG:NH2	3:A:203:GOL:O1	2.45	0.49
1:B:146:VAL:HG21	3:B:202:GOL:H31	1.94	0.49
1:D:145:ARG:HD2	2:K:1:ACE:H3	1.95	0.47
1:C:116:GLU:OE2	1:C:138:ASN:HB2	2.15	0.47
1:A:146:VAL:HG21	3:A:201:GOL:H2	1.98	0.46
1:B:54:ASP:OD2	2:K:10:ARG:NH2	2.49	0.46
1:C:113:VAL:HG22	1:C:139:ILE:HG12	1.97	0.46
1:A:33:ASP:CG	1:B:145:ARG:HH22	2.25	0.45
1:F:102:ARG:NH2	1:F:155:ASP:O	2.41	0.45
1:F:145:ARG:HD3	2:G:1:ACE:H3	1.99	0.45
1:B:30:CYS:SG	1:B:32:GLU:O	2.76	0.44
1:C:90:VAL:HG22	1:C:135:LEU:HD21	2.00	0.43
1:C:145:ARG:HB3	2:I:11:ILE:HG21	2.01	0.43
1:B:138:ASN:HD22	1:D:138:ASN:HD22	1.65	0.43
1:A:63:GLY:HA2	1:A:64:PRO:C	2.44	0.42
2:N:6:ARG:O	2:N:10:ARG:HG2	2.20	0.42
1:C:30:CYS:HB3	1:C:67:ALA:HA	2.01	0.42
2:L:6:ARG:O	2:L:10:ARG:HG2	2.20	0.42
1:A:64:PRO:HG3	2:L:9:ARG:HB2	2.01	0.42
2:K:6:ARG:O	2:K:10:ARG:HG2	2.20	0.42
1:F:4:VAL:HG21	1:F:38:LEU:HD11	2.03	0.41
1:F:90:VAL:HG22	1:F:135:LEU:HD21	2.03	0.41
1:F:145:ARG:CD	2:G:1:ACE:H3	2.50	0.41
1:F:39:GLU:HB3	1:F:100:THR:HB	2.01	0.41
1:A:101:TYR:CD2	1:A:102:ARG:HG3	2.56	0.41
2:H:6:ARG:O	2:H:10:ARG:HG2	2.21	0.41
1:A:150:HIS:HB3	3:A:204:GOL:H12	2.02	0.41
1:B:10:VAL:HB	1:B:25:GLU:HB3	2.02	0.41
1:E:4:VAL:CG2	1:E:34:LEU:HD11	2.51	0.41
1:B:138:ASN:ND2	1:D:138:ASN:HD22	2.20	0.40
1:D:145:ARG:NE	2:K:1:ACE:CH3	2.85	0.40
1:F:10:VAL:HB	1:F:25:GLU:HB3	2.04	0.40
1:C:126:PRO:HA	1:C:127:PRO:HD3	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	153/158 (97%)	153 (100%)	0	0	100	100
1	B	153/158 (97%)	152 (99%)	1 (1%)	0	100	100
1	C	150/158 (95%)	149 (99%)	1 (1%)	0	100	100
1	D	152/158 (96%)	151 (99%)	1 (1%)	0	100	100
1	E	152/158 (96%)	151 (99%)	1 (1%)	0	100	100
1	F	152/158 (96%)	150 (99%)	2 (1%)	0	100	100
2	G	10/13 (77%)	10 (100%)	0	0	100	100
2	H	10/13 (77%)	10 (100%)	0	0	100	100
2	I	10/13 (77%)	10 (100%)	0	0	100	100
2	K	10/13 (77%)	10 (100%)	0	0	100	100
2	L	10/13 (77%)	10 (100%)	0	0	100	100
2	N	10/13 (77%)	10 (100%)	0	0	100	100
All	All	972/1026 (95%)	966 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/141 (99%)	139 (99%)	1 (1%)	76	88
1	B	140/141 (99%)	138 (99%)	2 (1%)	59	78
1	C	137/141 (97%)	137 (100%)	0	100	100
1	D	139/141 (99%)	139 (100%)	0	100	100
1	E	139/141 (99%)	138 (99%)	1 (1%)	76	88
1	F	139/141 (99%)	139 (100%)	0	100	100
2	G	7/7 (100%)	7 (100%)	0	100	100
2	H	7/7 (100%)	7 (100%)	0	100	100
2	I	7/7 (100%)	7 (100%)	0	100	100
2	K	7/7 (100%)	7 (100%)	0	100	100
2	L	7/7 (100%)	7 (100%)	0	100	100
2	N	7/7 (100%)	7 (100%)	0	100	100
All	All	876/888 (99%)	872 (100%)	4 (0%)	81	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ILE
1	B	104	GLN
1	B	145	ARG
1	E	140	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
1	A	152	ASN
1	B	14	ASN
1	B	114	ASN
1	D	5	GLN
1	D	8	ASN
1	D	138	ASN
1	E	114	ASN
1	F	104	GLN
1	F	114	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALN	I	4	2	15,16,17	0.21	0	16,21,23	0.19	0
2	ALN	H	4	2	15,16,17	0.20	0	16,21,23	0.22	0
2	ALN	N	4	2	15,16,17	0.22	0	16,21,23	0.24	0
2	ALN	K	4	2	15,16,17	0.30	0	16,21,23	0.24	0
2	ALN	L	4	2	15,16,17	0.25	0	16,21,23	0.26	0
2	ALN	G	4	2	15,16,17	0.24	0	16,21,23	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALN	I	4	2	-	0/5/6/8	0/2/2/2
2	ALN	H	4	2	-	0/5/6/8	0/2/2/2
2	ALN	N	4	2	-	0/5/6/8	0/2/2/2
2	ALN	K	4	2	-	0/5/6/8	0/2/2/2
2	ALN	L	4	2	-	0/5/6/8	0/2/2/2
2	ALN	G	4	2	-	0/5/6/8	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	F	201	-	5,5,5	0.17	0	5,5,5	0.31	0
3	GOL	A	205	-	5,5,5	0.18	0	5,5,5	0.35	0
3	GOL	E	201	-	5,5,5	0.13	0	5,5,5	0.34	0
3	GOL	D	201	-	5,5,5	0.06	0	5,5,5	0.26	0
3	GOL	A	204	-	5,5,5	0.18	0	5,5,5	0.44	0
3	GOL	B	203	-	5,5,5	0.10	0	5,5,5	0.30	0
3	GOL	E	203	-	5,5,5	0.12	0	5,5,5	0.25	0
3	GOL	E	202	-	5,5,5	0.10	0	5,5,5	0.27	0
3	GOL	A	202	-	5,5,5	0.34	0	5,5,5	0.38	0
3	GOL	A	206	-	5,5,5	0.06	0	5,5,5	0.28	0
3	GOL	A	201	-	5,5,5	0.20	0	5,5,5	0.37	0
3	GOL	B	201	-	5,5,5	0.14	0	5,5,5	0.39	0
3	GOL	B	202	-	5,5,5	0.10	0	5,5,5	0.33	0
3	GOL	C	201	-	5,5,5	0.17	0	5,5,5	0.39	0
3	GOL	A	203	-	5,5,5	0.10	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	F	201	-	-	0/4/4/4	-
3	GOL	A	205	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	E	201	-	-	1/4/4/4	-
3	GOL	D	201	-	-	0/4/4/4	-
3	GOL	A	204	-	-	0/4/4/4	-
3	GOL	B	203	-	-	0/4/4/4	-
3	GOL	E	203	-	-	0/4/4/4	-
3	GOL	E	202	-	-	1/4/4/4	-
3	GOL	A	202	-	-	0/4/4/4	-
3	GOL	A	206	-	-	0/4/4/4	-
3	GOL	A	201	-	-	1/4/4/4	-
3	GOL	B	201	-	-	0/4/4/4	-
3	GOL	B	202	-	-	0/4/4/4	-
3	GOL	C	201	-	-	2/4/4/4	-
3	GOL	A	203	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	201	GOL	C1-C2-C3-O3
3	A	205	GOL	O1-C1-C2-C3
3	A	205	GOL	O1-C1-C2-O2
3	C	201	GOL	O2-C2-C3-O3
3	A	201	GOL	O2-C2-C3-O3
3	E	201	GOL	O2-C2-C3-O3
3	E	202	GOL	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	204	GOL	1	0
3	A	201	GOL	1	0
3	B	202	GOL	1	0
3	A	203	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	155/158 (98%)	-0.22	3 (1%) 66 59	54, 80, 109, 121	1 (0%)
1	B	155/158 (98%)	-0.08	2 (1%) 75 69	54, 85, 112, 124	1 (0%)
1	C	152/158 (96%)	0.13	3 (1%) 65 57	56, 100, 146, 161	1 (0%)
1	D	154/158 (97%)	0.05	3 (1%) 66 59	52, 97, 154, 164	2 (1%)
1	E	154/158 (97%)	0.06	5 (3%) 50 40	71, 99, 137, 146	0
1	F	154/158 (97%)	0.31	5 (3%) 50 40	75, 111, 173, 182	0
2	G	10/13 (76%)	-0.24	0 100 100	77, 80, 84, 94	0
2	H	10/13 (76%)	0.01	0 100 100	84, 92, 100, 103	0
2	I	10/13 (76%)	-0.02	0 100 100	79, 83, 91, 96	0
2	K	10/13 (76%)	-0.05	0 100 100	72, 75, 88, 92	0
2	L	10/13 (76%)	-0.34	0 100 100	69, 73, 82, 88	0
2	N	10/13 (76%)	-0.31	0 100 100	73, 75, 85, 97	0
All	All	984/1026 (95%)	0.03	21 (2%) 63 55	52, 92, 150, 182	5 (0%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	155	ASP	4.4
1	C	31	ILE	3.6
1	F	66	PRO	3.6
1	A	155	ASP	3.5
1	D	2	ALA	3.4
1	F	64	PRO	3.3
1	B	156	ASN	3.3
1	C	2	ALA	3.2
1	F	155	ASP	3.2
1	B	2	ALA	3.2
1	A	1	MET	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	31	ILE	3.0
1	E	116	GLU	2.7
1	E	31	ILE	2.5
1	F	2	ALA	2.4
1	D	3	LYS	2.4
1	E	2	ALA	2.4
1	D	155	ASP	2.2
1	E	154	GLU	2.2
1	A	123	ARG	2.1
1	C	123	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ALN	I	4	15/16	0.96	0.10	79,79,80,80	0
2	ALN	G	4	15/16	0.97	0.09	81,83,83,83	0
2	ALN	K	4	15/16	0.97	0.09	72,73,76,77	0
2	ALN	N	4	15/16	0.97	0.10	78,78,79,80	0
2	ALN	L	4	15/16	0.98	0.07	71,75,76,76	0
2	ALN	H	4	15/16	0.98	0.09	82,83,84,84	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	E	202	6/6	0.61	0.28	104,105,105,105	0
3	GOL	A	204	6/6	0.79	0.18	107,108,109,109	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	206	6/6	0.81	0.14	118,118,119,119	0
3	GOL	E	201	6/6	0.82	0.21	87,88,88,89	0
3	GOL	A	203	6/6	0.84	0.21	116,116,116,117	0
3	GOL	C	201	6/6	0.86	0.24	100,100,101,101	0
3	GOL	F	201	6/6	0.87	0.22	104,106,107,107	0
3	GOL	A	201	6/6	0.88	0.20	87,88,90,91	0
3	GOL	B	202	6/6	0.88	0.20	89,90,90,90	0
3	GOL	A	205	6/6	0.88	0.15	98,99,99,100	0
3	GOL	A	202	6/6	0.89	0.14	86,87,89,91	0
3	GOL	B	203	6/6	0.89	0.16	119,119,119,119	0
3	GOL	B	201	6/6	0.89	0.15	110,110,111,111	0
3	GOL	E	203	6/6	0.93	0.16	119,120,121,121	0
3	GOL	D	201	6/6	0.94	0.18	94,94,95,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.