



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:55 AM UTC

PDB ID : 3BWU / pdb_00003bwu
Title : Crystal structure of the ternary complex of FimD (N-Terminal Domain, FimDN) with FimC and the N-terminally truncated pilus subunit FimF (FimFt)
Authors : Eidam, O.; Grutter, M.G.; Capitani, G.
Deposited on : 2008-01-10
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	205	3% 87% 10% .
2	D	125	8% 86% 10% .
3	F	142	12% 78% 12% 10%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperone protein fimC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	200	Total	C	N	O	S	4	14	0
			1621	1030	276	307	8			

- Molecule 2 is a protein called Outer membrane usher protein FimD, N-terminal domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	121	Total	C	N	O	S	0	6	0
			938	587	157	186	8			

- Molecule 3 is a protein called Protein fimF.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	128	Total	C	N	O	S	12	5	0
			951	601	159	188	3			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	301	Total	O	0	0
			301	301		
6	D	173	Total	O	0	0
			173	173		
6	F	167	Total	O	0	0
			167	167		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chaperone protein fimC

Chain C: 




- Molecule 2: Outer membrane usher protein FimD, N-terminal domain

Chain D: 



- Molecule 3: Protein fimF

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	126.95Å 126.95Å 68.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.34 – 1.76 32.34 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.9 (32.34-1.76) 99.9 (32.34-1.76)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.49 (at 1.76Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.175 , 0.212 0.172 , 0.211	Depositor DCC
R_{free} test set	1285 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å ²)	15.3	Xtriage
Anisotropy	0.225	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4210	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.35	0/1695	0.70	0/2301
2	D	0.32	0/972	0.65	0/1326
3	F	0.32	0/983	0.71	0/1347
All	All	0.34	0/3650	0.69	0/4974

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1621	0	1683	20	0
2	D	938	0	909	10	0
3	F	951	0	953	16	0
4	C	32	0	48	6	0
4	D	8	0	12	3	0
4	F	12	0	18	3	0
5	C	7	0	10	3	0
6	C	301	0	0	10	0
6	D	173	0	0	1	0
6	F	167	0	0	5	0
All	All	4210	0	3633	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:200[B]:MET:HE3	6:C:463:HOH:O	1.82	0.80
2:D:94:THR:HG23	4:D:127:EDO:H12	1.65	0.79
1:C:34:GLN:HG3	4:C:213:EDO:H12	1.72	0.72
2:D:109[B]:GLN:NE2	6:D:177:HOH:O	2.29	0.64
2:D:93:LEU:HB3	4:D:127:EDO:H11	1.81	0.63
1:C:158:GLY:HA2	1:C:184[B]:ASN:HD21	1.66	0.60
1:C:165:ALA:C	4:C:208:EDO:H22	2.27	0.59
3:F:128:TYR:OH	4:F:155:EDO:H22	2.03	0.58
3:F:77:ASN:HB2	6:F:192:HOH:O	2.02	0.58
4:C:212:EDO:H12	6:C:227:HOH:O	2.03	0.57
2:D:79:VAL:HG11	2:D:82:MET:HE3	1.87	0.57
1:C:16:LYS:HG2	6:C:358:HOH:O	2.06	0.56
1:C:129:GLU:HG2	1:C:200[B]:MET:HE1	1.87	0.55
1:C:129:GLU:HA	1:C:200[B]:MET:HE1	1.88	0.54
3:F:91:LEU:HD23	3:F:133:ALA:HA	1.89	0.54
1:C:158:GLY:HA2	1:C:184[B]:ASN:ND2	2.24	0.53
1:C:93:MET:HB3	2:D:3:TYR:HD2	1.74	0.53
1:C:27:GLU:HG2	6:C:475:HOH:O	2.09	0.52
2:D:94:THR:H	4:D:127:EDO:C1	2.23	0.52
3:F:124:THR:HG21	6:F:185:HOH:O	2.10	0.51
1:C:37:VAL:HG23	1:C:50[A]:VAL:CG2	2.41	0.50
4:C:207:EDO:H21	5:C:214:PEG:H32	1.94	0.49
1:C:23[B]:THR:HG23	6:C:296:HOH:O	2.13	0.49
1:C:19[A]:GLN:NE2	6:C:304:HOH:O	2.46	0.49
1:C:200[B]:MET:HE2	6:C:278:HOH:O	2.13	0.49
6:C:450:HOH:O	3:F:55:PRO:HG3	2.12	0.48
3:F:104:PRO:HB3	4:F:5:EDO:H21	1.95	0.48
3:F:128:TYR:CZ	4:F:155:EDO:H11	2.48	0.48
1:C:101:ASN:HA	3:F:143:ILE:O	2.14	0.47
4:C:209:EDO:H21	6:F:312:HOH:O	2.14	0.47
1:C:36:TRP:CE2	5:C:214:PEG:H31	2.51	0.46
1:C:93:MET:HE2	1:C:104:GLN:OE1	2.16	0.46
4:C:210:EDO:H11	6:C:383:HOH:O	2.14	0.45
1:C:157:ALA:HA	1:C:185[B]:ILE:HD13	1.99	0.45
2:D:2:LEU:HD23	2:D:2:LEU:HA	1.89	0.43
1:C:204:MET:HG3	6:C:351:HOH:O	2.18	0.43
3:F:13:ASP:N	6:F:180:HOH:O	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:44:LYS:NZ	5:C:214:PEG:H42	2.34	0.43
3:F:96[A]:LEU:HD23	3:F:102:GLN:HA	1.99	0.42
3:F:104:PRO:HB2	3:F:107:ALA:HB2	2.01	0.42
3:F:13:ASP:N	6:F:216:HOH:O	2.53	0.42
2:D:98:GLN:O	2:D:99:ASP:HB2	2.20	0.41
3:F:95:LEU:O	3:F:96[B]:LEU:HD13	2.21	0.41
2:D:5:ASN:HA	2:D:6:PRO:HD3	1.93	0.41
3:F:40:ILE:HG13	3:F:136:VAL:HG22	2.02	0.41
2:D:92:PRO:HD2	2:D:96:MET:SD	2.61	0.41
3:F:65:VAL:HB	3:F:149:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	210/205 (102%)	210 (100%)	0	0	100	100
2	D	125/125 (100%)	124 (99%)	1 (1%)	0	100	100
3	F	129/142 (91%)	127 (98%)	2 (2%)	0	100	100
All	All	464/472 (98%)	461 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	185/176 (105%)	184 (100%)	1 (0%)	81	75
2	D	103/103 (100%)	103 (100%)	0	100	100
3	F	106/114 (93%)	106 (100%)	0	100	100
All	All	394/393 (100%)	393 (100%)	1 (0%)	86	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	8	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	164	ASN
2	D	52	ASN
2	D	75	ASN
2	D	122	ASN
3	F	101	ASN
3	F	135	GLN
3	F	144	ASN
3	F	154	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	C	209	-	3,3,3	0.45	0	2,2,2	0.31	0
4	EDO	C	206	-	3,3,3	0.39	0	2,2,2	0.44	0
4	EDO	F	9	-	3,3,3	0.38	0	2,2,2	0.63	0
4	EDO	C	210	-	3,3,3	0.42	0	2,2,2	0.32	0
4	EDO	C	211	-	3,3,3	0.43	0	2,2,2	0.49	0
4	EDO	C	212	-	3,3,3	0.43	0	2,2,2	0.35	0
4	EDO	D	126	-	3,3,3	0.43	0	2,2,2	0.28	0
5	PEG	C	214	-	6,6,6	0.55	0	5,5,5	0.45	0
4	EDO	F	5	-	3,3,3	0.43	0	2,2,2	0.37	0
4	EDO	C	213	-	3,3,3	0.49	0	2,2,2	0.14	0
4	EDO	C	208	-	3,3,3	0.49	0	2,2,2	0.24	0
4	EDO	C	207	-	3,3,3	0.42	0	2,2,2	0.34	0
4	EDO	F	155	-	3,3,3	0.39	0	2,2,2	0.41	0
4	EDO	D	127	-	3,3,3	0.42	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	209	-	-	0/1/1/1	-
4	EDO	C	206	-	-	0/1/1/1	-
4	EDO	F	9	-	-	1/1/1/1	-
4	EDO	C	210	-	-	0/1/1/1	-
4	EDO	C	211	-	-	0/1/1/1	-
4	EDO	C	212	-	-	1/1/1/1	-
4	EDO	D	126	-	-	0/1/1/1	-
5	PEG	C	214	-	-	1/4/4/4	-
4	EDO	F	5	-	-	1/1/1/1	-
4	EDO	C	213	-	-	0/1/1/1	-
4	EDO	C	208	-	-	1/1/1/1	-
4	EDO	C	207	-	-	0/1/1/1	-
4	EDO	F	155	-	-	0/1/1/1	-
4	EDO	D	127	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	208	EDO	O1-C1-C2-O2
5	C	214	PEG	C4-C3-O2-C2
4	C	212	EDO	O1-C1-C2-O2
4	F	5	EDO	O1-C1-C2-O2
4	F	9	EDO	O1-C1-C2-O2

There are no ring outliers.

10 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	209	EDO	1	0
4	C	210	EDO	1	0
4	C	212	EDO	1	0
5	C	214	PEG	3	0
4	F	5	EDO	1	0
4	C	213	EDO	1	0
4	C	208	EDO	1	0
4	C	207	EDO	1	0
4	F	155	EDO	2	0
4	D	127	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	C	200/205 (97%)	-0.35	7 (3%)	47 53	6, 12, 38, 78	16 (8%)
2	D	121/125 (96%)	0.33	10 (8%)	17 20	9, 21, 43, 74	6 (4%)
3	F	127/142 (89%)	0.38	17 (13%)	7 8	9, 19, 59, 75	7 (5%)
All	All	448/472 (94%)	0.04	34 (7%)	20 23	6, 16, 48, 78	29 (6%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	40	ILE	7.9
3	F	136	VAL	5.1
1	C	94	ASP	4.4
3	F	25	PHE	3.7
2	D	122	ASN	3.7
3	F	84	THR	3.7
3	F	86	SER	3.5
1	C	180	ASP	3.4
3	F	139	THR	3.3
3	F	140	ALA	3.3
3	F	137	PRO	3.2
2	D	16	VAL	3.1
3	F	42	ALA	2.9
1	C	101	ASN	2.9
1	C	100	GLU	2.7
2	D	98	GLN	2.7
2	D	13	PRO	2.7
2	D	2	LEU	2.6
2	D	3	TYR	2.6
3	F	142	HIS	2.6
3	F	141	GLY	2.6
2	D	14	GLN	2.5
3	F	135	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	55	ASP	2.5
3	F	87	ALA	2.4
1	C	179	SER	2.4
1	C	16	LYS	2.3
1	C	176	LYS	2.3
3	F	41	GLY	2.3
3	F	89	SER	2.2
2	D	24	ASN	2.2
3	F	83	ASN	2.1
3	F	74	HIS	2.1
2	D	10	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	C	209	4/4	0.68	0.29	33,49,82,86	0
4	EDO	D	127	4/4	0.75	0.22	38,44,51,72	0
5	PEG	C	214	7/7	0.84	0.18	38,55,65,78	0
4	EDO	C	213	4/4	0.86	0.25	15,37,47,62	0
4	EDO	C	207	4/4	0.87	0.19	30,35,45,61	0
4	EDO	C	208	4/4	0.88	0.20	18,24,28,34	0
4	EDO	F	9	4/4	0.89	0.16	25,30,46,51	0
4	EDO	F	5	4/4	0.90	0.14	34,36,45,84	0
4	EDO	C	210	4/4	0.91	0.17	11,28,31,41	0
4	EDO	C	212	4/4	0.93	0.15	17,25,49,85	0
4	EDO	C	206	4/4	0.93	0.15	27,34,62,77	0
4	EDO	F	155	4/4	0.94	0.14	19,22,24,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	D	126	4/4	0.94	0.11	20,31,39,40	0
4	EDO	C	211	4/4	0.99	0.04	8,9,10,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.