



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:39 AM UTC

PDB ID : 5BWD / pdb_00005bwd
Title : Benzylsuccinate alpha-gamma bound to fumarate
Authors : Funk, M.A.; Drennan, C.L.
Deposited on : 2015-06-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	878	13% 88% 9% .
2	C	60	5% 60% 5% 35%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called benzy succinate synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	853	Total	C	N	O	S	0	1	0
			6756	4265	1173	1282	36			


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	789	ILE	MET	engineered mutation	UNP O68395
A	866	SER	-	expression tag	UNP O68395
A	867	GLY	-	expression tag	UNP O68395
A	868	THR	-	expression tag	UNP O68395
A	869	GLY	-	expression tag	UNP O68395
A	870	SER	-	expression tag	UNP O68395
A	871	GLY	-	expression tag	UNP O68395
A	872	SER	-	expression tag	UNP O68395
A	873	SER	-	expression tag	UNP O68395
A	874	HIS	-	expression tag	UNP O68395
A	875	HIS	-	expression tag	UNP O68395
A	876	HIS	-	expression tag	UNP O68395
A	877	HIS	-	expression tag	UNP O68395
A	878	HIS	-	expression tag	UNP O68395
A	879	HIS	-	expression tag	UNP O68395

- Molecule 2 is a protein called benzy succinate synthase gamma chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	39	Total	C	N	O	S	0	0	0
			310	198	50	61	1			

- Molecule 3 is FUMARIC ACID (CCD ID: FUM) (formula: C₄H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	4	4		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	394	Total	O	0	0
			394	394		
7	C	20	Total	O	0	0
			20	20		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: benzy succinate synthase alpha chain



- Molecule 2: benzy succinate synthase gamma chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	154.86Å 154.86Å 82.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.63 – 2.00 34.63 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.9 (34.63-2.00) 83.8 (34.63-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 2.00Å)	Xtriage
Refinement program	PHENIX dev_1951	Depositor
R, R_{free}	0.195 , 0.223 0.198 , 0.226	Depositor DCC
R_{free} test set	3273 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.837	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7559	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, FUM, PG4, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/6911	0.68	2/9339 (0.0%)
2	C	0.26	0/318	0.66	0/428
All	All	0.29	0/7229	0.67	2/9767 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	747	GLY	CA-C-N	5.10	124.41	118.85
1	A	747	GLY	C-N-CA	5.10	124.41	118.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6756	0	6590	45	0
2	C	310	0	287	2	0
3	A	8	0	1	0	0
4	A	8	0	12	0	0
5	A	50	0	70	9	0
6	A	13	0	18	0	0
7	A	394	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	20	0	0	0	0
All	All	7559	0	6978	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:606:LEU:HD12	5:A:905:PGE:H42	1.62	0.81
1:A:666:PRO:HB2	1:A:675:ALA:HB2	1.66	0.77
1:A:251:LYS:NZ	7:A:1001:HOH:O	2.19	0.74
1:A:412:ASP:OD2	1:A:439:LYS:NZ	2.30	0.64
1:A:235:TRP:CE3	5:A:904:PGE:H6	2.33	0.63
1:A:652:TRP:HD1	1:A:655:PHE:HB2	1.64	0.62
1:A:737:ILE:HD13	1:A:755:VAL:HG13	1.84	0.58
1:A:313:LYS:NZ	1:A:366:GLU:OE2	2.36	0.56
1:A:443:TRP:HE1	5:A:908:PGE:H1	1.71	0.55
1:A:270:LEU:HD13	2:C:43:SER:HB2	1.89	0.54
1:A:683:TYR:O	1:A:762:GLN:NE2	2.41	0.52
1:A:494:MET:HG3	1:A:707:GLN:HB3	1.92	0.52
1:A:236:ASN:H	5:A:904:PGE:H5	1.73	0.52
1:A:31:PRO:HB2	1:A:34:GLU:HG2	1.91	0.52
1:A:502:ARG:HG2	1:A:503:LYS:HG3	1.92	0.52
1:A:439:LYS:HE2	5:A:908:PGE:H22	1.92	0.52
1:A:640:THR:HG23	1:A:643:GLN:H	1.76	0.51
1:A:476:ASN:HB3	5:A:905:PGE:H2	1.92	0.50
1:A:772:SER:HA	1:A:850:GLU:OE2	2.13	0.48
1:A:604:CYS:SG	7:A:1237:HOH:O	2.59	0.48
1:A:475:LEU:HD12	5:A:905:PGE:H52	1.96	0.47
1:A:614:HIS:O	1:A:616:PRO:HD3	2.15	0.47
1:A:253:MET:HE2	1:A:590:SER:HB3	1.96	0.46
2:C:29:CYS:HB2	2:C:46:ILE:HD13	1.98	0.46
1:A:623:ALA:HB1	1:A:679:ILE:HG12	1.97	0.46
1:A:419:LYS:HD2	1:A:446:GLU:HB3	1.97	0.46
1:A:552:GLU:O	1:A:556:GLU:N	2.49	0.46
1:A:852:ASP:OD1	1:A:853:PHE:N	2.49	0.46
1:A:655:PHE:HB3	1:A:658:MET:HB3	1.97	0.46
1:A:439:LYS:HG3	5:A:908:PGE:H32	1.98	0.45
1:A:456:SER:HB2	1:A:799:HIS:HE2	1.82	0.45
5:A:905:PGE:H32	5:A:905:PGE:H5	1.79	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:ALA:HB1	1:A:483:GLU:HB3	2.00	0.44
1:A:681:ARG:NE	1:A:685:GLU:OE2	2.31	0.44
1:A:569:CYS:HB3	1:A:614:HIS:CE1	2.52	0.44
1:A:517:PRO:HD2	1:A:617:ILE:O	2.17	0.43
1:A:647:ALA:HB1	1:A:652:TRP:CD1	2.53	0.43
1:A:174:PHE:CZ	1:A:572:THR:HA	2.56	0.41
1:A:629:ILE:HD11	1:A:723:PRO:HG2	2.01	0.41
1:A:275:ALA:HB2	1:A:288:LEU:HB2	2.02	0.41
1:A:363:VAL:HG11	1:A:414:ILE:HA	2.02	0.41
1:A:185:VAL:HB	1:A:718:ARG:NH2	2.35	0.41
1:A:722:THR:OG1	1:A:724:ASP:OD1	2.27	0.41
1:A:185:VAL:HB	1:A:718:ARG:HH22	1.85	0.41
1:A:459:HIS:HE1	1:A:461:GLU:HB2	1.86	0.40
1:A:518:ALA:HA	1:A:622:ALA:HB2	2.02	0.40
1:A:375:ALA:O	1:A:382:ARG:NH1	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	850/878 (97%)	825 (97%)	23 (3%)	2 (0%)	43	42
2	C	37/60 (62%)	36 (97%)	1 (3%)	0	100	100
All	All	887/938 (95%)	861 (97%)	24 (3%)	2 (0%)	43	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	827	VAL
1	A	828	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	714/741 (96%)	711 (100%)	3 (0%)	84	89
2	C	31/51 (61%)	31 (100%)	0	100	100
All	All	745/792 (94%)	742 (100%)	3 (0%)	84	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	VAL
1	A	827	VAL
1	A	863	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	499	HIS
1	A	769	GLN
1	A	786	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FUM	A	901	-	7,7,7	1.14	0	8,8,8	0.88	0
5	PGE	A	904	-	9,9,9	0.40	0	8,8,8	0.54	0
5	PGE	A	906	-	9,9,9	0.46	0	8,8,8	0.25	0
6	PG4	A	907	-	12,12,12	0.45	0	11,11,11	0.32	0
5	PGE	A	903	-	9,9,9	0.45	0	8,8,8	0.29	0
5	PGE	A	908	-	9,9,9	0.45	0	8,8,8	0.31	0
5	PGE	A	905	-	9,9,9	0.44	0	8,8,8	0.39	0
4	TRS	A	902	-	7,7,7	0.30	0	9,9,9	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FUM	A	901	-	-	0/5/5/5	-
5	PGE	A	904	-	-	1/7/7/7	-
5	PGE	A	906	-	-	0/7/7/7	-
6	PG4	A	907	-	-	0/10/10/10	-
5	PGE	A	903	-	-	0/7/7/7	-
5	PGE	A	908	-	-	1/7/7/7	-
5	PGE	A	905	-	-	3/7/7/7	-
4	TRS	A	902	-	-	4/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	902	TRS	N-C-C2-O2
5	A	905	PGE	O2-C3-C4-O3
5	A	908	PGE	O3-C5-C6-O4
5	A	904	PGE	C3-C4-O3-C5
4	A	902	TRS	C1-C-C2-O2
5	A	905	PGE	C3-C4-O3-C5
4	A	902	TRS	C2-C-C1-O1
5	A	905	PGE	O1-C1-C2-O2
4	A	902	TRS	N-C-C1-O1

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	904	PGE	2	0
5	A	908	PGE	3	0
5	A	905	PGE	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	853/878 (97%)	0.50	112 (13%) 7 6	18, 42, 110, 177	1 (0%)
2	C	39/60 (65%)	0.52	3 (7%) 19 18	37, 51, 81, 84	0
All	All	892/938 (95%)	0.50	115 (12%) 7 6	18, 43, 110, 177	1 (0%)

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	554	VAL	6.0
1	A	708	ALA	5.9
1	A	627	VAL	5.8
1	A	545	LEU	4.8
1	A	548	LEU	4.5
1	A	752	LEU	4.5
1	A	652	TRP	4.4
1	A	628	ALA	4.3
1	A	766	LEU	4.1
1	A	626	LEU	3.9
1	A	6	VAL	3.8
1	A	668	TRP	3.8
1	A	551	PHE	3.7
1	A	680	THR	3.7
1	A	7	SER	3.7
1	A	634	PHE	3.7
1	A	840	TYR	3.7
2	C	9	CYS	3.6
1	A	679	ILE	3.6
1	A	633	VAL	3.4
1	A	620	ILE	3.4
1	A	758	VAL	3.4
1	A	776	MET	3.4
1	A	641	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	385	PHE	3.4
1	A	755	VAL	3.3
1	A	666	PRO	3.2
1	A	654	GLY	3.2
1	A	740	TYR	3.1
1	A	772	SER	3.1
1	A	775	ILE	3.1
1	A	732	ALA	3.1
1	A	624	ASN	3.1
1	A	621	VAL	3.1
1	A	728	GLY	3.0
1	A	660	VAL	3.0
1	A	716	GLY	3.0
1	A	683	TYR	3.0
1	A	767	LEU	3.0
1	A	771	LEU	3.0
1	A	618	THR	3.0
1	A	682	PHE	3.0
1	A	675	ALA	3.0
1	A	733	ASP	3.0
1	A	739	PRO	3.0
1	A	662	PHE	3.0
1	A	632	LEU	2.9
1	A	10	VAL	2.9
1	A	845	ILE	2.9
1	A	655	PHE	2.9
1	A	8	ALA	2.9
1	A	622	ALA	2.9
1	A	65	ILE	2.9
1	A	667	LYS	2.9
1	A	727	PHE	2.9
1	A	674	TYR	2.8
1	A	665	ALA	2.8
1	A	555	TRP	2.8
1	A	751	VAL	2.8
1	A	853	PHE	2.8
1	A	865	ILE	2.8
1	A	860	PHE	2.8
1	A	731	ALA	2.7
1	A	830	TYR	2.7
1	A	827	VAL	2.7
1	A	678	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	686	ILE	2.7
1	A	25	PRO	2.7
1	A	792	TRP	2.6
1	A	384	ILE	2.6
1	A	639	TYR	2.6
1	A	643	GLN	2.6
1	A	151	ALA	2.6
1	A	863	VAL	2.5
1	A	669	GLY	2.5
1	A	749	THR	2.5
1	A	629	ILE	2.5
1	A	742	GLY	2.5
1	A	549	LYS	2.5
1	A	745	LYS	2.5
1	A	743	THR	2.5
1	A	741	MET	2.5
1	A	737	ILE	2.5
1	A	764	GLY	2.4
1	A	544	ASP	2.4
1	A	631	LYS	2.4
1	A	26	ALA	2.3
1	A	15	GLY	2.3
1	A	664	ARG	2.3
1	A	657	GLU	2.3
1	A	13	TYR	2.3
1	A	736	GLY	2.3
2	C	10	ALA	2.3
1	A	861	LEU	2.3
1	A	542	THR	2.2
1	A	858	LEU	2.2
1	A	844	THR	2.2
1	A	855	ALA	2.2
1	A	788	TYR	2.2
1	A	640	THR	2.2
1	A	849	GLN	2.2
1	A	753	ARG	2.2
1	A	543	GLY	2.1
1	A	623	ALA	2.1
1	A	754	SER	2.1
1	A	744	ASP	2.1
1	A	757	LYS	2.1
1	A	525	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	19	ASN	2.1
1	A	724	ASP	2.1
1	A	797	ILE	2.1
2	C	12	PHE	2.0
1	A	550	SER	2.0
1	A	734	ASP	2.0
1	A	29	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PGE	A	904	10/10	0.76	0.18	56,74,81,81	0
4	TRS	A	902	8/8	0.79	0.14	57,59,62,65	0
5	PGE	A	908	10/10	0.80	0.20	63,67,71,72	0
5	PGE	A	905	10/10	0.84	0.14	47,52,63,65	0
5	PGE	A	906	10/10	0.85	0.11	57,61,63,64	0
5	PGE	A	903	10/10	0.85	0.12	50,56,60,61	0
6	PG4	A	907	13/13	0.85	0.12	46,53,57,58	0
3	FUM	A	901	8/8	0.90	0.09	33,41,44,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.