



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:24 AM UTC

PDB ID : 4BX9 / pdb_00004bx9
Title : Human Vps33A in complex with a fragment of human Vps16
Authors : Graham, S.C.; Wartosch, L.; Gray, S.R.; Scourfield, E.J.; Deane, J.E.; Luzio, J.P.; Owen, D.J.
Deposited on : 2013-07-09
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	606	2% 87% 6% • 5%
1	B	606	4% 87% 8% • •
2	C	99	% 89% 7% •

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10193 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VACUOLAR PROTEIN SORTING-ASSOCIATED PROTEIN 33A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	574	Total	C	N	O	S	0	0	0
			4570	2918	780	853	19			
1	B	581	Total	C	N	O	S	0	0	0
			4637	2961	795	862	19			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	HIS	-	expression tag	UNP Q96AX1
A	598	HIS	-	expression tag	UNP Q96AX1
A	599	HIS	-	expression tag	UNP Q96AX1
A	600	HIS	-	expression tag	UNP Q96AX1
A	601	HIS	-	expression tag	UNP Q96AX1
A	602	HIS	-	expression tag	UNP Q96AX1
A	603	HIS	-	expression tag	UNP Q96AX1
A	604	HIS	-	expression tag	UNP Q96AX1
A	605	HIS	-	expression tag	UNP Q96AX1
A	606	HIS	-	expression tag	UNP Q96AX1
B	597	HIS	-	expression tag	UNP Q96AX1
B	598	HIS	-	expression tag	UNP Q96AX1
B	599	HIS	-	expression tag	UNP Q96AX1
B	600	HIS	-	expression tag	UNP Q96AX1
B	601	HIS	-	expression tag	UNP Q96AX1
B	602	HIS	-	expression tag	UNP Q96AX1
B	603	HIS	-	expression tag	UNP Q96AX1
B	604	HIS	-	expression tag	UNP Q96AX1
B	605	HIS	-	expression tag	UNP Q96AX1
B	606	HIS	-	expression tag	UNP Q96AX1


- Molecule 2 is a protein called VACUOLAR PROTEIN SORTING-ASSOCIATED PROTEIN 16 HOMOLOG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	95	Total	C	N	O	S	0	0	0
			766	483	144	138	1			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	638	GLY	-	expression tag	UNP Q9H269
C	639	PRO	-	expression tag	UNP Q9H269
C	640	HIS	-	expression tag	UNP Q9H269
C	641	MET	-	expression tag	UNP Q9H269

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		

- Molecule 4 is (2S)-2-hydroxybutanedioic acid (CCD ID: LMR) (formula: $\text{C}_4\text{H}_6\text{O}_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			9	4	5		

- Molecule 5 is MALONIC ACID (CCD ID: MLA) (formula: $C_3H_4O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			7	3	4		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	101	Total 101	O 101	0	0
6	B	84	Total 84	O 84	0	0
6	C	10	Total 10	O 10	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	128.41 Å 128.41 Å 263.36 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.84 – 2.60 85.84 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (85.84-2.60) 100.0 (85.84-2.60)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.92 (at 2.62 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.176 , 0.210 0.180 , 0.210	Depositor DCC
R_{free} test set	3465 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	62.9	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10193	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: LMR, MLA, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	3/4657 (0.1%)	1.04	3/6297 (0.0%)
1	B	1.11	6/4724 (0.1%)	1.06	6/6381 (0.1%)
2	C	1.20	0/777	1.14	0/1045
All	All	1.12	9/10158 (0.1%)	1.06	9/13723 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	143	ASP	C-O	-7.48	1.17	1.23
1	B	144	LEU	C-O	-6.94	1.15	1.24
1	B	211	ARG	C-O	-6.52	1.16	1.24
1	B	288	GLN	C-O	-5.43	1.17	1.24
1	A	134	PHE	C-O	-5.31	1.17	1.24
1	B	138	GLU	CA-C	-5.28	1.46	1.52
1	A	6	SER	C-O	-5.20	1.16	1.24
1	A	292	ALA	C-O	-5.15	1.17	1.24
1	B	590	ALA	C-O	-5.12	1.17	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	402	SER	CB-CA-C	-5.76	101.20	109.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	VAL	CB-CA-C	-5.51	104.87	110.93
1	B	207	ILE	N-CA-C	-5.45	105.40	110.53
1	B	534	GLN	N-CA-C	-5.33	106.61	113.01
1	B	307	VAL	N-CA-C	5.17	115.39	110.53
1	B	335	LYS	CA-C-N	5.10	127.42	120.54
1	B	335	LYS	C-N-CA	5.10	127.42	120.54
1	A	69	ASN	CB-CA-C	-5.07	109.20	115.89
1	A	336	GLN	N-CA-C	5.06	116.60	111.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	535	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4570	0	4588	20	0
1	B	4637	0	4670	22	0
2	C	766	0	778	4	0
3	A	3	0	1	0	0
3	B	6	0	3	0	0
4	A	9	0	4	1	0
5	C	7	0	2	0	0
6	A	101	0	0	0	0
6	B	84	0	0	0	0
6	C	10	0	0	0	0
All	All	10193	0	10046	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:442:LEU:CD2	1:A:593:GLU:HG3	2.15	0.76
1:B:324:GLU:HG2	1:B:337:PHE:CZ	2.21	0.75
1:A:442:LEU:HD21	1:A:593:GLU:HG3	1.71	0.73
1:A:582:MET:HE2	1:A:587:TRP:CD1	2.25	0.71
2:C:671:GLU:HA	2:C:674:MET:HE3	1.76	0.68
1:A:341:LEU:HA	1:A:344:MET:HE3	1.80	0.63
1:B:582:MET:HE2	1:B:587:TRP:CD1	2.37	0.59
1:B:341:LEU:HA	1:B:344:MET:HE3	1.85	0.57
1:A:442:LEU:HD21	1:A:593:GLU:CG	2.33	0.57
1:A:330:THR:OG1	1:A:333:GLU:CD	2.48	0.56
2:C:729:LEU:HD23	2:C:729:LEU:C	2.31	0.56
2:C:692:LEU:O	2:C:693:ASP:HB2	2.05	0.55
1:B:331:VAL:O	1:B:335:LYS:HG3	2.07	0.54
1:B:54:LEU:C	1:B:54:LEU:HD23	2.35	0.52
1:A:137:ARG:C	1:A:138:GLU:HG2	2.34	0.51
1:A:404:ILE:HD11	1:A:589:GLU:HG2	1.93	0.51
1:B:190:PRO:HG3	1:B:529:LEU:HD21	1.91	0.51
1:A:582:MET:CE	1:A:587:TRP:CD1	2.95	0.49
1:A:54:LEU:C	1:A:54:LEU:HD23	2.38	0.49
1:A:404:ILE:CD1	1:A:589:GLU:HG2	2.44	0.48
1:A:166:GLU:HA	1:B:528:PRO:HG3	1.95	0.48
1:B:533:LEU:HD12	1:B:533:LEU:O	2.16	0.46
1:B:108:ASP:HB3	1:B:135:ILE:HD12	1.97	0.46
1:A:269:ALA:HA	1:A:270:PRO:HA	1.85	0.45
1:A:108:ASP:HB3	1:A:135:ILE:HD12	1.98	0.45
1:B:394:GLU:OE2	1:B:427:TYR:OH	2.29	0.45
1:B:324:GLU:HG2	1:B:337:PHE:CE2	2.50	0.45
1:B:116:ARG:HA	1:B:139:GLU:HB2	1.98	0.44
1:B:41:TYR:CE1	1:B:84:ARG:HD3	2.51	0.44
1:B:534:GLN:HE21	1:B:534:GLN:CA	2.31	0.43
1:B:534:GLN:HE21	1:B:534:GLN:HA	1.84	0.43
1:A:70:ARG:HG2	1:A:71:LEU:O	2.18	0.43
1:A:509:ARG:NE	1:A:509:ARG:HA	2.33	0.43
1:B:70:ARG:HG2	1:B:71:LEU:O	2.19	0.43
1:B:509:ARG:NE	1:B:509:ARG:HA	2.35	0.42
1:B:195:LYS:NZ	1:B:520:GLY:O	2.50	0.42
1:A:173:TYR:OH	1:A:177:LYS:HE2	2.20	0.42
1:B:400:LYS:O	1:B:400:LYS:CG	2.68	0.41
2:C:705:LEU:HD23	2:C:705:LEU:HA	1.90	0.41
1:A:425:LEU:HG	1:A:429:LYS:HE2	2.03	0.41
1:B:427:TYR:CD2	1:B:427:TYR:C	2.98	0.41
1:B:341:LEU:N	1:B:342:PRO:CD	2.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:341:LEU:HA	1:A:344:MET:CE	2.49	0.40
4:A:1597:LMR:O1A	1:B:208:ARG:NH1	2.52	0.40
1:A:341:LEU:N	1:A:342:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	568/606 (94%)	554 (98%)	14 (2%)	0	100	100
1	B	575/606 (95%)	561 (98%)	14 (2%)	0	100	100
2	C	93/99 (94%)	91 (98%)	2 (2%)	0	100	100
All	All	1236/1311 (94%)	1206 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/528 (94%)	478 (96%)	18 (4%)	31	58
1	B	503/528 (95%)	486 (97%)	17 (3%)	32	60
2	C	76/80 (95%)	75 (99%)	1 (1%)	61	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1075/1136 (95%)	1039 (97%)	36 (3%)	33 61

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	PHE
1	A	67	LYS
1	A	106	THR
1	A	177	LYS
1	A	191	GLN
1	A	216	SER
1	A	330	THR
1	A	333	GLU
1	A	336	GLN
1	A	374	LEU
1	A	421	LYS
1	A	477	VAL
1	A	486	SER
1	A	513	GLU
1	A	524	GLU
1	A	531	THR
1	A	582	MET
1	A	593	GLU
1	B	46	PHE
1	B	67	LYS
1	B	88	GLU
1	B	106	THR
1	B	191	GLN
1	B	231	ASP
1	B	324	GLU
1	B	330	THR
1	B	374	LEU
1	B	421	LYS
1	B	486	SER
1	B	524	GLU
1	B	533	LEU
1	B	534	GLN
1	B	536	LYS
1	B	582	MET
1	B	593	GLU
2	C	647	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	ASN
1	B	96	ASN
1	B	201	GLN
1	B	399	GLN
1	B	527	GLN
1	B	534	GLN
2	C	709	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	LMR	A	1597	-	8,8,8	1.50	2 (25%)	10,10,10	1.31	1 (10%)
3	FMT	B	1598	-	2,2,2	0.44	0	1,1,1	1.23	0
3	FMT	A	1596	-	2,2,2	0.72	0	1,1,1	0.63	0
3	FMT	B	1599	-	2,2,2	1.42	0	1,1,1	0.29	0
5	MLA	C	1737	-	6,6,6	1.30	0	7,7,7	1.22	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LMR	A	1597	-	-	5/8/8/8	-
5	MLA	C	1737	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1597	LMR	O4B-C4	2.27	1.29	1.22
4	A	1597	LMR	O1A-C1	2.19	1.28	1.22

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1597	LMR	O1B-C1-C2	2.60	118.23	112.74
5	C	1737	MLA	O1A-C1-C2	2.08	120.95	114.51

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1597	LMR	O1B-C1-C2-C3
4	A	1597	LMR	O1B-C1-C2-O2
4	A	1597	LMR	O1A-C1-C2-C3
4	A	1597	LMR	O1A-C1-C2-O2
4	A	1597	LMR	O2-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1597	LMR	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/606 (94%)	-0.49	10 (1%) 69 64	40, 59, 109, 160	0
1	B	581/606 (95%)	-0.17	22 (3%) 44 38	42, 70, 123, 184	0
2	C	95/99 (95%)	-0.37	1 (1%) 78 74	53, 68, 101, 133	0
All	All	1250/1311 (95%)	-0.33	33 (2%) 57 51	40, 64, 117, 184	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	597	HIS	6.5
1	A	595	PRO	4.7
1	B	269	ALA	4.7
2	C	642	GLU	3.9
1	B	338	VAL	3.8
1	A	531	THR	3.6
1	B	570	GLY	3.6
1	B	595	PRO	3.6
1	B	337	PHE	3.6
1	B	438	TYR	3.5
1	A	331	VAL	3.1
1	B	594	LYS	2.8
1	B	87	LEU	2.8
1	B	329	LYS	2.8
1	B	125	LEU	2.8
1	A	270	PRO	2.7
1	B	331	VAL	2.7
1	B	332	GLY	2.5
1	B	330	THR	2.4
1	A	460	ARG	2.4
1	A	570	GLY	2.4
1	B	88	GLU	2.4
1	B	460	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	330	THR	2.3
1	A	594	LYS	2.3
1	B	326	HIS	2.2
1	B	474	MET	2.2
1	A	474	MET	2.2
1	A	269	ALA	2.1
1	B	536	LYS	2.1
1	B	538	GLN	2.1
1	B	105	PRO	2.0
1	B	126	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	LMR	A	1597	9/9	0.87	0.12	63,68,83,92	0
3	FMT	B	1599	3/3	0.91	0.22	74,74,75,92	0
3	FMT	B	1598	3/3	0.91	0.18	80,80,84,86	0
5	MLA	C	1737	7/7	0.92	0.09	68,78,85,93	0
3	FMT	A	1596	3/3	0.94	0.22	78,78,87,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.