



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:50 PM UTC

PDB ID : 4BXD / pdb_00004bxid
Title : CRYSTAL STRUCTURE OF AMPDH3 FROM PSEUDOMONAS AERUGINOSA IN COMPLEX WITH TETRASACCHARIDE PENTAPEPTIDE
Authors : Artola-Recolons, C.; Hermoso, J.A.
Deposited on : 2013-07-10
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	
1	B	255	
2	C	4	
2	D	4	
3	E	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	6CL	D	3	-	-	X	-
2	DAL	D	4	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8343 atoms, of which 4079 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMPDH3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	255	4005	1300	1969	354	379	3	0	0	0
1	B	255	4004	1300	1968	354	379	3	0	0	0

- Molecule 2 is a protein called PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	4	60	18	28	5	9	0	0	0
2	D	4	60	18	28	5	9	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-methyl 2-acetamido-2-deoxy-beta-D-glucopyranoside.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	E	3	87	25	43	3	16	0	0	0
3	F	3	87	25	43	3	16	0	0	0

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	22	Total	O	0	0
			22	22		
5	B	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AMPDH3



- Molecule 1: AMPDH3



- Molecule 2: PEPTIDE



- Molecule 2: PEPTIDE



A1
E2
K3
A4

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-methyl 2-acetamido-2-deoxy-beta-D-glucopyranoside

Chain E:  100%

MAG1
MAG2
MAG3

- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-methyl 2-acetamido-2-deoxy-beta-D-glucopyranoside

Chain F:  100%

MAG1
MAG2
MAG3

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.19Å 101.19Å 162.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.30 – 3.10 48.30 – 3.10	Depositor EDS
% Data completeness (in resolution range)	95.0 (48.30-3.10) 89.8 (48.30-3.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.95 (at 3.12Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.224 , 0.277 0.231 , 0.281	Depositor DCC
R_{free} test set	795 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtrriage
Anisotropy	0.538	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8343	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAG, ZN, FGA, DAL, 6CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.84	39/2091 (1.9%)	1.22	13/2842 (0.5%)
1	B	1.68	28/2092 (1.3%)	1.24	15/2845 (0.5%)
2	C	1.85	0/4	2.85	0/4
2	D	1.50	0/4	1.25	0/4
All	All	1.76	67/4191 (1.6%)	1.23	28/5695 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	218	PHE	C-N	10.81	1.48	1.33
1	A	46	HIS	CG-ND1	-8.07	1.29	1.38
1	A	29	LEU	C-N	7.96	1.43	1.33
1	A	79	HIS	CG-ND1	-7.81	1.29	1.38
1	A	146	HIS	CG-ND1	-7.76	1.29	1.38

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ASP	N-CA-C	9.43	124.16	112.47
1	A	129	LYS	N-CA-C	-7.68	102.85	111.14
1	B	213	VAL	N-CA-C	7.51	117.51	106.85
1	B	135	TYR	CA-C-N	-7.09	112.58	120.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	135	TYR	C-N-CA	-7.09	112.58	120.45

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	93	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2036	1969	1967	96	0
1	B	2036	1968	1968	118	0
2	C	32	28	29	5	0
2	D	32	28	29	10	0
3	E	44	43	41	5	0
3	F	44	43	41	10	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	22	0	0	1	0
5	B	16	0	0	0	0
All	All	4264	4079	4075	219	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 219 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:192:LEU:HD12	1:B:193:PRO:CD	1.83	1.07
1:B:192:LEU:HD12	1:B:193:PRO:HD3	1.41	1.03
1:B:154:LYS:NZ	3:F:1:MAG:O3	1.94	1.01
1:A:107:ASP:CA	1:A:108:ASP:N	2.28	0.97
1:A:93:ASP:OD2	1:B:88:ARG:NH2	2.03	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/255 (98%)	220 (88%)	31 (12%)	0	100	100
1	B	253/255 (99%)	218 (86%)	34 (13%)	1 (0%)	30	61
All	All	504/510 (99%)	438 (87%)	65 (13%)	1 (0%)	43	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	236	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/205 (100%)	195 (95%)	10 (5%)	22	53
1	B	205/205 (100%)	192 (94%)	13 (6%)	16	45
All	All	410/410 (100%)	387 (94%)	23 (6%)	19	49

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	94	THR
1	B	156	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	155	SER
1	B	189	ARG
1	A	185	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	HIS
1	A	92	ASN
1	A	122	ASN
1	B	228	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	6CL	C	3	2	9,11,12	1.48	2 (22%)	5,13,15	3.38	2 (40%)
2	6CL	D	3	2	9,11,12	1.71	2 (22%)	5,13,15	3.92	1 (20%)
2	FGA	D	2	2	6,8,9	1.14	0	6,9,11	1.17	1 (16%)
2	FGA	C	2	2	6,8,9	1.11	0	6,9,11	1.37	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6CL	C	3	2	-	9/11/12/14	-
2	6CL	D	3	2	-	5/11/12/14	-
2	FGA	D	2	2	-	0/8/8/9	-
2	FGA	C	2	2	-	2/8/8/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	6CL	CB-CA	-3.10	1.48	1.53
2	D	3	6CL	CA-N	-2.96	1.39	1.48
2	C	3	6CL	CB-CA	-2.39	1.49	1.53
2	C	3	6CL	OH2-CZ	-2.23	1.23	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	6CL	OH2-CZ-OH1	8.71	143.84	124.08
2	C	3	6CL	OH2-CZ-OH1	6.99	139.93	124.08
2	C	2	FGA	OXT-C-O	-2.88	117.55	124.08
2	D	2	FGA	OXT-C-O	-2.39	118.65	124.08
2	C	3	6CL	CD-CE-CZ	2.26	116.43	110.45

There are no chirality outliers.

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	3	6CL	N-CA-CB-CG
2	C	3	6CL	C-CA-CB-CG
2	C	3	6CL	CG-CD-CE-CZ
2	C	3	6CL	NZ-CE-CZ-OH1
2	D	3	6CL	CG-CD-CE-CZ

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	6CL	3	0
2	D	3	6CL	7	0
2	D	2	FGA	2	0
2	C	2	FGA	1	0

5.5 Carbohydrates i

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAG	E	1	3	16,16,16	1.35	2 (12%)	22,22,22	2.29	8 (36%)
3	NAG	E	2	3	14,14,15	1.45	3 (21%)	17,19,21	2.37	7 (41%)
3	NAG	E	3	3	14,14,15	1.46	2 (14%)	17,19,21	2.67	9 (52%)
3	MAG	F	1	3	16,16,16	1.30	2 (12%)	22,22,22	2.42	8 (36%)
3	NAG	F	2	3	14,14,15	1.57	3 (21%)	17,19,21	2.69	10 (58%)
3	NAG	F	3	3	14,14,15	1.51	2 (14%)	17,19,21	3.29	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAG	E	1	3	-	5/8/28/28	0/1/1/1
3	NAG	E	2	3	-	2/6/23/26	0/1/1/1
3	NAG	E	3	3	-	6/6/23/26	0/1/1/1
3	MAG	F	1	3	-	4/8/28/28	0/1/1/1
3	NAG	F	2	3	-	2/6/23/26	0/1/1/1
3	NAG	F	3	3	-	3/6/23/26	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	3	NAG	O5-C1	-3.19	1.38	1.43
3	E	3	NAG	C7-N2	3.16	1.44	1.34
3	E	1	MAG	C1-C2	-3.04	1.48	1.53
3	F	2	NAG	C7-N2	2.92	1.43	1.34
3	F	3	NAG	C3-C2	-2.87	1.46	1.52

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	3	NAG	C1-O5-C5	9.36	124.73	112.19
3	E	3	NAG	C8-C7-N2	6.04	126.13	116.12
3	F	2	NAG	C1-O5-C5	5.66	119.77	112.19
3	F	1	MAG	C8-C7-N2	5.64	125.47	116.12
3	E	1	MAG	C1-O5-C5	5.49	124.45	113.72

There are no chirality outliers.

5 of 22 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	1	MAG	C1-C2-N2-C7
3	F	3	NAG	C1-C2-N2-C7
3	E	2	NAG	O5-C5-C6-O6
3	E	3	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6


There are no ring outliers.

6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	1	MAG	5	0
3	F	3	NAG	4	0
3	E	2	NAG	1	0
3	F	2	NAG	4	0
3	E	3	NAG	1	0
3	E	1	MAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	107:ASP	C	108:ASP	N	2.29

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/255 (100%)	-0.18	2 (0%) 82 65	19, 33, 53, 98	0
1	B	255/255 (100%)	-0.06	0 100 100	19, 38, 67, 95	0
2	C	1/4 (25%)	0.31	0 100 100	59, 59, 59, 59	0
2	D	1/4 (25%)	0.55	0 100 100	59, 59, 59, 59	0
All	All	512/518 (98%)	-0.12	2 (0%) 88 76	19, 35, 64, 98	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	255	ALA	2.4
1	A	108	ASP	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DAL	D	4	6/6	0.70	0.19	29,42,58,70	0
2	FGA	C	2	9/10	0.80	0.17	24,33,58,80	0
2	6CL	C	3	12/13	0.83	0.15	29,46,55,60	0
2	6CL	D	3	12/13	0.85	0.12	23,49,59,59	0
2	FGA	D	2	9/10	0.88	0.11	32,53,65,65	0
2	DAL	C	4	6/6	0.90	0.13	13,18,32,33	0


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	F	2	14/15	0.52	0.19	46,73,91,93	0
3	NAG	F	3	14/15	0.66	0.16	57,75,95,101	0
3	MAG	E	1	16/16	0.72	0.21	53,81,103,114	0
3	NAG	E	3	14/15	0.77	0.15	58,88,103,112	0
3	MAG	F	1	16/16	0.81	0.16	52,75,93,94	0
3	NAG	E	2	14/15	0.85	0.14	51,81,99,101	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q < 0.9
4	ZN	A	1260	1/1	0.95	0.07	35,35,35,35	0
4	ZN	B	1256	1/1	0.97	0.10	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.