



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:29 PM UTC

PDB ID : 7BY1 / pdb_00007by1
Title : Crystal structure of GCN5 PCAF N-terminal domain
Authors : Hibi, R.; Toma-Fukai, S.; Shimizu, T.
Deposited on : 2020-04-21
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	10% 85% 6% 9%
1	B	317	8% 82% 8% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5077 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase KAT2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	0	0	0
			2336	1495	410	414	17			
1	B	285	Total	C	N	O	S	0	0	0
			2322	1485	408	412	17			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	GLY	-	expression tag	UNP Q9JHD2
A	63	PRO	-	expression tag	UNP Q9JHD2
A	64	LEU	-	expression tag	UNP Q9JHD2
A	65	GLY	-	expression tag	UNP Q9JHD2
A	66	SER	-	expression tag	UNP Q9JHD2
B	62	GLY	-	expression tag	UNP Q9JHD2
B	63	PRO	-	expression tag	UNP Q9JHD2
B	64	LEU	-	expression tag	UNP Q9JHD2
B	65	GLY	-	expression tag	UNP Q9JHD2
B	66	SER	-	expression tag	UNP Q9JHD2

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total 211	O 211	0	0
3	B	204	Total 204	O 204	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Histone acetyltransferase KAT2A



• Molecule 1: Histone acetyltransferase KAT2A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.07Å 148.86Å 59.00Å 90.00° 101.03° 90.00°	Depositor
Resolution (Å)	45.70 – 1.80 45.70 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.70-1.80) 100.0 (45.70-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.196 , 0.231 0.205 , 0.238	Depositor DCC
R_{free} test set	3322 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5077	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.18	2/2394 (0.1%)	1.12	4/3234 (0.1%)
1	B	1.23	5/2378 (0.2%)	1.16	3/3210 (0.1%)
All	All	1.20	7/4772 (0.1%)	1.14	7/6444 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	356	SER	C-O	6.19	1.31	1.24
1	A	123	PRO	CA-C	5.83	1.57	1.52
1	B	289	ARG	N-CA	5.59	1.53	1.46
1	B	318	ARG	C-O	5.43	1.30	1.24
1	B	326	ARG	CZ-NH1	5.43	1.40	1.32
1	B	219	PRO	CA-C	5.29	1.54	1.51
1	B	296	VAL	CA-C	5.21	1.57	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	367	SER	N-CA-C	6.73	119.39	109.50
1	B	367	SER	CA-C-N	6.43	126.98	120.04
1	B	367	SER	C-N-CA	6.43	126.98	120.04
1	A	367	SER	N-CA-C	5.69	117.86	109.50
1	A	367	SER	CA-C-N	5.31	125.78	120.04
1	A	367	SER	C-N-CA	5.31	125.78	120.04
1	A	239	SER	CA-CB-OG	-5.25	100.60	111.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2336	0	2339	11	0
1	B	2322	0	2325	12	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	211	0	0	3	0
3	B	204	0	0	2	0
All	All	5077	0	4664	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:GLU:HG3	3:A:652:HOH:O	1.95	0.66
1:A:349:THR:HG23	1:A:350:HIS:ND1	2.15	0.62
1:B:349:THR:HG23	1:B:350:HIS:ND1	2.14	0.61
1:A:108:LYS:HE3	3:A:584:HOH:O	2.06	0.55
1:A:263:TRP:NE1	1:A:327:GLN:OE1	2.39	0.53
1:B:263:TRP:NE1	1:B:327:GLN:OE1	2.40	0.53
1:B:227:GLU:O	1:B:231:LEU:HD23	2.13	0.49
1:A:249:MET:HE1	1:A:350:HIS:HB3	1.95	0.48
1:B:213:GLU:OE2	1:B:215:SER:HB3	2.13	0.48
1:A:249:MET:HE2	1:A:249:MET:N	2.30	0.46
1:A:223:LYS:HA	1:A:224:PRO:C	2.41	0.45
1:A:108:LYS:CE	3:A:584:HOH:O	2.64	0.45
1:B:231:LEU:HD22	3:B:547:HOH:O	2.17	0.44
1:A:249:MET:CE	1:A:350:HIS:HD2	2.32	0.43
1:B:276:GLN:NE2	3:B:504:HOH:O	2.51	0.43
1:B:249:MET:N	1:B:249:MET:HE2	2.35	0.42
1:B:108:LYS:HE2	1:B:155:HIS:HB3	2.00	0.42
1:B:249:MET:CE	1:B:350:HIS:HD2	2.32	0.42
1:A:108:LYS:HE2	1:A:155:HIS:HB3	2.01	0.42
1:B:207:MET:HE2	1:B:207:MET:HB3	1.95	0.42
1:A:249:MET:HE3	1:A:350:HIS:HD2	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:249:MET:HE1	1:B:350:HIS:HB3	2.02	0.41
1:B:223:LYS:HA	1:B:224:PRO:C	2.45	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/317 (89%)	281 (99%)	2 (1%)	0	100	100
1	B	281/317 (89%)	280 (100%)	1 (0%)	0	100	100
All	All	564/634 (89%)	561 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/288 (91%)	258 (98%)	4 (2%)	57	49
1	B	260/288 (90%)	255 (98%)	5 (2%)	50	41
All	All	522/576 (91%)	513 (98%)	9 (2%)	53	45

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	277	SER
1	A	333	ARG
1	A	375	THR
1	B	84	GLN
1	B	231	LEU
1	B	277	SER
1	B	333	ARG
1	B	375	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	228	GLN
1	A	235	GLN
1	A	276	GLN
1	A	350	HIS
1	B	228	GLN
1	B	276	GLN
1	B	310	HIS
1	B	350	HIS
1	B	366	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	287/317 (90%)	0.47	31 (10%) 11 9	14, 27, 51, 76	0
1	B	285/317 (89%)	0.44	25 (8%) 15 13	13, 27, 50, 68	0
All	All	572/634 (90%)	0.45	56 (9%) 13 11	13, 27, 51, 76	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	124	PRO	6.0
1	B	83	SER	5.9
1	A	83	SER	5.4
1	B	134	PRO	5.0
1	A	378	PRO	4.7
1	B	334	VAL	4.7
1	A	134	PRO	4.4
1	A	124	PRO	4.3
1	B	263	TRP	4.0
1	A	339	LEU	3.7
1	A	349	THR	3.7
1	B	348	LEU	3.7
1	B	349	THR	3.7
1	A	332	PHE	3.6
1	A	348	LEU	3.6
1	B	182	HIS	3.5
1	B	207	MET	3.5
1	A	342	GLU	3.4
1	A	344	ARG	3.3
1	A	345	THR	3.2
1	B	84	GLN	3.1
1	A	346	LEU	2.9
1	B	375	THR	2.8
1	A	84	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	207	MET	2.8
1	A	347	ILE	2.7
1	B	333	ARG	2.7
1	A	182	HIS	2.7
1	A	114	LYS	2.6
1	A	263	TRP	2.6
1	A	111	GLU	2.6
1	A	340	VAL	2.6
1	B	366	ASN	2.5
1	B	277	SER	2.4
1	B	347	ILE	2.4
1	A	244	ARG	2.3
1	B	109	ALA	2.3
1	B	114	LYS	2.3
1	A	112	THR	2.3
1	A	137	ASN	2.3
1	A	85	ARG	2.2
1	A	136	ALA	2.2
1	B	332	PHE	2.2
1	B	179	MET	2.2
1	A	367	SER	2.2
1	A	341	PRO	2.1
1	A	350	HIS	2.1
1	B	185	GLU	2.1
1	B	376	MET	2.1
1	B	178	PHE	2.1
1	B	367	SER	2.1
1	A	135	ALA	2.1
1	B	250	PHE	2.0
1	B	137	ASN	2.0
1	A	178	PHE	2.0
1	A	187	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	401	1/1	0.99	0.04	28,28,28,28	0
2	ZN	A	402	1/1	0.99	0.02	23,23,23,23	0
2	ZN	B	401	1/1	0.99	0.05	26,26,26,26	0
2	ZN	B	402	1/1	0.99	0.05	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around ZN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**6.5 Other polymers** ⓘ

There are no such residues in this entry.