



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:54 AM UTC

PDB ID : 5BZB / pdb_00005bzb
Title : NavMs voltage-gated sodium channel pore and C-terminal domain
Authors : Naylor, C.E.; Bagneris, C.; Wallace, B.A.
Deposited on : 2015-06-11
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	10% 57% 5% 38%
1	B	149	11% 56% . 40%
1	C	149	10% 57% . 39%
1	D	149	12% 55% 6% 39%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3269 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ion transport protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	92	Total	C	N	O	S	0	0	0
			718	488	106	119	5			
1	B	90	Total	C	N	O	S	0	0	0
			711	483	104	119	5			
1	C	91	Total	C	N	O	S	0	0	0
			715	485	105	120	5			
1	D	91	Total	C	N	O	S	0	0	0
			712	482	105	120	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	126	GLY	-	expression tag	UNP A0L5S6
A	127	SER	-	expression tag	UNP A0L5S6
A	128	HIS	-	expression tag	UNP A0L5S6
A	129	MET	-	expression tag	UNP A0L5S6
B	126	GLY	-	expression tag	UNP A0L5S6
B	127	SER	-	expression tag	UNP A0L5S6
B	128	HIS	-	expression tag	UNP A0L5S6
B	129	MET	-	expression tag	UNP A0L5S6
C	126	GLY	-	expression tag	UNP A0L5S6
C	127	SER	-	expression tag	UNP A0L5S6
C	128	HIS	-	expression tag	UNP A0L5S6
C	129	MET	-	expression tag	UNP A0L5S6
D	126	GLY	-	expression tag	UNP A0L5S6
D	127	SER	-	expression tag	UNP A0L5S6
D	128	HIS	-	expression tag	UNP A0L5S6
D	129	MET	-	expression tag	UNP A0L5S6

- Molecule 2 is DODECAETHYLENE GLYCOL (CCD ID: 12P) (formula: C₂₄H₅₀O₁₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	10	6		
2	B	1	Total	C	O	0	0
			16	10	6		
2	C	1	Total	C	O	0	0
			16	10	6		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Na	0	0
			3	3		
3	C	2	Total	Na	0	0
			2	2		
3	D	1	Total	Na	0	0
			1	1		

- Molecule 4 is HEGA-10 (CCD ID: 2CV) (formula: C₁₈H₃₇NO₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total 26	C 18	N 1	O 7	0	0
4	B	1	Total 26	C 18	N 1	O 7	0	0
4	D	1	Total 26	C 18	N 1	O 7	0	0
4	D	1	Total 26	C 18	N 1	O 7	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	60	Total O 60 60	0	0
5	B	61	Total O 61 61	0	0
5	C	66	Total O 66 66	0	0
5	D	68	Total O 68 68	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Ion transport protein




• Molecule 1: Ion transport protein



• Molecule 1: Ion transport protein



• Molecule 1: Ion transport protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	80.34Å 332.76Å 80.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.81 – 2.70 57.81 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (57.81-2.70) 99.7 (57.81-2.70)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.92 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.178 , 0.214 0.186 , 0.218	Depositor DCC
R_{free} test set	1548 reflections (3.47%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	1.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 81.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	3269	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2CV, 12P, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	0/740	1.42	0/1015
1	B	0.89	0/733	1.39	0/1005
1	C	0.87	0/737	1.37	0/1010
1	D	0.88	0/734	1.38	0/1006
All	All	0.89	0/2944	1.39	0/4036

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	718	0	726	3	0
1	B	711	0	721	4	0
1	C	715	0	724	2	0
1	D	712	0	715	4	0
2	A	16	0	21	0	0
2	B	16	0	21	0	0
2	C	16	0	21	0	0
3	A	3	0	0	0	0
3	C	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	0	0	0	0
4	B	52	0	74	2	0
4	D	52	0	74	0	0
5	A	60	0	0	0	0
5	B	61	0	0	0	0
5	C	66	0	0	0	0
5	D	68	0	0	0	0
All	All	3269	0	3097	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:LYS:HG2	4:B:301:2CV:H41	1.78	0.66
1:B:189:MET:HB3	4:B:302:2CV:H431	1.90	0.54
1:B:200:ILE:HG22	1:B:204:MET:HE2	1.91	0.52
1:C:194:ASN:O	1:C:197:VAL:HG12	2.10	0.52
1:D:200:ILE:HG22	1:D:204:MET:HE2	1.97	0.46
1:B:157:PHE:CZ	1:B:188:VAL:HA	2.50	0.45
1:A:157:PHE:CZ	1:A:188:VAL:HA	2.53	0.44
1:A:142:PHE:HZ	1:A:172:PHE:HA	1.84	0.43
1:A:200:ILE:HG22	1:A:204:MET:HE2	2.00	0.42
1:D:189:MET:HA	1:D:192:HIS:O	2.20	0.41
1:D:157:PHE:CZ	1:D:188:VAL:HA	2.55	0.41
1:C:189:MET:HA	1:C:192:HIS:O	2.20	0.41
1:D:180:TRP:O	1:D:185:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/149 (60%)	86 (96%)	4 (4%)	0	100	100
1	B	88/149 (59%)	86 (98%)	2 (2%)	0	100	100
1	C	89/149 (60%)	85 (96%)	4 (4%)	0	100	100
1	D	89/149 (60%)	85 (96%)	4 (4%)	0	100	100
All	All	356/596 (60%)	342 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/129 (60%)	76 (99%)	1 (1%)	61	83
1	B	78/129 (60%)	78 (100%)	0	100	100
1	C	78/129 (60%)	76 (97%)	2 (3%)	40	70
1	D	77/129 (60%)	76 (99%)	1 (1%)	61	83
All	All	310/516 (60%)	306 (99%)	4 (1%)	61	83

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	ILE
1	C	142	PHE
1	C	211	LEU
1	D	211	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	212	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	12P	A	301	-	15,15,36	0.14	0	14,14,35	0.10	0
4	2CV	B	301	-	25,25,25	0.39	0	29,30,30	0.71	2 (6%)
2	12P	B	303	-	15,15,36	0.14	0	14,14,35	0.14	0
2	12P	C	301	-	15,15,36	0.12	0	14,14,35	0.13	0
4	2CV	D	301	-	25,25,25	0.23	0	29,30,30	0.84	1 (3%)
4	2CV	D	302	-	25,25,25	0.18	0	29,30,30	0.67	0
4	2CV	B	302	-	25,25,25	0.22	0	29,30,30	1.03	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	12P	A	301	-	-	8/13/13/34	-
4	2CV	B	301	-	-	7/34/34/34	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	12P	B	303	-	-	6/13/13/34	-
2	12P	C	301	-	-	7/13/13/34	-
4	2CV	D	301	-	-	13/34/34/34	-
4	2CV	D	302	-	-	11/34/34/34	-
4	2CV	B	302	-	-	9/34/34/34	-

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	301	2CV	C35-N33-C36	3.24	119.81	116.14
4	B	302	2CV	O47-C37-C40	3.18	116.70	109.25
4	B	302	2CV	C35-N33-C36	2.61	119.09	116.14
4	B	302	2CV	C37-C40-C41	2.52	116.35	112.48
4	B	301	2CV	O47-C37-C40	2.32	114.67	109.25
4	B	301	2CV	C37-C40-C41	2.12	115.73	112.48

There are no chirality outliers.

All (61) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	301	2CV	N33-C36-C37-C40
4	B	301	2CV	N33-C36-C37-O47
4	B	301	2CV	C36-C37-C40-C41
4	B	301	2CV	C36-C37-C40-O49
4	B	301	2CV	O47-C37-C40-C41
4	B	301	2CV	O47-C37-C40-O49
4	B	302	2CV	N33-C36-C37-C40
4	B	302	2CV	N33-C36-C37-O47
4	B	302	2CV	C36-C37-C40-C41
4	B	302	2CV	C36-C37-C40-O49
4	B	302	2CV	O47-C37-C40-C41
4	B	302	2CV	O47-C37-C40-O49
4	B	302	2CV	O49-C40-C41-C42
4	D	301	2CV	N33-C36-C37-O47
4	D	301	2CV	C36-C37-C40-C41
4	D	301	2CV	C36-C37-C40-O49
4	D	301	2CV	O47-C37-C40-C41
4	D	301	2CV	O47-C37-C40-O49
4	D	301	2CV	C37-C40-C41-C42

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	D	301	2CV	C37-C40-C41-O51
4	D	301	2CV	O49-C40-C41-C42
4	D	301	2CV	O49-C40-C41-O51
4	D	302	2CV	N33-C36-C37-O47
4	D	302	2CV	C36-C37-C40-C41
4	D	302	2CV	C36-C37-C40-O49
4	D	302	2CV	C37-C40-C41-C42
4	D	302	2CV	O49-C40-C41-C42
4	D	302	2CV	O47-C37-C40-C41
4	D	302	2CV	O49-C40-C41-O51
4	D	302	2CV	O47-C37-C40-O49
4	B	302	2CV	C37-C40-C41-C42
4	D	302	2CV	C37-C40-C41-O51
2	C	301	12P	O7-C8-C9-O10
2	B	303	12P	C12-C11-O10-C9
2	A	301	12P	C12-C11-O10-C9
2	A	301	12P	C8-C9-O10-C11
2	C	301	12P	C5-C6-O7-C8
2	A	301	12P	C15-C14-O13-C12
2	B	303	12P	C15-C14-O13-C12
2	C	301	12P	C8-C9-O10-C11
2	B	303	12P	C9-C8-O7-C6
2	B	303	12P	C5-C6-O7-C8
2	A	301	12P	C6-C5-O4-C3
4	D	301	2CV	C37-C36-N33-C35
2	B	303	12P	C8-C9-O10-C11
2	A	301	12P	C2-C3-O4-C5
2	C	301	12P	C12-C11-O10-C9
2	C	301	12P	C2-C3-O4-C5
2	A	301	12P	C11-C12-O13-C14
2	C	301	12P	O4-C5-C6-O7
4	D	301	2CV	O51-C41-C42-C43
2	A	301	12P	O7-C8-C9-O10
2	C	301	12P	C6-C5-O4-C3
4	D	301	2CV	O51-C41-C42-O53
4	D	302	2CV	O51-C41-C42-C43
4	B	301	2CV	C41-C42-C43-O44
2	A	301	12P	O4-C5-C6-O7
4	D	301	2CV	C9-C12-C15-C18
2	B	303	12P	C6-C5-O4-C3
4	B	302	2CV	O49-C40-C41-O51
4	D	302	2CV	O51-C41-C42-O53


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	2CV	1	0
4	B	302	2CV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	92/149 (61%)	0.52	15 (16%) 4 4	22, 32, 110, 163	0
1	B	90/149 (60%)	0.49	17 (18%) 3 3	23, 32, 103, 161	0
1	C	91/149 (61%)	0.51	15 (16%) 4 4	22, 31, 104, 158	0
1	D	91/149 (61%)	0.62	18 (19%) 3 2	24, 32, 107, 159	0
All	All	364/596 (61%)	0.54	65 (17%) 3 3	22, 32, 108, 163	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	214	PHE	10.1
1	C	214	PHE	8.4
1	A	222	MET	7.9
1	D	210	VAL	5.5
1	B	211	LEU	5.3
1	B	131	VAL	5.2
1	B	220	ASP	5.2
1	A	131	VAL	5.0
1	C	218	ILE	4.8
1	A	211	LEU	4.8
1	C	217	ILE	4.8
1	D	217	ILE	4.7
1	A	220	ASP	4.7
1	A	214	PHE	4.4
1	D	211	LEU	4.1
1	D	218	ILE	4.0
1	D	132	GLY	3.9
1	D	215	ILE	3.9
1	B	214	PHE	3.9
1	C	215	ILE	3.9
1	C	213	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	130	GLY	3.9
1	C	130	GLY	3.8
1	C	210	VAL	3.8
1	B	208	PHE	3.8
1	A	218	ILE	3.7
1	B	204	MET	3.7
1	A	221	ALA	3.6
1	C	131	VAL	3.6
1	C	219	VAL	3.6
1	A	204	MET	3.5
1	D	219	VAL	3.5
1	D	220	ASP	3.5
1	A	207	THR	3.5
1	A	215	ILE	3.4
1	A	219	VAL	3.4
1	A	217	ILE	3.4
1	B	219	VAL	3.4
1	B	218	ILE	3.3
1	C	207	THR	3.3
1	D	131	VAL	3.3
1	B	207	THR	3.2
1	D	207	THR	3.0
1	C	220	ASP	3.0
1	C	132	GLY	2.9
1	A	208	PHE	2.8
1	D	209	THR	2.8
1	C	211	LEU	2.7
1	D	213	LEU	2.7
1	B	140	VAL	2.7
1	C	212	ASN	2.7
1	B	215	ILE	2.6
1	B	217	ILE	2.5
1	B	144	ILE	2.4
1	C	209	THR	2.4
1	D	208	PHE	2.3
1	D	133	SER	2.3
1	B	137	LEU	2.2
1	D	143	TYR	2.2
1	D	134	VAL	2.2
1	B	132	GLY	2.2
1	B	134	VAL	2.1
1	A	216	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	141	VAL	2.0
1	B	159	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	12P	C	301	16/37	0.81	0.27	67,75,84,84	0
3	NA	A	303	1/1	0.86	0.30	58,58,58,58	1
4	2CV	B	301	26/26	0.88	0.18	43,59,71,72	0
2	12P	B	303	16/37	0.89	0.19	61,65,67,68	0
4	2CV	B	302	26/26	0.89	0.17	42,56,64,67	0
2	12P	A	301	16/37	0.90	0.17	56,67,70,71	0
3	NA	C	303	1/1	0.90	0.24	47,47,47,47	1
4	2CV	D	301	26/26	0.90	0.17	45,66,74,75	0
4	2CV	D	302	26/26	0.90	0.19	47,70,79,82	0
3	NA	A	304	1/1	0.92	0.21	62,62,62,62	1
3	NA	A	302	1/1	0.97	0.39	67,67,67,67	1
3	NA	C	302	1/1	0.98	0.03	23,23,23,23	1
3	NA	D	303	1/1	0.99	0.26	70,70,70,70	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 12P C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 12P B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV B 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 12P A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)


**Electron density around 2CV D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 2CV D 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.