



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 04:32 PM UTC

PDB ID : 6BZN / pdb_00006bzn
Title : Crystal structure of halogenase PltM
Authors : Pang, A.H.; Garneau-Tsodikova, S.; Tsodikov, O.V.
Deposited on : 2017-12-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	522	
1	B	522	
1	C	522	
1	D	522	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	603	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Halogenase PltM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	501	Total	C	N	O	S	0	3	0
			3983	2540	704	728	11			
1	B	499	Total	C	N	O	S	0	1	0
			3942	2515	697	719	11			
1	C	500	Total	C	N	O	S	0	3	0
			3971	2532	701	727	11			
1	D	498	Total	C	N	O	S	0	2	0
			3956	2524	700	721	11			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP Q4KCZ3
A	-18	GLY	-	expression tag	UNP Q4KCZ3
A	-17	SER	-	expression tag	UNP Q4KCZ3
A	-16	SER	-	expression tag	UNP Q4KCZ3
A	-15	HIS	-	expression tag	UNP Q4KCZ3
A	-14	HIS	-	expression tag	UNP Q4KCZ3
A	-13	HIS	-	expression tag	UNP Q4KCZ3
A	-12	HIS	-	expression tag	UNP Q4KCZ3
A	-11	HIS	-	expression tag	UNP Q4KCZ3
A	-10	HIS	-	expression tag	UNP Q4KCZ3
A	-9	SER	-	expression tag	UNP Q4KCZ3
A	-8	SER	-	expression tag	UNP Q4KCZ3
A	-7	GLY	-	expression tag	UNP Q4KCZ3
A	-6	LEU	-	expression tag	UNP Q4KCZ3
A	-5	VAL	-	expression tag	UNP Q4KCZ3
A	-4	PRO	-	expression tag	UNP Q4KCZ3
A	-3	ARG	-	expression tag	UNP Q4KCZ3
A	-2	GLY	-	expression tag	UNP Q4KCZ3
A	-1	SER	-	expression tag	UNP Q4KCZ3
A	0	HIS	-	expression tag	UNP Q4KCZ3
B	-19	MET	-	initiating methionine	UNP Q4KCZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP Q4KCZ3
B	-17	SER	-	expression tag	UNP Q4KCZ3
B	-16	SER	-	expression tag	UNP Q4KCZ3
B	-15	HIS	-	expression tag	UNP Q4KCZ3
B	-14	HIS	-	expression tag	UNP Q4KCZ3
B	-13	HIS	-	expression tag	UNP Q4KCZ3
B	-12	HIS	-	expression tag	UNP Q4KCZ3
B	-11	HIS	-	expression tag	UNP Q4KCZ3
B	-10	HIS	-	expression tag	UNP Q4KCZ3
B	-9	SER	-	expression tag	UNP Q4KCZ3
B	-8	SER	-	expression tag	UNP Q4KCZ3
B	-7	GLY	-	expression tag	UNP Q4KCZ3
B	-6	LEU	-	expression tag	UNP Q4KCZ3
B	-5	VAL	-	expression tag	UNP Q4KCZ3
B	-4	PRO	-	expression tag	UNP Q4KCZ3
B	-3	ARG	-	expression tag	UNP Q4KCZ3
B	-2	GLY	-	expression tag	UNP Q4KCZ3
B	-1	SER	-	expression tag	UNP Q4KCZ3
B	0	HIS	-	expression tag	UNP Q4KCZ3
C	-19	MET	-	initiating methionine	UNP Q4KCZ3
C	-18	GLY	-	expression tag	UNP Q4KCZ3
C	-17	SER	-	expression tag	UNP Q4KCZ3
C	-16	SER	-	expression tag	UNP Q4KCZ3
C	-15	HIS	-	expression tag	UNP Q4KCZ3
C	-14	HIS	-	expression tag	UNP Q4KCZ3
C	-13	HIS	-	expression tag	UNP Q4KCZ3
C	-12	HIS	-	expression tag	UNP Q4KCZ3
C	-11	HIS	-	expression tag	UNP Q4KCZ3
C	-10	HIS	-	expression tag	UNP Q4KCZ3
C	-9	SER	-	expression tag	UNP Q4KCZ3
C	-8	SER	-	expression tag	UNP Q4KCZ3
C	-7	GLY	-	expression tag	UNP Q4KCZ3
C	-6	LEU	-	expression tag	UNP Q4KCZ3
C	-5	VAL	-	expression tag	UNP Q4KCZ3
C	-4	PRO	-	expression tag	UNP Q4KCZ3
C	-3	ARG	-	expression tag	UNP Q4KCZ3
C	-2	GLY	-	expression tag	UNP Q4KCZ3
C	-1	SER	-	expression tag	UNP Q4KCZ3
C	0	HIS	-	expression tag	UNP Q4KCZ3
D	-19	MET	-	initiating methionine	UNP Q4KCZ3
D	-18	GLY	-	expression tag	UNP Q4KCZ3
D	-17	SER	-	expression tag	UNP Q4KCZ3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP Q4KCZ3
D	-15	HIS	-	expression tag	UNP Q4KCZ3
D	-14	HIS	-	expression tag	UNP Q4KCZ3
D	-13	HIS	-	expression tag	UNP Q4KCZ3
D	-12	HIS	-	expression tag	UNP Q4KCZ3
D	-11	HIS	-	expression tag	UNP Q4KCZ3
D	-10	HIS	-	expression tag	UNP Q4KCZ3
D	-9	SER	-	expression tag	UNP Q4KCZ3
D	-8	SER	-	expression tag	UNP Q4KCZ3
D	-7	GLY	-	expression tag	UNP Q4KCZ3
D	-6	LEU	-	expression tag	UNP Q4KCZ3
D	-5	VAL	-	expression tag	UNP Q4KCZ3
D	-4	PRO	-	expression tag	UNP Q4KCZ3
D	-3	ARG	-	expression tag	UNP Q4KCZ3
D	-2	GLY	-	expression tag	UNP Q4KCZ3
D	-1	SER	-	expression tag	UNP Q4KCZ3
D	0	HIS	-	expression tag	UNP Q4KCZ3

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Ca 2 2	0	0
2	B	1	Total Ca 1 1	0	0
2	C	2	Total Ca 2 2	0	0
2	D	2	Total Ca 2 2	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	510	Total	O	0	0
			510	510		
4	B	423	Total	O	0	0
			423	423		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	555	Total 555	O 555	0	0
4	D	527	Total 527	O 527	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Halogenase PltM



- Molecule 1: Halogenase PltM



- Molecule 1: Halogenase PltM



- Molecule 1: Halogenase PltM



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.23Å 157.11Å 214.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.88 – 1.80 39.88 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.88-1.80) 99.6 (39.88-1.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.161 , 0.189 (Not available) , 0.182	Depositor DCC
R_{free} test set	10092 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	17934	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/4083	0.78	0/5544
1	B	0.56	0/4041	0.77	0/5486
1	C	0.61	1/4071 (0.0%)	0.78	0/5528
1	D	0.58	0/4055	0.78	0/5503
All	All	0.58	1/16250 (0.0%)	0.78	0/22061

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	107	VAL	C-O	-5.02	1.19	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3983	0	3922	6	0
1	B	3942	0	3877	3	0
1	C	3971	0	3903	8	0
1	D	3956	0	3899	8	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	18	0	24	0	0
3	B	12	0	15	0	0
3	C	18	0	24	0	0
3	D	12	0	16	0	0
4	A	510	0	0	1	0
4	B	423	0	0	1	0
4	C	555	0	0	2	0
4	D	527	0	0	1	0
All	All	17934	0	15680	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:39[B]:ARG:HG2	1:D:39[B]:ARG:HH11	1.22	1.03
1:D:39[B]:ARG:HH11	1:D:39[B]:ARG:CG	1.90	0.82
1:D:39[B]:ARG:HG2	1:D:39[B]:ARG:NH1	2.00	0.75
1:D:39[B]:ARG:CG	1:D:39[B]:ARG:NH1	2.46	0.75
1:D:39[B]:ARG:HG2	1:D:39[B]:ARG:O	2.04	0.56
1:D:39[B]:ARG:NH2	1:D:124:ASP:OD2	2.43	0.52
1:D:36:GLN:OE1	1:D:141:ASN:ND2	2.43	0.51
1:B:462:ARG:NH1	4:B:703:HOH:O	2.43	0.51
1:C:92:PHE:O	1:C:106:LEU:HA	2.14	0.47
1:C:106:LEU:C	1:C:106:LEU:HD12	2.39	0.47
1:C:185:LEU:HB2	4:C:706:HOH:O	2.14	0.47
1:C:493:ARG:NH1	4:C:705:HOH:O	2.47	0.47
1:C:350[A]:ASP:OD2	1:C:356:ARG:NH2	2.50	0.44
1:B:142:ILE:C	1:B:142:ILE:HD12	2.43	0.44
1:A:171:ASP:HB2	1:A:179:LEU:HD12	2.02	0.42
1:A:466:ARG:NH1	1:A:470:GLU:OE1	2.47	0.42
1:A:185:LEU:HB2	4:A:706:HOH:O	2.20	0.41
1:A:311:ASN:HB2	1:A:318:LEU:HB2	2.02	0.41
1:D:141:ASN:ND2	4:D:710:HOH:O	2.54	0.41
1:A:142:ILE:HD12	1:A:142:ILE:C	2.46	0.41
1:C:171:ASP:OD2	1:C:177:SER:OG	2.29	0.41
1:B:4:TYR:O	1:B:166:ALA:HA	2.21	0.41
1:C:106:LEU:HD11	1:C:498:PHE:CZ	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:215:LEU:HD21	1:A:223:GLU:HA	2.03	0.40
1:C:142:ILE:C	1:C:142:ILE:HD12	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	502/522 (96%)	491 (98%)	11 (2%)	0	100	100
1	B	496/522 (95%)	485 (98%)	11 (2%)	0	100	100
1	C	501/522 (96%)	491 (98%)	10 (2%)	0	100	100
1	D	496/522 (95%)	490 (99%)	6 (1%)	0	100	100
All	All	1995/2088 (96%)	1957 (98%)	38 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	425/440 (97%)	422 (99%)	3 (1%)	76	73
1	B	397/440 (90%)	391 (98%)	6 (2%)	57	49
1	C	112/440 (26%)	110 (98%)	2 (2%)	51	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	D	404/440 (92%)	397 (98%)	7 (2%)	53 45
All	All	1338/1760 (76%)	1320 (99%)	18 (1%)	61 54

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	ARG
1	A	224	LEU
1	A	305	ASN
1	B	39	ARG
1	B	87	LYS
1	B	145	GLU
1	B	206	LYS
1	B	305	ASN
1	B	331	SER
1	C	3	GLN
1	C	39	ARG
1	D	39[A]	ARG
1	D	39[B]	ARG
1	D	141	ASN
1	D	182	GLN
1	D	305	ASN
1	D	320	PRO
1	D	413	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	96	GLN
1	A	246	HIS
1	A	321	GLN
1	A	376	ASN
1	A	460	GLN
1	A	473	ASN
1	B	2	ASN
1	B	358	GLN
1	B	376	ASN
1	C	2	ASN
1	D	28	ASN
1	D	36	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	141	ASN
1	D	259	GLN
1	D	321	GLN
1	D	376	ASN
1	D	378	GLN
1	D	412	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 7 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	B	603	-	5,5,5	0.48	0	5,5,5	1.85	3 (60%)
3	GOL	A	604	-	5,5,5	0.67	0	5,5,5	1.17	0
3	GOL	A	603	-	5,5,5	0.24	0	5,5,5	0.62	0
3	GOL	D	603	-	5,5,5	0.66	0	5,5,5	1.06	0
3	GOL	B	602	-	5,5,5	0.45	0	5,5,5	0.95	0
3	GOL	D	604	-	5,5,5	0.58	0	5,5,5	0.60	0
3	GOL	C	605	-	5,5,5	0.36	0	5,5,5	0.28	0
3	GOL	C	604	-	5,5,5	0.45	0	5,5,5	1.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	C	603	-	5,5,5	0.56	0	5,5,5	0.53	0
3	GOL	A	605	-	5,5,5	0.46	0	5,5,5	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	603	-	-	3/4/4/4	-
3	GOL	A	604	-	-	1/4/4/4	-
3	GOL	A	603	-	-	0/4/4/4	-
3	GOL	D	603	-	-	0/4/4/4	-
3	GOL	B	602	-	-	1/4/4/4	-
3	GOL	D	604	-	-	2/4/4/4	-
3	GOL	C	605	-	-	2/4/4/4	-
3	GOL	C	604	-	-	0/4/4/4	-
3	GOL	C	603	-	-	2/4/4/4	-
3	GOL	A	605	-	-	4/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	GOL	O2-C2-C1	2.97	121.49	109.18
3	B	603	GOL	C3-C2-C1	2.00	119.15	111.80
3	B	603	GOL	O2-C2-C3	2.00	117.47	109.18

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	603	GOL	C1-C2-C3-O3
3	C	605	GOL	O1-C1-C2-C3
3	A	605	GOL	O1-C1-C2-C3
3	A	605	GOL	C1-C2-C3-O3
3	C	603	GOL	O1-C1-C2-C3
3	C	605	GOL	O1-C1-C2-O2
3	B	603	GOL	O1-C1-C2-O2
3	B	603	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	602	GOL	O2-C2-C3-O3
3	A	605	GOL	O2-C2-C3-O3
3	D	604	GOL	O1-C1-C2-C3
3	D	604	GOL	O1-C1-C2-O2
3	A	605	GOL	O1-C1-C2-O2
3	C	603	GOL	O1-C1-C2-O2
3	A	604	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/522 (95%)	-0.29	10 (1%) 65 65	7, 17, 31, 48	3 (0%)
1	B	499/522 (95%)	-0.09	9 (1%) 67 68	10, 20, 39, 59	1 (0%)
1	C	500/522 (95%)	-0.36	9 (1%) 67 68	8, 15, 30, 48	3 (0%)
1	D	498/522 (95%)	-0.24	4 (0%) 82 83	8, 17, 35, 58	2 (0%)
All	All	1998/2088 (95%)	-0.24	32 (1%) 70 71	7, 18, 35, 59	9 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	ALA	5.1
1	C	174	ALA	5.1
1	D	176	GLY	4.8
1	B	174	ALA	3.8
1	C	2	ASN	3.8
1	A	2	ASN	3.5
1	A	175	GLN	3.5
1	A	178	PRO	3.4
1	C	178	PRO	3.2
1	C	3	GLN	3.2
1	C	176	GLY	3.1
1	A	502	VAL	3.0
1	A	177	SER	3.0
1	B	2	ASN	3.0
1	D	3[A]	GLN	2.8
1	B	500	LEU	2.7
1	B	176	GLY	2.6
1	B	461	ALA	2.6
1	B	416	HIS	2.6
1	C	179	LEU	2.6
1	B	350	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	175	GLN	2.4
1	D	2	ASN	2.4
1	D	160	ASN	2.4
1	A	112	LYS	2.3
1	A	179	LEU	2.3
1	B	501	LYS	2.3
1	C	177	SER	2.2
1	C	106	LEU	2.2
1	A	180	SER	2.1
1	B	160	ASN	2.1
1	A	420	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	603	6/6	0.72	0.16	46,47,48,48	0
3	GOL	B	603	6/6	0.72	0.18	52,52,53,53	0
3	GOL	C	605	6/6	0.79	0.17	44,49,50,52	0
3	GOL	A	605	6/6	0.85	0.12	32,35,36,38	0
3	GOL	D	604	6/6	0.87	0.12	31,35,36,38	0
3	GOL	C	603	6/6	0.89	0.12	24,27,29,30	0
2	CA	A	602	1/1	0.93	0.11	36,36,36,36	0
2	CA	D	602	1/1	0.93	0.13	36,36,36,36	0
3	GOL	D	603	6/6	0.94	0.08	16,17,18,18	0
2	CA	C	602	1/1	0.94	0.12	34,34,34,34	0
2	CA	B	601	1/1	0.95	0.08	25,25,25,25	0
3	GOL	A	604	6/6	0.95	0.07	16,17,18,18	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	C	604	6/6	0.97	0.06	14,15,15,16	0
3	GOL	B	602	6/6	0.97	0.06	17,19,20,20	0
2	CA	D	601	1/1	0.99	0.05	18,18,18,18	0
2	CA	C	601	1/1	0.99	0.05	18,18,18,18	0
2	CA	A	601	1/1	0.99	0.05	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.