



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 11:38 AM UTC

PDB ID : 4C0D / pdb_00004c0d
Title : Structure of the NOT module of the human CCR4-NOT complex (CNOT1-CNOT2-CNOT3)
Authors : Raisch, T.; Jonas, S.; Boland, A.; Chen, Y.; Izaurralde, E.; Weichenrieder, O.
Deposited on : 2013-08-01
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	812	
2	B	201	
3	C	166	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	512	4096	2653	702	720	21	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1560	GLY	-	expression tag	UNP A5YKK6
A	1561	HIS	-	expression tag	UNP A5YKK6
A	1562	MET	-	expression tag	UNP A5YKK6
A	1563	LEU	-	expression tag	UNP A5YKK6
A	1564	GLU	-	expression tag	UNP A5YKK6

- Molecule 2 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	191	1558	1008	259	283	8	0	0	0

There are 4 discrepancies between the modelled and reference sequences:




Chain	Residue	Modelled	Actual	Comment	Reference
B	340	GLY	-	expression tag	UNP Q9NZN8
B	341	PRO	-	expression tag	UNP Q9NZN8
B	342	HIS	-	expression tag	UNP Q9NZN8
B	343	MET	-	expression tag	UNP Q9NZN8

- Molecule 3 is a protein called CCR4-NOT TRANSCRIPTION COMPLEX SUBUNIT 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	142	1210	789	198	217	6	0	0	0

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	588	MET	-	expression tag	UNP O75175
C	589	GLY	-	expression tag	UNP O75175
C	590	SER	-	expression tag	UNP O75175
C	591	SER	-	expression tag	UNP O75175
C	592	HIS	-	expression tag	UNP O75175
C	593	HIS	-	expression tag	UNP O75175
C	594	HIS	-	expression tag	UNP O75175
C	595	HIS	-	expression tag	UNP O75175
C	596	HIS	-	expression tag	UNP O75175
C	597	HIS	-	expression tag	UNP O75175
C	598	SER	-	expression tag	UNP O75175
C	599	SER	-	expression tag	UNP O75175
C	600	GLY	-	expression tag	UNP O75175
C	601	THR	-	expression tag	UNP O75175
C	602	GLY	-	expression tag	UNP O75175
C	603	SER	-	expression tag	UNP O75175
C	604	GLY	-	expression tag	UNP O75175
C	605	HIS	-	expression tag	UNP O75175
C	606	MET	-	expression tag	UNP O75175



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	91.57Å 165.92Å 78.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.48 – 3.20 48.48 – 3.20	Depositor EDS
% Data completeness (in resolution range)	98.8 (48.48-3.20) 98.9 (48.48-3.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.224 , 0.273 0.221 , 0.269	Depositor DCC
R_{free} test set	1035 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	62.8	Xtrriage
Anisotropy	0.751	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6864	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.27	0/4206	0.71	0/5738
2	B	0.25	0/1606	0.67	0/2186
3	C	0.26	0/1259	0.69	2/1714 (0.1%)
All	All	0.27	0/7071	0.70	2/9638 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	653	PRO	CA-C-N	5.21	123.46	119.66
3	C	653	PRO	C-N-CA	5.21	123.46	119.66

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4096	0	4038	32	0
2	B	1558	0	1489	18	0
3	C	1210	0	1090	14	0
All	All	6864	0	6617	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their

clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:436:LEU:HB3	2:B:463:GLU:HG3	1.80	0.63
2:B:424:ASN:HD21	3:C:636:GLN:HG2	1.66	0.60
3:C:674:ILE:HG23	3:C:678:LEU:HD12	1.83	0.59
1:A:2163:ARG:NH1	1:A:2276:TYR:OH	2.36	0.58
2:B:534:ASN:HB3	2:B:537:GLN:HG3	1.88	0.55

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/812 (63%)	488 (96%)	21 (4%)	1 (0%)	43	73
2	B	189/201 (94%)	181 (96%)	8 (4%)	0	100	100
3	C	140/166 (84%)	134 (96%)	6 (4%)	0	100	100
All	All	839/1179 (71%)	803 (96%)	35 (4%)	1 (0%)	48	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1866	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	441/711 (62%)	437 (99%)	4 (1%)	70	81
2	B	164/175 (94%)	159 (97%)	5 (3%)	36	66
3	C	123/151 (82%)	123 (100%)	0	100	100
All	All	728/1037 (70%)	719 (99%)	9 (1%)	63	79

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	500	TYR
2	B	538	GLN
1	A	2241	THR
2	B	363	PHE
2	B	387	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	451	ASN
2	B	426	HIS
1	A	2231	HIS
2	B	411	GLN
1	A	2199	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	512/812 (63%)	0.10	9 (1%) 67 48	36, 58, 111, 145	0
2	B	191/201 (95%)	0.36	4 (2%) 63 43	54, 77, 112, 132	0
3	C	142/166 (85%)	0.22	2 (1%) 73 54	46, 80, 145, 157	0
All	All	845/1179 (71%)	0.18	15 (1%) 67 48	36, 65, 121, 157	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	540	PHE	3.3
3	C	715	ILE	2.7
1	A	2149	ASN	2.5
1	A	2152	VAL	2.5
2	B	531	PHE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.