



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:10 PM UTC

PDB ID : 5C1D / pdb_00005c1d
Title : Human OGT in complex with UDP-5S-GlcNAc and substrate peptide (RB2L)
Authors : Schimpl, M.; Rafie, K.; van Aalten, D.M.F.
Deposited on : 2015-06-13
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	723	
2	C	8	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	1102	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	695	Total	C	N	O	S	0	3	0
			5515	3497	966	1015	37			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	309	GLY	-	expression tag	UNP O15294
A	310	PRO	-	expression tag	UNP O15294
A	311	GLY	-	expression tag	UNP O15294
A	312	SER	-	expression tag	UNP O15294


- Molecule 2 is a protein called Retinoblastoma-like protein 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	0	0	1
			47	29	8	10			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	8	ALA	THR	conflict	UNP Q08999

- Molecule 3 is (2S,3R,4R,5S,6R)-3-(acetylamino)-4,5-dihydroxy-6-(hydroxymethyl)tetrahydro-2H-thiopyran-2-yl [(2R,3S,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methyl dihydrogen diphosphate (CCD ID: 12V) (formula: C₁₇H₂₇N₃O₁₆P₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	S	0	0
			39	17	3	16	2	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	251	Total 251	O 251	0	0
5	C	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-N-acetylglucosamine--peptide N-acetylglucosaminyltransferase 110 kDa subunit



- Molecule 2: Retinoblastoma-like protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	138.53Å 151.61Å 200.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.20 – 2.05 46.20 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.20-2.05) 100.0 (46.20-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.195 , 0.229 0.205 , 0.235	Depositor DCC
R_{free} test set	3160 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5865	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, 12V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/5641	0.85	2/7649 (0.0%)
2	C	0.94	0/47	1.11	0/66
All	All	0.70	0/5688	0.85	2/7715 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	763	ASN	N-CA-C	7.02	119.74	109.07
1	A	991	ARG	CG-CD-NE	-5.62	99.63	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5515	0	5484	32	1
2	C	47	0	51	3	0
3	A	39	0	25	7	0
4	A	10	0	0	2	1
5	A	251	0	0	2	2
5	C	3	0	0	0	0
All	All	5865	0	5560	38	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1101:12V:C1'	3:A:1101:12V:O1'	1.67	1.43
1:A:859:PRO:O	1:A:860:ASN:ND2	2.00	0.94
3:A:1101:12V:H1'	3:A:1101:12V:O1A	1.79	0.82
1:A:859:PRO:O	1:A:860:ASN:CG	2.29	0.76
3:A:1101:12V:C1'	3:A:1101:12V:PB	2.74	0.75
3:A:1101:12V:S5'	2:C:5:SER:OG	2.46	0.74
1:A:850:MET:HE1	1:A:966:TYR:HB3	1.75	0.69
1:A:654:GLY:HA2	3:A:1101:12V:H6'A	1.79	0.64
1:A:315:THR:N	5:A:1202:HOH:O	2.31	0.63
1:A:432:SER:HB2	4:A:1102:PO4:O3	2.00	0.61
1:A:895:VAL:HG11	2:C:2:THR:HG21	1.84	0.59
1:A:895:VAL:CG1	2:C:2:THR:HG21	2.33	0.58
1:A:321:ASN:HD21	1:A:356:ASN:HD22	1.53	0.56
1:A:1021:ASN:HD22	1:A:1021:ASN:H	1.55	0.54
3:A:1101:12V:H1'	3:A:1101:12V:PB	2.48	0.52
1:A:1015:GLU:OE2	1:A:1015:GLU:HA	2.10	0.52
1:A:429:HIS:ND1	4:A:1102:PO4:O1	2.44	0.51
3:A:1101:12V:H6'	5:A:1336:HOH:O	2.11	0.49
1:A:859:PRO:O	1:A:860:ASN:CB	2.61	0.49
1:A:467:CYS:SG	1:A:876:ILE:HD11	2.53	0.48
1:A:961:LYS:HG3	1:A:965:GLU:OE1	2.15	0.46
1:A:1022:LYS:HB3	1:A:1023:PRO:HD2	1.99	0.45
1:A:547:ARG:HB2	1:A:624:HIS:HD2	1.82	0.45
1:A:558:HIS:CG	1:A:559:PRO:HD2	2.53	0.44
1:A:435:ILE:N	1:A:436:PRO:CD	2.80	0.44
1:A:669:THR:OG1	1:A:670:ASP:N	2.45	0.44
1:A:850:MET:SD	1:A:963:ARG:NH1	2.92	0.43
1:A:850:MET:HE1	1:A:966:TYR:CB	2.47	0.42
1:A:515:GLU:OE1	1:A:516:ARG:NH1	2.50	0.42
1:A:524:LYS:O	1:A:527:VAL:HG12	2.19	0.42
1:A:1021:ASN:HD22	1:A:1021:ASN:N	2.17	0.42
1:A:877:GLN:O	1:A:881:GLN:HG2	2.20	0.41
1:A:471:ASP:HB3	1:A:474:GLU:CG	2.50	0.41
1:A:1015:GLU:OE2	1:A:1015:GLU:CA	2.69	0.41
1:A:547:ARG:H	1:A:624:HIS:CD2	2.39	0.41
1:A:471:ASP:O	1:A:474:GLU:HG2	2.21	0.40
1:A:479:LEU:O	1:A:483:VAL:HG23	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1018:ALA:C	1:A:1020:GLY:H	2.29	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1344:HOH:O	5:A:1406:HOH:O[3_555]	1.64	0.56
4:A:1102:PO4:O2	4:A:1102:PO4:O2[8_544]	1.78	0.42
1:A:620:GLN:OE1	1:A:620:GLN:OE1[11_554]	1.82	0.38
5:A:1278:HOH:O	5:A:1406:HOH:O[3_555]	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	692/723 (96%)	671 (97%)	19 (3%)	2 (0%)	36	30
2	C	6/8 (75%)	6 (100%)	0	0	100	100
All	All	698/731 (96%)	677 (97%)	19 (3%)	2 (0%)	36	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	860	ASN
1	A	1019	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	600/618 (97%)	583 (97%)	17 (3%)	38	34
2	C	6/6 (100%)	5 (83%)	1 (17%)	2	0
All	All	606/624 (97%)	588 (97%)	18 (3%)	36	32

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	430	LYS
1	A	501	MET
1	A	519	ASN
1	A	534	GLU
1	A	541	LEU
1	A	567	ILE
1	A	677	GLU
1	A	715	SER
1	A	742	LYS
1	A	778	GLU
1	A	887	GLN
1	A	914	THR
1	A	926	VAL
1	A	961	LYS
1	A	1008	ARG
1	A	1021	ASN
1	A	1027	ILE
2	C	2	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	325	ASN
1	A	356	ASN
1	A	364	GLN
1	A	565	GLN
1	A	624	HIS
1	A	727	ASN
1	A	763	ASN
1	A	906	GLN
1	A	1021	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	12V	A	1101	-	38,41,41	1.87	9 (23%)	52,62,62	1.98	8 (15%)
4	PO4	A	1103	-	4,4,4	0.73	0	6,6,6	0.77	0
4	PO4	A	1102	-	4,4,4	1.01	0	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	12V	A	1101	-	-	2/25/63/63	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	12V	C4'-C5'	7.65	1.60	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	12V	C3'-C2'	3.43	1.59	1.53
3	A	1101	12V	C8'-C7'	2.59	1.55	1.50
3	A	1101	12V	O3'-C3'	2.46	1.49	1.43
3	A	1101	12V	C6'-C5'	2.44	1.54	1.52
3	A	1101	12V	C2'-N2'	2.32	1.49	1.45
3	A	1101	12V	C5-C4	-2.20	1.38	1.43
3	A	1101	12V	O4B-C1B	2.09	1.46	1.42
3	A	1101	12V	C2-N1	-2.00	1.35	1.38

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1101	12V	O1'-C1'-C2'	7.08	118.28	107.52
3	A	1101	12V	C4-N3-C2	-5.33	119.99	126.61
3	A	1101	12V	N3-C2-N1	4.94	121.32	114.89
3	A	1101	12V	C5-C4-N3	3.64	119.90	114.80
3	A	1101	12V	C6'-C5'-C4'	-3.58	99.22	114.44
3	A	1101	12V	O4B-C1B-N1	-3.42	100.62	108.36
3	A	1101	12V	O3'-C3'-C2'	3.02	115.58	109.58
3	A	1101	12V	O2-C2-N3	-3.00	115.96	121.49

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1101	12V	C3B-C4B-C5B-O5B
3	A	1101	12V	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1101	12V	7	0
4	A	1102	PO4	2	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	695/723 (96%)	0.44	44 (6%) 26 26	15, 42, 72, 87	5 (0%)
2	C	8/8 (100%)	0.91	1 (12%) 8 7	33, 44, 49, 66	0
All	All	703/731 (96%)	0.44	45 (6%) 25 25	15, 42, 72, 87	5 (0%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	329	GLU	10.0
2	C	1	VAL	5.6
1	A	762	LEU	4.7
1	A	1027	ILE	4.4
1	A	715	SER	4.4
1	A	713	PHE	3.9
1	A	860	ASN	3.2
1	A	714	LYS	3.0
1	A	680	GLU	3.0
1	A	527	VAL	2.8
1	A	342	LYS	2.8
1	A	531	PRO	2.8
1	A	1018	ALA	2.7
1	A	529	HIS	2.5
1	A	317	ALA	2.5
1	A	677	GLU	2.4
1	A	740	ASP	2.4
1	A	474	GLU	2.4
1	A	1026	MET	2.4
1	A	1019	ALA	2.4
1	A	885	LEU	2.4
1	A	326	ILE	2.4
1	A	534	GLU	2.3
1	A	1016	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	315	THR	2.3
1	A	887	GLN	2.3
1	A	620	GLN	2.2
1	A	347	PHE	2.2
1	A	323	LEU	2.2
1	A	886	PRO	2.2
1	A	939	GLU	2.2
1	A	521	CYS	2.2
1	A	662	LEU	2.2
1	A	614	ALA	2.1
1	A	520	LEU	2.1
1	A	528	LEU	2.1
1	A	1017	TYR	2.1
1	A	532	PRO	2.1
1	A	625	ILE	2.1
1	A	346	VAL	2.1
1	A	539	LEU	2.0
1	A	881	GLN	2.0
1	A	769	MET	2.0
1	A	610	CYS	2.0
1	A	1025	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	A	1102	5/5	0.85	0.24	56,97,140,145	1
4	PO4	A	1103	5/5	0.86	0.16	78,84,91,91	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	12V	A	1101	39/39	0.96	0.10	24,28,63,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.