



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:14 PM UTC

PDB ID : 8C2P / pdb_00008c2p
Title : FMDV 3D polymerase in complex with 3B3
Authors : Ferrer-Orta, C.; Veraguer, N.
Deposited on : 2022-12-22
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	14% 84% 14%
2	B	24	25% 25% 75%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase 3D-POL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	468	Total	C	N	O	S	0	0	0
			3673	2339	633	680	21			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	471	ALA	-	expression tag	UNP P03311
A	472	ALA	-	expression tag	UNP P03311
A	473	LEU	-	expression tag	UNP P03311
A	474	GLU	-	expression tag	UNP P03311
A	475	HIS	-	expression tag	UNP P03311
A	476	HIS	-	expression tag	UNP P03311
A	477	HIS	-	expression tag	UNP P03311
A	478	HIS	-	expression tag	UNP P03311
A	479	HIS	-	expression tag	UNP P03311
A	480	HIS	-	expression tag	UNP P03311

- Molecule 2 is a protein called Protein 3B-3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	0	0	0
			31	19	6	6			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	51	Total	O	0	0
			51	51		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-directed RNA polymerase 3D-POL



• Molecule 2: Protein 3B-3



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.51Å 93.51Å 100.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.76 – 1.85 46.76 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.76-1.85) 99.3 (46.76-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.20rc4_4425	Depositor
R, R_{free}	0.243 , 0.262 0.243 , 0.265	Depositor DCC
R_{free} test set	2046 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 29.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3755	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/3762 (0.0%)	0.64	5/5099 (0.1%)
2	B	0.59	0/31	0.79	0/42
All	All	0.47	1/3793 (0.0%)	0.64	5/5141 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	355	PRO	N-CD	-14.00	1.28	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	GLU	N-CA-C	-8.03	96.62	109.39
1	A	238	ASP	O-C-N	5.96	130.40	123.30
1	A	355	PRO	N-CD-CG	5.73	111.79	103.20
1	A	238	ASP	CA-C-N	-5.63	115.29	123.06
1	A	238	ASP	C-N-CA	-5.63	115.29	123.06

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3673	0	3580	55	0
2	B	31	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	51	0	0	0	0
All	All	3755	0	3596	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:HIS:HB2	1:A:356:HIS:CE1	1.90	1.06
1:A:18:LYS:O	1:A:39:LEU:HD12	1.74	0.87
1:A:322:HIS:HB2	1:A:356:HIS:HE1	1.46	0.81
1:A:47:ASN:HB2	1:A:50:VAL:CG2	2.19	0.72
1:A:18:LYS:O	1:A:39:LEU:HB2	1.90	0.72
1:A:322:HIS:CB	1:A:356:HIS:CE1	2.72	0.70
1:A:47:ASN:HB2	1:A:50:VAL:HG23	1.73	0.69
1:A:322:HIS:CB	1:A:356:HIS:HE1	2.06	0.68
1:A:322:HIS:HB2	1:A:356:HIS:ND1	2.09	0.68
1:A:147:LEU:O	1:A:151:GLU:HG3	1.93	0.68
1:A:115:THR:O	1:A:128:ARG:HD3	1.95	0.66
1:A:16:MET:HG3	1:A:39:LEU:HD13	1.80	0.64
1:A:17:ARG:HG3	1:A:17:ARG:O	1.97	0.63
1:A:121:TRP:CH2	1:A:142:GLU:HB3	2.34	0.62
1:A:15:VAL:HG22	1:A:16:MET:H	1.66	0.60
1:A:121:TRP:HH2	1:A:142:GLU:HB3	1.65	0.60
1:A:189:ILE:HG23	1:A:301:SER:HB3	1.86	0.57
1:A:15:VAL:HG22	1:A:16:MET:N	2.20	0.57
1:A:6:THR:HG23	1:A:290:ILE:HG12	1.87	0.56
1:A:16:MET:O	1:A:163:LEU:HD12	2.05	0.56
1:A:248:HIS:HE1	1:A:307:ASN:OD1	1.88	0.56
1:A:248:HIS:HD2	1:A:314:TYR:OH	1.91	0.54
1:A:18:LYS:O	1:A:39:LEU:CD1	2.53	0.54
1:A:13:VAL:HG22	1:A:123:LEU:HD21	1.89	0.53
1:A:18:LYS:O	1:A:39:LEU:CB	2.55	0.53
1:A:33:GLU:O	1:A:34:PHE:CD1	2.63	0.52
1:A:121:TRP:CZ3	1:A:131:LEU:HD21	2.46	0.51
1:A:43:ASP:HB3	1:A:46:LEU:HD23	1.93	0.51
1:A:20:LYS:O	1:A:416:ARG:HB3	2.12	0.50
1:A:47:ASN:HB2	1:A:50:VAL:HG21	1.94	0.50
1:A:12:ARG:HG2	1:A:287:ASN:OD1	2.12	0.50
1:A:43:ASP:O	1:A:46:LEU:HG	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:HIS:CD2	1:A:314:TYR:OH	2.66	0.48
1:A:52:LEU:O	1:A:56:ILE:HG23	2.14	0.47
1:A:248:HIS:CE1	1:A:307:ASN:OD1	2.66	0.47
1:A:322:HIS:CA	1:A:356:HIS:HE1	2.27	0.47
1:A:27:HIS:CD2	1:A:32:PRO:HD2	2.51	0.46
1:A:15:VAL:HG21	1:A:285:TYR:CE1	2.51	0.46
1:A:150:MET:HE2	1:A:155:TYR:CE2	2.51	0.46
1:A:168:ARG:NH1	1:A:177:LYS:O	2.47	0.46
1:A:275:LEU:HD22	1:A:306:ILE:HD11	1.99	0.45
1:A:423:LYS:O	1:A:427:VAL:HG23	2.16	0.45
1:A:322:HIS:CA	1:A:356:HIS:CE1	3.00	0.45
1:A:347:ASP:HA	1:A:376:LEU:CD2	2.47	0.45
1:A:411:ILE:O	1:A:423:LYS:NZ	2.49	0.45
1:A:17:ARG:O	1:A:17:ARG:CG	2.65	0.44
1:A:85:LEU:HD11	1:A:203:MET:SD	2.58	0.44
1:A:121:TRP:CE3	1:A:131:LEU:HD21	2.53	0.44
1:A:229:HIS:O	1:A:232:GLN:HG2	2.17	0.43
1:A:421:GLN:O	1:A:425:ILE:HG12	2.20	0.42
1:A:224:GLN:CD	1:A:398:PHE:CD1	2.98	0.42
1:A:296:MET:HE3	1:A:296:MET:HB3	1.75	0.41
1:A:444:GLU:HG3	1:A:445:PRO:HD3	2.02	0.41
1:A:321:ARG:O	1:A:321:ARG:HG3	2.22	0.40
1:A:86:HIS:CD2	1:A:265:PHE:CD2	3.10	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/480 (97%)	456 (98%)	10 (2%)	0	100	100
2	B	4/24 (17%)	2 (50%)	2 (50%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	470/504 (93%)	458 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/403 (96%)	387 (100%)	0	100	100
2	B	1/20 (5%)	1 (100%)	0	100	100
All	All	388/423 (92%)	388 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	HIS
1	A	27	HIS
1	A	248	HIS
1	A	356	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	468/480 (97%)	0.98	68 (14%) 6 5	26, 45, 74, 88	0
2	B	6/24 (25%)	3.84	6 (100%) 0 0	49, 56, 70, 75	0
All	All	474/504 (94%)	1.02	74 (15%) 5 5	26, 46, 74, 88	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	MET	7.2
1	A	46	LEU	6.7
1	A	50	VAL	5.8
1	A	15	VAL	5.6
1	A	52	LEU	5.2
1	A	51	VAL	5.1
1	A	394	TYR	5.0
2	B	5	PRO	5.0
1	A	371	ASP	5.0
2	B	9	PRO	4.6
1	A	19	THR	4.5
1	A	17	ARG	4.5
1	A	368	ASP	4.4
1	A	18	LYS	4.3
1	A	136	ASN	4.3
1	A	370	SER	4.1
2	B	6	VAL	4.0
1	A	20	LYS	4.0
1	A	468	GLY	3.7
1	A	285	TYR	3.7
2	B	7	LYS	3.6
1	A	123	LEU	3.5
1	A	49	GLY	3.5
1	A	372	LYS	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	14	HIS	3.4
2	B	4	GLY	3.3
1	A	373	GLY	3.3
1	A	234	ARG	3.2
1	A	447	GLN	3.1
1	A	398	PHE	2.9
1	A	28	GLY	2.9
1	A	300	CYS	2.9
1	A	42	LYS	2.8
1	A	9	VAL	2.8
1	A	7	ARG	2.8
1	A	416	ARG	2.8
1	A	11	GLU	2.7
1	A	173	VAL	2.7
1	A	21	LEU	2.7
1	A	167	ILE	2.7
1	A	299	GLY	2.6
1	A	122	ALA	2.6
1	A	44	PRO	2.6
1	A	170	MET	2.6
2	B	8	LYS	2.5
1	A	39	LEU	2.4
1	A	1	GLY	2.4
1	A	356	HIS	2.4
1	A	13	VAL	2.4
1	A	467	CYS	2.4
1	A	41	ASN	2.4
1	A	40	SER	2.4
1	A	369	LYS	2.3
1	A	420	ILE	2.3
1	A	10	GLU	2.2
1	A	449	LEU	2.2
1	A	176	GLY	2.2
1	A	139	VAL	2.2
1	A	466	VAL	2.2
1	A	38	ALA	2.2
1	A	55	VAL	2.2
1	A	132	ILE	2.1
1	A	48	GLU	2.1
1	A	134	PHE	2.1
1	A	2	LEU	2.1
1	A	127	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	137	GLY	2.1
1	A	121	TRP	2.1
1	A	114	ASP	2.1
1	A	324	GLU	2.0
1	A	375	VAL	2.0
1	A	244	PHE	2.0
1	A	418	GLY	2.0
1	A	12	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.