



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 11:33 PM UTC

PDB ID : 2C39 / pdb_00002c39
Title : RNase PH core of the archaeal exosome in complex with ADP
Authors : Lorentzen, E.; Conti, E.
Deposited on : 2005-10-05
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 2% 81% 12% • 5%
1	C	275	 % 74% 19% • 5%
1	E	275	 75% 18% • 5%
1	G	275	 3% 78% 15% • 5%
1	I	275	 % 77% 16% • 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	275	 3% 74% 19% • 5%
1	M	275	 2% 80% 12% • 5%
1	O	275	 2% 77% 16% • 5%
1	Q	275	 0% 80% 12% • 5%
1	S	275	 4% 76% 15% • 6%
1	U	275	 9% 76% 15% • 7%
1	W	275	 3% 78% 14% • 6%
2	B	248	 0% 71% 23% • •
2	D	248	 2% 69% 26% •
2	F	248	 2% 70% 25% • •
2	H	248	 3% 67% 23% • 7%
2	J	248	 3% 72% 22% • •
2	L	248	 0% 72% 24% •
2	N	248	 2% 74% 19% • •
2	P	248	 3% 67% 28% 5%
2	R	248	 2% 71% 25% •
2	T	248	 2% 68% 28% •
2	V	248	 5% 71% 22% • •
2	X	248	 5% 74% 21% • •

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 45814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROBABLE EXOSOME COMPLEX EXONUCLEASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	260	Total 1944	C 1239	N 320	O 380	S 5	0	0	0
1	C	260	Total 1947	C 1243	N 321	O 378	S 5	0	0	0
1	E	260	Total 1968	C 1253	N 325	O 385	S 5	0	0	0
1	G	260	Total 1961	C 1250	N 323	O 383	S 5	0	0	0
1	I	260	Total 1954	C 1247	N 323	O 379	S 5	0	0	0
1	K	260	Total 1958	C 1249	N 323	O 381	S 5	0	0	0
1	M	260	Total 1960	C 1249	N 324	O 382	S 5	0	0	0
1	O	260	Total 1951	C 1246	N 323	O 377	S 5	0	0	0
1	Q	260	Total 1953	C 1245	N 322	O 381	S 5	0	0	0
1	S	259	Total 1955	C 1245	N 323	O 382	S 5	0	0	0
1	U	255	Total 1908	C 1219	N 317	O 367	S 5	0	0	0
1	W	259	Total 1950	C 1245	N 322	O 378	S 5	0	0	0

- Molecule 2 is a protein called PROBABLE EXOSOME COMPLEX EXONUCLEASE 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	241	Total 1812	C 1145	N 317	O 340	S 10	0	0	0
2	D	248	Total 1900	C 1198	N 330	O 360	S 12	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	241	Total	C	N	O	S	0	0	0
			1838	1161	317	350	10			
2	H	230	Total	C	N	O	S	0	0	0
			1732	1094	302	327	9			
2	J	241	Total	C	N	O	S	0	0	0
			1844	1165	321	348	10			
2	L	247	Total	C	N	O	S	0	0	0
			1890	1194	328	357	11			
2	N	239	Total	C	N	O	S	0	0	0
			1809	1144	313	342	10			
2	P	248	Total	C	N	O	S	0	0	0
			1904	1202	332	358	12			
2	R	248	Total	C	N	O	S	0	0	0
			1880	1188	332	348	12			
2	T	248	Total	C	N	O	S	0	0	0
			1884	1193	328	351	12			
2	V	239	Total	C	N	O	S	0	0	0
			1815	1148	318	339	10			
2	X	239	Total	C	N	O	S	0	0	0
			1821	1148	315	348	10			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total C N O P 27 10 5 10 2	0	0
3	F	1	Total C O P 17 5 10 2	0	0
3	H	1	Total C N O P 27 10 5 10 2	0	0
3	J	1	Total O P 9 7 2	0	0
3	L	1	Total C N O P 27 10 5 10 2	0	0
3	N	1	Total C N O P 27 10 5 10 2	0	0
3	P	1	Total C N O P 27 10 5 10 2	0	0
3	R	1	Total C N O P 27 10 5 10 2	0	0
3	T	1	Total C N O P 27 10 5 10 2	0	0
3	V	1	Total C O P 17 5 10 2	0	0
3	X	1	Total C O P 17 5 10 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2




- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2




- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2





- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2








- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2






- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2




- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2




- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2





- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2

Chain Q: 





- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2

Chain S: 





- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2

Chain U: 




- Molecule 1: PROBABLE EXOSOME COMPLEX EXONUCLEASE 2

Chain W: 




- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1

Chain B:  71% 23%




• Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1

Chain D:  2% 69% 26%




• Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1

Chain F:  2% 70% 25%




• Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1


Chain H:  3% 67% 23% 7%




- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1




- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1



- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1




- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1




- Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1








● Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1



● Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1



● Molecule 2: PROBABLE EXOSOME COMPLEX EXONUCLEASE 1



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	206.20Å 214.00Å 432.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.25 – 3.30 93.25 – 3.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (93.25-3.30) 93.5 (93.25-3.30)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 3.33Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.274 , 0.295 0.265 , 0.283	Depositor DCC
R_{free} test set	4015 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	86.3	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.048 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	45814	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/1969	0.86	2/2678 (0.1%)
1	C	0.66	0/1972	0.87	2/2680 (0.1%)
1	E	0.59	0/1993	0.87	2/2706 (0.1%)
1	G	0.60	0/1986	0.86	2/2697 (0.1%)
1	I	0.57	0/1979	0.85	1/2688 (0.0%)
1	K	0.66	0/1983	0.87	0/2693
1	M	0.59	0/1985	0.84	3/2696 (0.1%)
1	O	0.67	0/1976	0.88	2/2684 (0.1%)
1	Q	0.61	0/1978	0.84	0/2688
1	S	0.60	0/1980	0.84	2/2690 (0.1%)
1	U	0.60	0/1933	0.86	0/2628
1	W	0.59	0/1975	0.85	2/2682 (0.1%)
2	B	0.57	0/1840	0.88	0/2494
2	D	0.64	0/1928	0.94	2/2606 (0.1%)
2	F	0.60	0/1866	0.89	0/2524
2	H	0.59	0/1757	0.88	0/2377
2	J	0.57	0/1872	0.85	0/2530
2	L	0.64	0/1918	0.94	2/2592 (0.1%)
2	N	0.59	0/1837	0.89	0/2487
2	P	0.64	0/1932	0.93	2/2609 (0.1%)
2	R	0.64	0/1908	0.94	5/2578 (0.2%)
2	T	0.62	0/1912	0.90	5/2584 (0.2%)
2	V	0.64	0/1843	0.88	0/2494
2	X	0.60	0/1848	0.89	0/2501
All	All	0.61	0/46170	0.88	34/62586 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	2
2	L	0	1
2	P	0	1
2	R	0	2
2	T	0	1
All	All	0	7

There are no bond length outliers.

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	9	ARG	CA-C-N	9.14	129.19	119.78
2	R	9	ARG	C-N-CA	9.14	129.19	119.78
2	D	18	LYS	N-CA-C	8.12	121.01	110.53
2	L	246	GLU	N-CA-C	7.08	117.94	108.38
2	D	246	GLU	N-CA-C	6.85	117.54	107.88

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	1	MET	Peptide
2	D	17	GLY	Peptide
2	L	17	GLY	Peptide
2	P	17	GLY	Peptide
2	R	5	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1944	0	1969	21	0
1	C	1947	0	1982	30	0
1	E	1968	0	2013	35	0
1	G	1961	0	2003	34	0
1	I	1954	0	1997	33	0
1	K	1958	0	2001	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	M	1960	0	2003	20	0
1	O	1951	0	1995	26	0
1	Q	1953	0	1988	23	0
1	S	1955	0	1996	30	0
1	U	1908	0	1950	26	0
1	W	1950	0	1997	26	0
2	B	1812	0	1810	43	0
2	D	1900	0	1920	52	0
2	F	1838	0	1852	49	0
2	H	1732	0	1740	51	0
2	J	1844	0	1870	49	0
2	L	1890	0	1915	52	0
2	N	1809	0	1817	43	0
2	P	1904	0	1938	51	0
2	R	1880	0	1904	44	0
2	T	1884	0	1913	52	0
2	V	1815	0	1836	42	0
2	X	1821	0	1820	36	0
3	B	27	0	12	4	0
3	D	27	0	12	4	0
3	F	17	0	7	1	0
3	H	27	0	12	2	0
3	J	9	0	0	1	0
3	L	27	0	12	3	0
3	N	27	0	12	3	0
3	P	27	0	12	3	0
3	R	27	0	12	1	0
3	T	27	0	12	3	0
3	V	17	0	7	2	0
3	X	17	0	7	1	0
All	All	45814	0	46346	823	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 823 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:R:99:ARG:NH1	3:R:404:ADP:O2A	1.85	1.09
2:T:1:MET:H2	2:T:2:ARG:HA	1.24	1.00
2:T:1:MET:N	2:T:2:ARG:HA	1.75	1.00

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:243:LYS:HG2	2:F:112:GLU:OE1	1.61	0.98
2:D:99:ARG:NH1	3:D:404:ADP:O2A	2.03	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/275 (92%)	241 (95%)	10 (4%)	3 (1%)	10	37
1	C	254/275 (92%)	240 (94%)	8 (3%)	6 (2%)	4	24
1	E	254/275 (92%)	241 (95%)	11 (4%)	2 (1%)	16	45
1	G	254/275 (92%)	239 (94%)	11 (4%)	4 (2%)	7	31
1	I	254/275 (92%)	242 (95%)	10 (4%)	2 (1%)	16	45
1	K	254/275 (92%)	240 (94%)	11 (4%)	3 (1%)	10	37
1	M	254/275 (92%)	240 (94%)	11 (4%)	3 (1%)	10	37
1	O	254/275 (92%)	245 (96%)	7 (3%)	2 (1%)	16	45
1	Q	254/275 (92%)	238 (94%)	13 (5%)	3 (1%)	10	37
1	S	253/275 (92%)	242 (96%)	9 (4%)	2 (1%)	16	45
1	U	249/275 (90%)	236 (95%)	11 (4%)	2 (1%)	16	45
1	W	253/275 (92%)	241 (95%)	9 (4%)	3 (1%)	10	37
2	B	239/248 (96%)	221 (92%)	15 (6%)	3 (1%)	9	35
2	D	246/248 (99%)	231 (94%)	10 (4%)	5 (2%)	6	27
2	F	239/248 (96%)	226 (95%)	10 (4%)	3 (1%)	9	35
2	H	226/248 (91%)	216 (96%)	8 (4%)	2 (1%)	14	43
2	J	239/248 (96%)	225 (94%)	10 (4%)	4 (2%)	7	30
2	L	245/248 (99%)	231 (94%)	13 (5%)	1 (0%)	30	60

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	N	237/248 (96%)	224 (94%)	12 (5%)	1 (0%)	30 60
2	P	246/248 (99%)	230 (94%)	13 (5%)	3 (1%)	10 37
2	R	246/248 (99%)	230 (94%)	13 (5%)	3 (1%)	10 37
2	T	246/248 (99%)	233 (95%)	11 (4%)	2 (1%)	16 45
2	V	237/248 (96%)	218 (92%)	15 (6%)	4 (2%)	7 30
2	X	235/248 (95%)	223 (95%)	11 (5%)	1 (0%)	30 60
All	All	5922/6276 (94%)	5593 (94%)	262 (4%)	67 (1%)	11 39

5 of 67 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	8	GLN
2	J	15	ASP
2	P	16	ASP
1	Q	8	GLN
2	V	91	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	212/242 (88%)	201 (95%)	11 (5%)	21 49
1	C	212/242 (88%)	201 (95%)	11 (5%)	21 49
1	E	218/242 (90%)	209 (96%)	9 (4%)	27 55
1	G	216/242 (89%)	207 (96%)	9 (4%)	26 55
1	I	214/242 (88%)	206 (96%)	8 (4%)	30 58
1	K	215/242 (89%)	203 (94%)	12 (6%)	19 47
1	M	216/242 (89%)	205 (95%)	11 (5%)	21 50
1	O	213/242 (88%)	204 (96%)	9 (4%)	26 55
1	Q	214/242 (88%)	204 (95%)	10 (5%)	23 52
1	S	216/242 (89%)	204 (94%)	12 (6%)	19 47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	U	208/242 (86%)	198 (95%)	10 (5%)	23	52
1	W	214/242 (88%)	203 (95%)	11 (5%)	21	50
2	B	186/208 (89%)	175 (94%)	11 (6%)	18	46
2	D	200/208 (96%)	185 (92%)	15 (8%)	12	38
2	F	193/208 (93%)	182 (94%)	11 (6%)	18	47
2	H	179/208 (86%)	168 (94%)	11 (6%)	17	45
2	J	194/208 (93%)	184 (95%)	10 (5%)	21	49
2	L	199/208 (96%)	186 (94%)	13 (6%)	15	43
2	N	188/208 (90%)	179 (95%)	9 (5%)	23	52
2	P	201/208 (97%)	184 (92%)	17 (8%)	10	33
2	R	194/208 (93%)	183 (94%)	11 (6%)	18	47
2	T	196/208 (94%)	180 (92%)	16 (8%)	10	35
2	V	189/208 (91%)	181 (96%)	8 (4%)	26	55
2	X	191/208 (92%)	180 (94%)	11 (6%)	18	47
All	All	4878/5400 (90%)	4612 (94%)	266 (6%)	19	48

5 of 266 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	U	191	LEU
2	V	99	ARG
2	X	99	ARG
2	J	20	THR
1	I	255	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	183	ASN
1	Q	83	ASN
2	V	216	GLN
2	N	178	ASN
1	O	183	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	T	404	-	28,29,29	1.43	6 (21%)	43,45,45	1.85	10 (23%)
3	ADP	V	404	-	16,17,29	1.22	2 (12%)	22,26,45	1.11	1 (4%)
3	ADP	F	404	-	16,17,29	1.86	3 (18%)	22,26,45	1.07	1 (4%)
3	ADP	X	404	-	16,17,29	2.09	3 (18%)	22,26,45	1.32	2 (9%)
3	ADP	B	404	-	28,29,29	1.62	5 (17%)	43,45,45	1.81	7 (16%)
3	ADP	L	404	-	28,29,29	1.51	3 (10%)	43,45,45	1.93	11 (25%)
3	ADP	H	404	-	28,29,29	1.47	4 (14%)	43,45,45	1.95	10 (23%)
3	ADP	N	404	-	28,29,29	1.57	5 (17%)	43,45,45	1.80	8 (18%)
3	ADP	D	404	-	28,29,29	1.51	4 (14%)	43,45,45	2.00	12 (27%)
3	ADP	R	404	-	28,29,29	1.63	5 (17%)	43,45,45	1.81	10 (23%)
3	ADP	P	404	-	28,29,29	1.49	5 (17%)	43,45,45	2.09	13 (30%)
3	ADP	J	404	-	6,8,29	0.94	0	12,13,45	1.05	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	T	404	-	-	3/16/32/32	0/3/3/3
3	ADP	V	404	-	-	0/12/25/32	0/1/1/3
3	ADP	F	404	-	-	2/12/25/32	0/1/1/3
3	ADP	X	404	-	-	6/12/25/32	0/1/1/3
3	ADP	B	404	-	-	6/16/32/32	0/3/3/3
3	ADP	L	404	-	-	3/16/32/32	0/3/3/3
3	ADP	H	404	-	-	3/16/32/32	0/3/3/3
3	ADP	N	404	-	-	5/16/32/32	0/3/3/3
3	ADP	D	404	-	-	1/16/32/32	0/3/3/3
3	ADP	R	404	-	-	3/16/32/32	0/3/3/3
3	ADP	P	404	-	-	5/16/32/32	0/3/3/3
3	ADP	J	404	-	-	0/6/6/32	-

The worst 5 of 45 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	404	ADP	O4'-C1'	6.32	1.57	1.43
3	F	404	ADP	O4'-C1'	5.47	1.55	1.43
3	B	404	ADP	C5-C4	5.29	1.48	1.39
3	R	404	ADP	C5-C4	5.23	1.48	1.39
3	N	404	ADP	C5-C4	5.12	1.48	1.39

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	404	ADP	C5-C4-N3	-6.15	118.25	126.72
3	H	404	ADP	C5-C4-N3	-6.02	118.42	126.72
3	N	404	ADP	C5-C4-N3	-5.98	118.49	126.72
3	L	404	ADP	C5-C4-N3	-5.77	118.77	126.72
3	R	404	ADP	C5-C4-N3	-5.68	118.89	126.72

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	404	ADP	C5'-O5'-PA-O2A
3	B	404	ADP	C5'-O5'-PA-O3A
3	H	404	ADP	C5'-O5'-PA-O3A

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	L	404	ADP	PB-O3A-PA-O5'
3	N	404	ADP	C5'-O5'-PA-O1A


There are no ring outliers.



12 monomers are involved in 28 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	T	404	ADP	3	0
3	V	404	ADP	2	0
3	F	404	ADP	1	0
3	X	404	ADP	1	0
3	B	404	ADP	4	0
3	L	404	ADP	3	0
3	H	404	ADP	2	0
3	N	404	ADP	3	0
3	D	404	ADP	4	0
3	R	404	ADP	1	0
3	P	404	ADP	3	0
3	J	404	ADP	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/275 (94%)	0.31	6 (2%) 61 42	46, 53, 67, 74	0
1	C	260/275 (94%)	0.30	4 (1%) 72 53	46, 53, 67, 75	0
1	E	260/275 (94%)	0.17	1 (0%) 88 79	46, 53, 67, 74	0
1	G	260/275 (94%)	0.45	7 (2%) 56 37	46, 53, 67, 74	0
1	I	260/275 (94%)	0.36	4 (1%) 72 53	46, 53, 67, 73	0
1	K	260/275 (94%)	0.42	8 (3%) 51 35	46, 53, 67, 75	0
1	M	260/275 (94%)	0.32	5 (1%) 66 48	46, 53, 67, 75	0
1	O	260/275 (94%)	0.48	6 (2%) 61 42	46, 53, 67, 75	0
1	Q	260/275 (94%)	0.21	1 (0%) 88 79	46, 53, 67, 74	0
1	S	259/275 (94%)	0.48	12 (4%) 37 25	46, 53, 67, 73	0
1	U	255/275 (92%)	0.80	24 (9%) 14 11	46, 53, 66, 73	0
1	W	259/275 (94%)	0.48	8 (3%) 51 35	46, 53, 67, 75	0
2	B	241/248 (97%)	0.20	1 (0%) 88 79	46, 52, 68, 80	0
2	D	248/248 (100%)	0.34	4 (1%) 70 52	45, 52, 68, 80	0
2	F	241/248 (97%)	0.19	4 (1%) 69 50	46, 52, 68, 80	0
2	H	230/248 (92%)	0.52	8 (3%) 47 31	46, 52, 69, 80	0
2	J	241/248 (97%)	0.40	7 (2%) 53 35	46, 52, 68, 80	0
2	L	247/248 (99%)	0.36	2 (0%) 82 68	45, 52, 68, 80	0
2	N	239/248 (96%)	0.26	4 (1%) 69 50	46, 52, 69, 79	0
2	P	248/248 (100%)	0.40	7 (2%) 55 36	45, 52, 68, 80	0
2	R	248/248 (100%)	0.30	4 (1%) 70 52	45, 52, 69, 80	0
2	T	248/248 (100%)	0.36	6 (2%) 59 40	45, 52, 69, 79	0
2	V	239/248 (96%)	0.69	12 (5%) 34 23	46, 52, 68, 79	0
2	X	239/248 (96%)	0.61	12 (5%) 34 23	46, 52, 69, 79	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	6022/6276 (95%)	0.39	157 (2%) 57 38	45, 52, 68, 80	0

The worst 5 of 157 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	166	TYR	4.1
2	L	248	VAL	3.7
1	O	134	GLY	3.7
1	C	12	PRO	3.7
2	X	172	VAL	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ADP	V	404	17/27	0.78	0.14	90,93,95,95	0
3	ADP	X	404	17/27	0.81	0.14	86,87,89,89	0
3	ADP	F	404	17/27	0.84	0.12	91,94,95,95	0
3	ADP	H	404	27/27	0.84	0.14	116,117,119,119	0
3	ADP	J	404	9/27	0.89	0.07	100,100,101,101	0
3	ADP	B	404	27/27	0.92	0.08	79,90,95,95	0
3	ADP	N	404	27/27	0.92	0.09	59,62,64,64	0
3	ADP	L	404	27/27	0.94	0.08	52,54,56,58	0
3	ADP	D	404	27/27	0.94	0.08	48,50,51,51	0
3	ADP	R	404	27/27	0.95	0.08	47,55,58,58	0
3	ADP	P	404	27/27	0.96	0.07	33,39,41,42	0
3	ADP	T	404	27/27	0.96	0.07	55,60,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ADP X 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP F 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP H 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP J 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP B 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP N 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around ADP L 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADP D 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP R 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around ADP P 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP T 404:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.