



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:48 AM UTC

PDB ID : 4C3P / pdb_00004c3p
Title : Structure of dephosphorylated Aurora A (122-403) bound to TPX2 and AMP-PCP
Authors : Zorba, A.; Kutter, S.; Kern, D.
Deposited on : 2013-08-26
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	
1	D	282	
2	B	43	
2	E	43	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4662 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called AURORA KINASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			2168	1392	384	386	6			
1	D	260	Total	C	N	O	S	0	0	0
			2136	1377	375	378	6			

- Molecule 2 is a protein called TARGETING PROTEIN FOR XKLP2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	17	Total	C	N	O	0	0	0
			134	86	18	30			
2	E	17	Total	C	N	O	0	0	0
			136	88	19	29			

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
3	D	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	13	Total	O	0	0
			13	13		
5	B	3	Total	O	0	0
			3	3		
5	D	5	Total	O	0	0
			5	5		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.93Å 86.72Å 153.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.48 – 2.69 47.48 – 2.69	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.48-2.69) 100.0 (47.48-2.69)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.201 , 0.289 0.204 , 0.285	Depositor DCC
R_{free} test set	1035 reflections (5.41%)	wwPDB-VP
Wilson B-factor (Å ²)	55.1	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4662	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/2222	1.03	7/3003 (0.2%)
1	D	0.75	0/2190	0.91	0/2959
2	B	0.89	0/138	0.87	0/187
2	E	0.78	0/140	0.85	0/189
All	All	0.78	0/4690	0.97	7/6338 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	PRO	CA-C-N	6.78	126.73	119.28
1	A	327	PRO	C-N-CA	6.78	126.73	119.28
1	A	273	ALA	N-CA-C	6.67	118.95	110.61
1	A	254	HIS	N-CA-C	-5.40	102.49	110.48
1	A	385	ALA	N-CA-C	5.08	117.48	111.33
1	A	126	ARG	N-CA-C	-5.06	103.33	111.02
1	A	332	ASN	N-CA-C	5.00	116.73	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2168	0	2178	36	0
1	D	2136	0	2147	15	0
2	B	134	0	113	3	0
2	E	136	0	119	1	0
3	A	31	0	14	1	0
3	D	31	0	14	0	0
4	D	5	0	0	0	0
5	A	13	0	0	0	0
5	B	3	0	0	0	0
5	D	5	0	0	0	0
All	All	4662	0	4585	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:TYR:CD1	1:A:270:LEU:HD13	2.35	0.61
1:D:290:CYS:SG	1:D:291:GLY:N	2.75	0.60
1:D:312:LEU:HD22	1:D:372:PRO:O	2.02	0.59
1:A:217:THR:HG22	1:A:263:LEU:CD2	2.32	0.59
1:A:333:THR:HG23	1:A:336:GLU:OE1	2.04	0.58
1:A:151:ARG:NH2	1:A:158:ILE:HD11	2.19	0.58
1:A:151:ARG:CZ	1:A:158:ILE:HD11	2.33	0.57
1:A:337:THR:HG21	1:D:293:LEU:HD13	1.87	0.56
1:A:300:MET:HE2	1:A:300:MET:HA	1.89	0.54
1:A:310:VAL:HG13	1:D:297:PRO:CG	2.39	0.53
1:A:275:PHE:HA	1:A:278:SER:OG	2.09	0.52
1:A:320:TYR:CG	1:A:328:PRO:HG3	2.44	0.52
1:A:178:LEU:HD23	2:B:10:TYR:OH	2.09	0.52
1:A:293:LEU:HA	1:A:296:LEU:HD11	1.92	0.52
1:A:253:ILE:HD11	1:A:281:ALA:HB2	1.93	0.51
1:A:296:LEU:HD13	1:A:301:ILE:HG12	1.92	0.51
1:A:297:PRO:HG3	1:D:310:VAL:HG23	1.94	0.50
1:A:285:ARG:HA	1:A:288:THR:HG22	1.93	0.50
1:A:310:VAL:HG13	1:D:297:PRO:HG2	1.94	0.50
1:A:240:LEU:HD23	1:A:272:ILE:HD11	1.96	0.48
1:A:373:MET:HE3	1:A:375:ARG:HG2	1.96	0.47
2:E:19:PHE:O	2:E:20:SER:CB	2.62	0.47
1:D:269:GLU:OE1	1:D:269:GLU:N	2.41	0.47
1:D:293:LEU:O	1:D:294:ASP:C	2.56	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:THR:O	1:A:237:ILE:HG12	2.16	0.46
1:A:296:LEU:HD13	1:A:301:ILE:CG1	2.46	0.46
1:A:343:ARG:HB2	1:A:345:GLU:HG3	1.98	0.46
1:A:254:HIS:O	1:A:311:ASP:OD1	2.34	0.45
1:D:253:ILE:HD13	1:D:307:ASP:HB3	1.98	0.45
1:A:338:TYR:CE2	1:D:301:ILE:CG2	3.00	0.45
1:A:341:ILE:HG21	1:D:301:ILE:CD1	2.47	0.44
1:A:289:LEU:HD13	1:A:304:ARG:HD2	2.00	0.44
1:A:341:ILE:HD13	1:D:301:ILE:HD11	2.00	0.44
1:D:240:LEU:HB3	1:D:315:LEU:HD11	1.98	0.44
1:A:375:ARG:CD	1:A:375:ARG:H	2.31	0.44
1:A:179:ARG:HG3	2:B:10:TYR:CD1	2.54	0.43
1:A:348:PHE:CD2	1:A:357:ARG:HG2	2.53	0.43
1:D:179:ARG:O	1:D:183:GLU:HG3	2.18	0.43
1:A:162:LYS:HB3	1:A:208:LEU:HB2	2.01	0.42
1:A:321:GLU:OE1	1:D:295:TYR:OH	2.35	0.42
1:A:373:MET:HE3	1:A:375:ARG:CG	2.49	0.42
1:A:217:THR:HG22	1:A:263:LEU:HD23	2.01	0.41
1:A:132:ASP:OD1	1:A:132:ASP:N	2.53	0.41
2:B:10:TYR:O	2:B:12:ALA:N	2.52	0.41
1:A:367:ASN:O	1:A:368:PRO:C	2.64	0.40
3:A:1389:ACP:H8	3:A:1389:ACP:O5'	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/282 (93%)	244 (93%)	17 (6%)	1 (0%)	30	54
1	D	256/282 (91%)	238 (93%)	17 (7%)	1 (0%)	30	54
2	B	15/43 (35%)	14 (93%)	1 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	15/43 (35%)	15 (100%)	0	0	100	100
All	All	548/650 (84%)	511 (93%)	35 (6%)	2 (0%)	30	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	255	ARG
1	A	372	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/250 (93%)	211 (91%)	22 (9%)	8	21
1	D	229/250 (92%)	211 (92%)	18 (8%)	11	28
2	B	16/40 (40%)	14 (88%)	2 (12%)	4	11
2	E	16/40 (40%)	16 (100%)	0	100	100
All	All	494/580 (85%)	452 (92%)	42 (8%)	10	25

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	LYS
1	A	126	ARG
1	A	159	LEU
1	A	179	ARG
1	A	188	LEU
1	A	226	SER
1	A	232	ARG
1	A	240	LEU
1	A	249	SER
1	A	257	ILE
1	A	258	LYS
1	A	280	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	296	LEU
1	A	299	GLU
1	A	300	MET
1	A	305	MET
1	A	312	LEU
1	A	315	LEU
1	A	330	GLU
1	A	340	ARG
1	A	354	GLU
1	A	375	ARG
2	B	14	SER
2	B	15	ASP
1	D	125	LYS
1	D	127	GLN
1	D	164	LEU
1	D	188	LEU
1	D	194	LEU
1	D	195	ARG
1	D	220	ARG
1	D	226	SER
1	D	255	ARG
1	D	257	ILE
1	D	289	LEU
1	D	292	THR
1	D	293	LEU
1	D	305	MET
1	D	315	LEU
1	D	318	LEU
1	D	354	GLU
1	D	362	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	306	HIS
1	A	370	GLN
1	D	127	GLN
2	E	18	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACP	A	1389	-	31,33,33	1.99	10 (32%)	47,52,52	1.84	11 (23%)
4	SO4	D	1392	-	4,4,4	0.55	0	6,6,6	0.34	0
3	ACP	D	1391	-	31,33,33	2.07	10 (32%)	47,52,52	1.79	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACP	A	1389	-	-	3/19/38/38	0/3/3/3
3	ACP	D	1391	-	-	6/19/38/38	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1389	ACP	PG-O1G	6.28	1.62	1.50
3	D	1391	ACP	PG-O1G	5.79	1.61	1.50
3	D	1391	ACP	C5-C4	4.93	1.47	1.39
3	A	1389	ACP	C5-C4	3.87	1.46	1.39
3	A	1389	ACP	PG-O2G	3.50	1.62	1.55
3	D	1391	ACP	PB-O3A	3.31	1.62	1.58
3	A	1389	ACP	C5-N7	-3.23	1.33	1.39
3	D	1391	ACP	PG-O2G	3.06	1.61	1.55
3	D	1391	ACP	PB-O2B	2.68	1.63	1.56
3	A	1389	ACP	PG-O3G	-2.68	1.49	1.55
3	D	1391	ACP	C8-N7	2.67	1.36	1.31
3	D	1391	ACP	PG-O3G	-2.59	1.49	1.55
3	A	1389	ACP	PB-O3A	2.50	1.61	1.58
3	D	1391	ACP	C5-N7	-2.49	1.34	1.39
3	D	1391	ACP	PA-O3A	2.44	1.62	1.59
3	D	1391	ACP	C5-C6	2.40	1.47	1.41
3	A	1389	ACP	C4-N9	-2.23	1.33	1.37
3	A	1389	ACP	PB-O2B	2.15	1.61	1.56
3	A	1389	ACP	C8-N9	-2.12	1.33	1.37
3	A	1389	ACP	C8-N7	2.05	1.35	1.31

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1389	ACP	C5-C4-N3	-5.52	119.12	126.72
3	D	1391	ACP	C5-C4-N3	-5.32	119.39	126.72
3	A	1389	ACP	N3-C4-N9	4.43	134.70	127.17
3	D	1391	ACP	N3-C2-N1	-4.17	122.28	128.58
3	D	1391	ACP	N3-C4-N9	3.95	133.88	127.17
3	A	1389	ACP	PB-O3A-PA	-3.76	120.11	132.37
3	D	1391	ACP	C2-N3-C4	3.75	121.00	111.83
3	A	1389	ACP	C2-N3-C4	3.57	120.54	111.83
3	A	1389	ACP	N3-C2-N1	-3.55	123.20	128.58
3	D	1391	ACP	C4-C5-N7	-3.53	106.55	110.58
3	A	1389	ACP	C4-C5-N7	-3.16	106.97	110.58
3	D	1391	ACP	PB-O3A-PA	-2.83	123.14	132.37
3	A	1389	ACP	O2G-PG-C3B	2.80	113.18	106.40
3	A	1389	ACP	C4-N9-C8	2.65	108.52	105.74
3	D	1391	ACP	C2-N1-C6	2.61	123.01	118.73
3	A	1389	ACP	C5-N7-C8	2.54	107.45	103.45
3	D	1391	ACP	C5-N7-C8	2.52	107.41	103.45
3	D	1391	ACP	C6-C5-N7	2.39	136.70	132.09
3	D	1391	ACP	C4-N9-C8	2.17	108.02	105.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1389	ACP	N9-C8-N7	-2.15	110.88	113.94
3	A	1389	ACP	C6-C5-N7	2.13	136.19	132.09

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1389	ACP	PB-C3B-PG-O1G
3	A	1389	ACP	PB-C3B-PG-O2G
3	A	1389	ACP	PB-C3B-PG-O3G
3	D	1391	ACP	C5'-O5'-PA-O1A
3	D	1391	ACP	C5'-O5'-PA-O2A
3	D	1391	ACP	C5'-O5'-PA-O3A
3	D	1391	ACP	O4'-C4'-C5'-O5'
3	D	1391	ACP	C3'-C4'-C5'-O5'
3	D	1391	ACP	PB-C3B-PG-O1G

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1389	ACP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/282 (93%)	0.07	7 (2%) 56 53	29, 49, 90, 123	0
1	D	260/282 (92%)	-0.02	7 (2%) 56 53	32, 51, 91, 128	0
2	B	17/43 (39%)	0.44	1 (5%) 28 25	33, 51, 92, 92	0
2	E	17/43 (39%)	0.16	1 (5%) 28 25	39, 53, 93, 99	0
All	All	558/650 (85%)	0.04	16 (2%) 53 50	29, 50, 91, 128	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	289	LEU	5.6
1	D	305	MET	3.7
2	B	22	LEU	3.7
1	A	282	PRO	3.3
1	A	291	GLY	3.0
1	D	390	PRO	2.8
1	D	282	PRO	2.6
1	A	288	THR	2.5
1	D	300	MET	2.4
1	A	256	ASP	2.4
2	E	4	VAL	2.2
1	A	281	ALA	2.2
1	A	125	LYS	2.2
1	A	222	LEU	2.1
1	D	256	ASP	2.0
1	D	266	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	D	1392	5/5	0.83	0.14	90,94,106,107	0
3	ACP	D	1391	31/31	0.90	0.08	40,54,92,95	0
3	ACP	A	1389	31/31	0.92	0.08	35,52,82,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.